

SEVIN BAIT

FOR THE CONTROL OF INSECT PESTS:
Cutworms, slugs, snails, armyworms, field crickets, darkling ground beetles, grasshoppers, sowbugs and ants.

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl-N-methylcarbamate) 5.00%

INERT INGREDIENTS: 95.00%

TOTAL 100.00%

Sevin® is the registered trademark of Union Carbide Corporation for the active ingredient carbaryl-(1-naphthyl-N-methylcarbamate).

KEEP OUT OF REACH OF CHILDREN
CAUTION

See additional cautions and warnings below.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if inhaled or swallowed. Avoid breathing of dust. Do not get in eyes. Avoid contact with skin. Wear regular long-sleeved work clothing when applying. Do not eat, drink, or smoke while applying. Wash hands and face before eating. Wash thoroughly after handling. Avoid contamination of food and feed. Keep pets off treated areas.

STATEMENTS OF PRACTICAL TREATMENT:

If in eyes: Flush with plenty of water at least 15 minutes. Get medical attention.
If swallowed: Induce vomiting by drinking 1 or 2 glasses of water and touching back of throat with finger. Get a physician or Poison Control Center immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidote. Do not use 2-PAM, apocain or cholinesterase inhibiting drugs.

If on skin: Wash thoroughly with soap and water.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic organisms. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues. The product may kill bees in substantial numbers. Do not use when value of bees as pollinators is more important than insect control. Before applying warn beekeepers to locate hives beyond bee flight range until one week after application or to take other equally effective precautions.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage, cleaning of equipment, or disposal. Open dumping is prohibited.

Pesticide or rinsate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

Dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place.

Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

Important Pre-Harvest and Grazing Use Information: Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label directions.

If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat.

Amount to Use: Broadcast applications may be made with either ground equipment or aircraft. Treatments may be repeated as necessary.

Dosages from 20-40 pounds per acre of Bait will control cutworms, slugs, snails, armyworms, field crickets, darkling ground beetles, grasshoppers, sowbugs and ants.

Crop	Dosage Limitations to Meet Residue Tolerances	Days Between Final Application and Harvest
Cucumbers, squash, melons, peaches	20 pounds per acre (7.3 oz./1,000 sq. ft.)	None
Alfalfa, peas, cotton	30 pounds per acre (11 oz./1,000 sq. ft.)	None
Beans, carrots, corn, eggplants, lima, peppers, potatoes, tomatoes	40 pounds per acre (14.7 oz./1,000 sq. ft.)	None
Asparagus, strawberries	40 pounds per acre (14.7 oz./1,000 sq. ft.)	3 Days
Broccoli, Brussels sprouts, cabbage, cauliflower, head lettuce, garden beets (roots), horseradish, parsnips, radishes, rutabagas, and turnips (roots)	40 pounds per acre (14.7 oz./1,000 sq. ft.)	14 Days
Sugar beets, callards, endive (escarole), garden beets (tops), kale, leaf lettuce, parsley, spinach, Swiss chard and turnips (tops), Dandelions	40 pounds per acre (14.7 oz./1,000 sq. ft.)	None
Tobacco — Field — Bud	20-40 pounds/acre 1/2-1 pound/1,000 sq.	None
(All insects listed above except darkling ground beetles) for the following crops: Apples, apricots, cherries, citrus, grapes, nectarines, peaches, pears, plums, and annual plants — herbaceous plants and shrubs, trees and woody ornamental plants	40-48 pounds (14.7-17.6 oz./1,000 sq. ft.)	None

*Includes Field, Sweet & Popcorn (including N-Till corn)

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given hereon.

NOTE: This pesticide is to be sold ONLY in the original unbroken package.

EPA Reg No. 8590-510

EPA Est No. 8590 PA 1



AGWAY INC., CHEMICAL DIVISION
PO BOX 4933, SYRACUSE, NY 13221

Net Weight 50 lbs.

A 3/82

BEST AVAILABLE COPY