AGWAY
GREENLAWN PLUS 21-5-7 Fertilizer
with Broadleaf Weed Killer Formula III

Contains 2,4-D and MCPP
Weeds and Feeds 465 sq. meters (5,000 sq. ft.)
Kills dandelions, clover, plantain, chickweed, and many other Broadleaf Weeds as listed.
Feeds and builds beautiful, thick green lawns.
25% of the nitrogen is in a slow release form.

(((picture)))

Ingredient statement as a Herbicide

ACTIVE INGREDIENTS:

Total 100.00%

(1) Equivalent to

2,4-dichlorophenoxyacetic acid

(2) Equivalent to

2,(2-methy1-4chlorophenoxy)-propionic acid

* ISOMER SPECIFIC BY AOAC METHOD NO.6. DO1-5

EPA Reg. No. 8590-522 EPA Est. No. 8590-NY-1

KEEP OUT OF REACH OF CHILDREN (((12 pt.)))

CAUTION (((18 pt.)))

See additional precautionary statements on rear panel

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS. AND

DIRECTIONS.

Net Wt. 18.16 kg (40 lbs.) Covers 5,000 sq. ft. (465 sq. meters)

> CODE 86-5416-31 (1 of 3) 9/10/84 A0067-142

Weeds and feeds 465 square meters (5,000 square feet) Net Wt. 18.16 kg (40 lbs.)

**FERTILIZER - GUARANTEED ANALYSIS:

Total Nitrogen (N)	10.00%
1.75 Coated Slow Release Ures Nitrogen	
8.25 Water Soluble Nitrogen	
Available Phosphoric Acid (P,O,)	6.00%
Available Phosphoric Acid (P ₂ O ₅)	4.00%
Z	

(((LOGO)))

AGWAY INC., BOX 4741, SYRACUSE, NY 13221

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Keep bag closed when not in use - store in a dry place.
KEEP THIS FERTILIZER-PESTICIDE MIXTURE IN ORIGINAL CONTAINER.

STORAGE & DISPOSAL: (((12 pt))) Do not contaminate water, food, or feed by storage or disposal.

Store only in original container in a cool, dry area that can be locked and is out of the reach of children.

Do not reuse empty bag. Discard bag in trash.

rejuvinates under-

nourished and weedy lawns by:

- 1. Providing a controlled feeding fertilizer which gives a green lawn, and
- 2. Controlling unsightly broadleaf weeds such as:

aster
buckhorn
buttercress
black medic
burdock
buttercup
buttonweed
chickweed
chicory
carpet weed
cinquefoil
cudweed
curly dock
daisy fleabane
dandelion
english daisy
evening primrose

field bindweed filaree galinsoga ground ivy beal all henbit hop clover knotweed (seedling) lambsquarters matchweed creeping charlie (moneywort) morning glory mustard oxalis pennywort pepper weed

pigweed pineapple weed plantain prickly lettuce purslane ragweed shadowf.lower shepard s purse sow thistle smartweed stitchwort. thyme leaf. (sandwort) vetch white clover wild radish yellow rocket

> CODE 86~5416-31 (2 of 3) 9/10/84 A0067-146

September 1

young and growing vigorously. Do not use this product during hot summer months since injury may occur. To avoid injury to new growth, new lawns should not be treated until after the fourth mowing. Established lawns may be injured if treated when in need of water. Water thoroughly 3 to 4 days prior to treatment and again the day before treatment. Grass may be injured if treated in periods of high heat and humidity. For this reason, apply only when the daytime temperature is not expected to exceed 80°F for 48 hours. To a oid drift, apply when air is calm. Do not apply if rain is expected within 24 hours.

HOW TO APPLY: The contents of this bag will cover an area of 5,000 sq. ft. Apply at rate of 3.90 kg/100 square meters (8.0 lbs./1,000 sq. ft.) Apply to lawn when grass is wet (after rain, sprinkling, or dew) so pellets will stick to the weeds. Do not water or mow lawn for at least 2 days after treatment, then resume normal watering program. Supplement with additional feeding of Turfood Special 10-6-4 Fertilizer as recommended for year-round nutrition of the lawn.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Avoid contact with skin, eyes, and clothing. Avoid breathing dust. Wash thoroughly after handling. In case of eye contact, flush with water for 15 minutes. Get medical attention if irritation persists. In skin contact, wash with soap and water.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not apply when weather conditions favor drift from target area.

IMPORTANT NOTICE: Since this product kills desirable as well as undesirable broadleaf plants, it may injure flowers, ornamentals, and trees. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not exceed recommended rate. Avoid contact with flower beds or shrubs. If accidentally deposited on flowers or shrubs, hose off immediately. Do not use on Bentgrass, Dichondra, Carpetgrass, St. Augustine, or ground covers. Delay planting grass seed until 4 weeks after treatment. Read and follow directions carefully.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

NOT: This pesticide is to be sold ONLY in this original unbroken package.

SPREADER SETTING GUIDES **

		Cyclone Bl		0rt	tho	Sco	otts Ort		tho	So	Scotts.	
Rate P	er	Spinne	r Type	Spinne	er Type	Spinne	r Type	Drop	Type	Drop	·Type	
100 Sq.	1000	Once	Twice	Once	Twice	Once	Twice	Once	Twitte			
Meters	Sq. Ft.	Over	Over	Over	Over	Over	Over	Over	Over .	Over '	Over	
3.90 kg	8.0 lbs.	!		1		Į		-		٠.		

**NOTE:

These settings are furnished as a guide only. Age, condition of spreader, speed of operation, and roughness of terrain can cause variation in rates applied.



CODE 86-5416-31 (3 of 3) 9/10/84 A0067-147



COMPANY / O/48-NUMBER

PRODUCT SERIAL NO.

LABEL APPROVAL 6-/3-85 DATE