



# AGWAY 3% RABON DUST

A Ready To Use Insecticide for direct application to DAIRY & BEEF CATTLE and HORSES.

Controls Horn flies, stable flies, horse flies, deer flies, house flies, fly larvae, lice, gnats and ticks. Hand Dusting. Apply approximately 2 oz. of dust by shaker can, rotary duster or by spray to the upper portions of the back, neck and poll, and to the face area in aid in the control of face flies. Rub in lightly to carry dust beneath the hair. Repeat as necessary.

Also controls ticks.

ACTIVE INGREDIENTS  
2-Chloro-1-(2,4,5-trichlorophenyl)vinyl dimethyl phosphorothioate . . . . . 3.0%

INERT INGREDIENTS . . . . . 97.0%

TOTAL . . . . . 100.0%

BY WEIGHT

\*Rabon is the registered trademark of Shell Chemical Company.

## CAUTION:

**KEEP OUT OF REACH OF CHILDREN.**  
See additional cautions on side panel

## DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

## STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL** Pesticide spray mixture or rinseate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

**CONTAINER DISPOSAL** (1# canister) (3# bag) Dispose of canister (bag) according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

## DAIRY AND BEEF CATTLE

3# RABON DUST is a ready to use insecticide dust which can be applied by hand or through the use of self treating dust bags. There is no withholding period from last application to slaughter.

For horn flies, stable flies, horse flies, deer flies, house flies, fly larvae, lice, gnats and ticks. Hand Dusting. Apply approximately 2 oz. of dust by shaker can, rotary duster or by spray to the upper portions of the back, neck and poll, and to the face area in aid in the control of face flies. Rub in lightly to carry dust beneath the hair. Repeat as necessary.

## SELF TREATING DUST BAG

**FORCED USE** P. 1-3 RABON DUST in cotton cloth or

double burlap bags or use pre-packed weatherproof cattle dust bags and hang in barn door exits or alleyways leading from animal buildings, salt or mineral blocks, or watering holes. Protect cloth or burlap bags from weather.

**FREE CHOICE USE** Use the same dust bags as above but place in loafing sheds, holding pens, feedlots, near watering holes, or other areas where cattle gather. The free choice use aids in the control of lice.

## HORSES

For use against most common insect pests of horses including:

- Horn Flies • Gnats • Horse Flies • Face Flies • Ticks
- Stable Flies • Lice • Deer Flies • House Flies • Fly Larvae

To help control the above insects, apply approximately 2 oz. by shaker can, hand duster, grooming brush or dust mitt. As with any pest control materials, do not use bare hands. Cover upper portions of back and neck. Apply to the face, mane and tail for added control of face flies. Rub or brush in lightly to carry dust beneath the hair. Repeat as necessary.

**PREMISE DUSTING** Once a week apply at the rate of 1 lb. per 100 sq. ft. (an area 10 ft. by 10 ft.) to barn or stall area floors where manure accumulates, to control fly larvae and flies in the stable area.

**Preventative Maintenance** Since flies breed in decaying organic matter, moisture, dirt and manure, it's important to keep stable and paddocks free of these materials. Stalls should be mucked out daily and low, wet spots either drained or filled in. Pay particular attention to fences and gates where low spots often go unnoticed, and water for potholes in pasture and fence corners. Clean up spoiled feeds immediately. And, of course, sprinkle the premises as required to further control flies at the source.

## SWINE

For lice. Hand Dusting. Apply 1-4 oz. of dust by conventional hand or power duster to each animal with special attention given to the neck and around the ears. Repeat as necessary but not more often than once every 14 days. In severe infestations, both individual animals and a bedding may be treated. One lb. of dust should be applied per 150 sq. ft. of bedding.

## PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS & DOMESTIC ANIMALS**  
**CAUTION** Harmful if swallowed. Avoid breathing dust. Do not get in eyes, on skin or on clothing. Wear protective glove or mitt when hand dusting. Avoid contamination of feed and foodstuffs. Do not apply in dwellings.

Do not apply to animals less than one month old. Do not contaminate milk or milking equipment.

## STATEMENT OF PRACTICAL TREATMENT

**If swallowed** Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Repeat until fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

**If inhaled** Remove victim to fresh air. Apply artificial respiration if indicated.

**If on skin** Remove contaminated clothing and wash affected areas with soap and water.

**If in eyes** Flush eyes with plenty of water. Call a physician if irritation persists.

## ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, ponds, streams or public waters. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DISCLAIMER** Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

**NOTE** This product is to be sold ONLY in its original, unbroken package.

EPA Reg. No. 8590-524 EPA Est. No. 5590-PA-1



AGWAY INC., CHEMICAL DIVISION  
BOX 4933, SYRACUSE, NY 13221

Net Weight 1 lb.