

and tobacco moths. Direct spray at an upward angle distributing uniformly throughout the entire area at a rate of 3.6 ounces per 1000 cubic feet of space.

FOGGING: To kill exposed or accessible stages of the above named crawling insects, use a rate of one gallon to 20,000 to 40,000 cubic feet of space. Contact insects directly whenever possible.

ON ANIMALS: To protect cattle and horses from horn flies, house flies, mosquitoes and gnats, apply a light mist sufficient to wet the surface of the hair. To control stable flies, horse flies and deer flies, apply at a rate of 2 ounces per adult cow or horse, sufficient to wet the hair thoroughly. Repeat treatment daily or at intervals necessary to give continued protection.

To control blood-sucking lice, apply to the infested areas of the animal, using a stiff brush to get the spray to the base of the hair. Repeat every 2 to 3 weeks if required.

OUTDOOR ADULT MOSQUITO CONTROL: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest. Low marshy places, stagnant pools or ponds and shorelines. Apply at a rate of 1.5 gallons per acre (4.5 ounces per 1000 square feet).

Using truck-mounted conventional misting equipment such as Rotomist models 51C, 91 or 100 HT traveling at 5 mph to treat a 300-foot swath, calibrate equipment to deliver 4.5 gallons per minute.

PRECAUTIONARY STATEMENTS

CAUTION: Hazardous to Humans & Domestic Animals. Harmful if swallowed. Avoid contamination of feed and foodstuffs.

ACCEPTED

01/2/01

874 533

STATEMENT OF PRACTICAL TREATMENT

If swallowed: drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If in eyes: wash with plenty of water for at least 15 minutes. If irritation persists, get medical attention.

If on skin: wash with soap and water. Remove any contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Remove pets, birds and cover fish aquaria before spraying.

This product is toxic to fish. Keep out of lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

EPA Reg. No. 8590-533

EPA Est. No. 8590-PA-1



AGWAY INC., CHEMICAL DIVISION
BOX 4933, SYRACUSE, NY 13221

Net Contents 1 gal.

AGWAY GENERAL PURPOSE AQUEOUS INSECTICIDE

A Concentrated Micro Emulsion - Clear
- Stable
- Leaves No Oily or Objectionable Residues

For Use in Homes, Institutions, Restaurants, Food Processing Plants, Livestock and Outdoors for Mosquito Control.

Apply with Conventional Mechanical or Compressed Air Equipment.

ACTIVE INGREDIENTS

Pyrethrins
*Piperonyl Butoxide, Technical
Petroleum Distillate

INERT INGREDIENTS

TOTAL

100% active ingredients are in the form of pyrethrins, ether and D-24 related compounds.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

See side panel for additional cautions.

AGWAY GENERAL PURPOSE AQUEOUS INSECTICIDE

A Concentrated Micro Emulsion — Clear
— Stable
— Leaves No Oily or Objectionable Residues

For Use in Homes, Institutions, Restaurants, Food Processing Plants, Dairies,
Livestock and Outdoors for Mosquito Control

Apply with Conventional Mechanical or Compressed Air Equipment

ACTIVE INGREDIENTS

Pyrethrins	0.1%
*Piperonyl Butoxide Technical	1.0%
Petroleum Distillate	0.4%
INERT INGREDIENTS	98.5%
TOTAL	100.0%

*Equivalent to 0.6% of total active ingredients.

**KEEP OUT OF REACH OF CHILDREN.
CAUTION**

See side panel for additional cautions

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection. Do not use near open food areas.

Indoor Application — The use of this product in food processing establishments should be confined to time periods when the plant is not in operation. Food should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using.

Vacate treated areas and ventilate before reoccupying.

For maximum effectiveness, a combination of spot surface treatment and fogging is recommended.

SURFACE SPRAYING To control roaches, ants, silverfish, spiders, crickets, clover mites, cheese mites, granary weevils, rice weevils, confused flour beetles, rust red flour beetles, saw-toothed grain beetles, spider beetles, cigarette beetles, Drug-store beetles, meal worms, grain mites and cadellies. Use a good sprayer adjusted to a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application is desired, spray floors and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To control carpet beetles: Spray edges of carpeting and under carpeting and rugs. Make localized application to floor and baseboards. Spray directly into cracks, closets and infested areas of shelving. Repeat treatment as needed.

To control bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, mouldings and floors. Repeat treatment as needed.

To control fleas and brown dog ticks: Thoroughly spray infested areas per beds, resting quarters, nearby cracks and crevices, along and behind baseboards, moldings, window and door frames, and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To control clothes moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight, if possible. Apply spray liberally to empty chests, closets, bureaus and other storages directing the spray into cracks, joints and crevices. After airing infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining.

SPACE SPRAYING Close doors and windows and shut off ventilating systems. Apply with a good machine adjusted to deliver a fine mist or fog. For rapid control of flies, fruit flies, gnats, mosquitoes, skipper flies, wasps, hornets, Angoumois grain moths, Mediterranean flour moths, Indian meal moths.