

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY CAPTAN 80WP

DATE: 9/9/87

((FRONT PANEL))

NET WEIGHT
5#, 30#

PICA - 50 CHARACTERS
ELITE - 60 CHARACTERS

AGWAY
CAPTAN 80WP

Fungicide for disease control of fruits, vegetables & ornamentals

ACTIVE INGREDIENTS:

*CAPTAN:	78.3%
Related Derivative	1.7%
INERT INGREDIENT	20.0%
TOTAL	100.0%

ACCEPTED
SEP 30 1987
 Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act, as amended, for the pesticide registered under EPA Reg. No. 8590-553

*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

KEEP OUT OF REACH OF CHILDREN (12 pt.)
DANGER (18 pt) PELIGRO (18 pt)

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a physician.

If swallowed: Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol.

If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

PRECAUCION AL USUARIO: Si Usted No lee ingles, No use este producto hasta que le etiqueta haya sido explicado ampliamente

BEST AVAILABLE COPY

O'K BY

2016

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY CAPTAN 80WP

DATE: 9/9/87

See additional precautionary statements below.

AGWAY INC.
P.O. BOX 4741
SYRACUSE, NEW YORK 13221

EPA Reg. No. 8590-553

EPA Est. No. 8590-PA-1

PICA 50 CHARACTERS
ELITE 60 CHARACTERS

OK BY	
.....
.....
.....
.....
.....
.....

FOR: AGWAY CAPTAN BOWP

DATE: 9/9/87

((SIDE PANEL))

PICA - 50 CHARACTERS
ELITE - 60 CHARACTERS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin & clothing. Remove & separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

This Pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Drift & runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

OK BY

ADVERTISING AND SALES PROMOTION SERVICES

40616
COPY

FOR: AGWAY CAPTAN 80WP

DATE: 9/9/87

((REAR & SIDE PANEL))

PICA 50 CHARACTERS
ELITE 60 CHARACTERS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse.

Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

Fieldworkers and harvesters must wear chemical resistant gloves. Leather or fabric gloves are not acceptable.

A dust mask must be worn when mixing and loading this product.


AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

During application only, from a tractor with a completely enclosed cab or from an aircraft with an enclosed cockpit a long sleeved shirt and long legged pants must be worn. Chemically resistant gloves must be available in the cab or cockpit and worn while exiting.

RE-ENTRY

Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.



ADVERTISING AND SALES PROMOTION SERVICES

20/10
COPY

FOR: AGWAY CAPTAN 80WP

DATE: 9/9/87

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

DANGER

(Insert area or field description) treated with Captan on (insert data of application). Do not enter treated areas within 4 days following application unless protective clothing is worn. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Captan and describe his condition. For further information see the "Statement of Practical Treatment".

Oral warnings must include the same information that is in the written warnings.

CAPTAN 80WP is a microfine wettable powder for use in water as a spray for the control of certain fungus diseases of fruit, vegetables, and ornamental crops, and as a soil treatment for the control of certain seed rots and damping-off diseases.

PICA 50 CHARACTERS
ELITE 50 CHARACTERS

OK BY

SHEET NO.

4-15

ADVERTISING AND SALES PROMOTION SERVICES

60816
COPY

FOR: AGWAY CAPTAN BOWP

DATE: 9/9/87

OBSERVE THE FOLLOWING USE PRECAUTIONS


Read all precautions and directions for use before using. Use only for claims listed and only as specified on this label.

In order that pesticide residues on food and forage crops will not exceed federal tolerances, use only at recommended rates and intervals, and do not apply closer to harvest than specified.

Do not apply or allow to drift to adjoining food fiber or pasture crops.

Consult State Agricultural Experiment Stations or State Agricultural Extension Service for additional information, as the timing of applications needed will vary with the local conditions.

PICA - 50 CHARACTERS
ELITE - 60 CHARACTERS



ADVERTISING AND SALES PROMOTION SERVICES

7 of 16
COPY

FOR: AGWAY CAPTAN 80WP

DATE: 9/9/87

COMPATIBILITY AND PLANT SAFETY: CAPTAN 80WP

can be combined safely and effectively at recommended dosage rates with most commonly used fungicides and insecticides, with the exception of oil and strongly alkaline materials. Alkaline materials such as spray lime, lime-sulfur and bordeaux mixture will reduce the fungicidal activity of CAPTAN. Do not apply CAPTAN in combination with or immediately before or closely following oil sprays. The time factor governing the safe interval between CAPTAN and oil sprays varies due to general climatic conditions, therefore, consult local agricultural spray programs and authorities to determine the proper timing. The use of spreaders which cause excessive wetting is not advised. Combinations with solvent formulations of organic phosphates should not be used. Combinations of CAPTAN and sulfur should not be used on crops sensitive to sulfur. Used at high rates or in drenching sprays, CAPTAN may cause a necrotic spotting of tender, immature leaves of certain varieties of apples, peaches, plums and cherries.

PICA - 50 CHARACTERS
ELITE - 60 CHARACTERS

O'K BY

SHEET NO.

FOR: AGWAY CAPTAN 80WP

DATE: 9/9/87


This type of injury is most likely to occur in the early cover sprays during long periods of warm, cloudy, humid weather. To avoid the hazard of leaf spotting under such conditions, use CAPTAN and other spray materials at lowest recommended rates and avoid drenching trees. Applications can be made by aircraft or ground power equipment (including concentrate and semi-concentrate equipment). Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants. (Read compatibility and plant safety information given under USE PRECAUTIONS.)

RECOMMENDATIONS

Unless otherwise specified, rates are given in terms of CAPTAN 80WP per 100 gallons water for fruit and nut crops and ornamentals. For vegetable crops, rates are expressed as CAPTAN 80WP per acre in sufficient water to provide thorough coverage. Except as specified begin applications before or at first sign of disease and repeat as needed to maintain control but observe use limitations. Unless otherwise specified, application can be made on the day of harvest.

PICA 50 CHARACTERS
ELITE - 60 CHARACTERS

O'KBY



ADVERTISING AND SALES PROMOTION SERVICES

90616
COPY

FOR: AGWAY CAPTAN 80WP

DATE: 9/9/87

Maximum permissible rate per acre, per application, expressed as CAPTAN 80WP is given in parentheses (), after each crop claim.

FRUITS

APPLES (eastern U.S.): Primary scab, black rot (frog-eye), Botrytis blossom end rot - 1 to 1½ pounds at 5 to 7 day intervals as needed to maintain cover in prebloom, bloom, petal fall and first cover sprays. Use at 1½ pounds rate if conditions favor scab or if active scab can be found. Secondary scab, Brooks fruit spot, sooty blotch, fly speck, black rot, black pox, Botryosphaeria rot, bitter rot - ¾ to 1½ pounds at 10 to 14 day intervals in second and later cover sprays up to start of harvest. (10 pounds). Powdery mildew - if powdery mildew is a problem add 1½ to 3 pounds sulfur or ¼ to ½ pound karathane to all pre-bloom and bloom sprays and ¼ to ½ pound karathane to all post bloom sprays until the foliage matures. Use ½ pound karathane if interval between sprays is longer than 7 days. Special Sooty Blotch Sprays (August and September) 10 oz. plus 1 lb. Zineb Wettable Powder Fungicide (such as FMC's Zineb 75 Wettable Powder or Rohm & Haas' Dithane Z-78). Where lead arsenate has been used in the apple maggot cover sprays, use at 1½ lbs. per 100 gals. spray. Do not use lead arsenate within 30 days of harvest.

PICA 50 CHARACTERS
ELITE - 60 CHARACTERS

OK BY

SHEET NO.

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY CAPTAN BOWP

DATE: 9/9/87

Caution - Do not use CAPTAN in combination with or closely following or in alternation with wettable sulfur products on sulfur sensitive varieties of apples such as Red Delicious, Stayman, Baldwin, King, etc. as severe injury and defoliation may occur.


PICA - 50 CHARACTERS
ELITE - 60 CHARACTERS

APRICOTS: Brown rot (twig blight), jacket rot - 1 1/2 pounds. Apply in red bud, bloom, 75% petal fall, and cover stages. (6 1/2 pounds)

CHERRIES (eastern U.S.): Brown rot, leaf spot, Botrytis rot - Use 1 1/2 pounds in pre-bloom, bloom, petal fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control blossom blight. Repeat applications at 7 to 10 day intervals as needed to maintain cover up to start of harvest. If powdery mildew is a problem add 3 pounds sulfur to the petal fall, shuck and early cover sprays.

If sulfur is added, CAPTAN may be reduced to 3/4 pound in these sprays. Post harvest sprays. Use 1 1/2 pounds immediately after harvest for leaf spot. Repeat application in 10 to 14 days. (12 1/2 pounds)

O'K BY



1108/6

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY CAPTAN 80WP


DATE: 9/9/87

GRAPES (eastern U.S.): Black rot, downy mildew
 1½ pounds per 100 gallons. Make first applica-
 tion just before bloom, repeat just after bloom
 and at 10 to 14 day intervals for 1 to 3 more
 applications depending upon the susceptibility
 of the grape variety. Dead Arm (current season
 infection) use 1½ to 2½ pounds per 100 gallons
 dilute spray. Apply 50 gallons per acre when
 shoots are ½ to 1 inch long. Repeat application
 when shoots are 4 to 6 inches long. (6½ pounds)

PICA 50 CHARACTERS
ELITE - 60 CHARACTERS

PEACHES, NECTARINES: Brown rot, scab - Use 1½
 pounds in full pink, bloom, petal fall, shuck,
 cover and preharvest sprays. Applications at
 3 to 4 day intervals may be necessary during bloom
 to control blossom blight. Repeat application
 at 7 to 14 day intervals as needed to maintain
 cover. Continue applications through harvest if
 conditions favor brown rot. If powdery mildew is
 a problem, add 3 pounds sulfur to the petal fall,
 shuck and early cover sprays. If sulfur is
 added, CAPTAN may be reduced to ¾ pound in
 these sprays. Coryneum blight (peach blight,
 shothole) - Use 1½ pounds, apply in pink bud,
 full bloom, petal fall and cover periods as
 necessary and as a post harvest spray, but
 before leaves drop. (6½ pounds)

O.K BY.



SHEET NO.

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY CAPTAN 80WP


DATE: 9/9/87

PEARS (eastern U.S.): Scab, leaf spot - Use 1 1/2 pounds in pre-blossom, petal fall and cover sprays. Repeat cover applications at 7 to 14 day intervals under severe scab conditions. Russetting may be reduced on Bosc pears. Do not use on D'Anjou pears. (10 pounds)

PICA 50 CHARACTERS
ELITE - 60 CHARACTERS

PLUMS (eastern U.S.): Brown rot - Use 1 1/2 pounds in full pink, bloom and petal fall sprays. Repeat applications at 7 to 14 day intervals as needed to maintain cover. Continue applications through harvest if conditions favor brown rot. The addition of a neutral spreader has improved coverage. (6 1/2 pounds)

STRAWBERRIES: Botrytis (gray mold), leaf spot - Apply 2 1/2 to 3 3/4 pounds per acre in sufficient water for thorough coverage. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat at weekly intervals. Under conditions favorable to fruit rot continue applications through harvest period treating immediately after each picking. (5 pounds)



130616

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY CAPTAN BOWE

DATE: 9/9/87

CANTALOUPE, CUCUMBER,

An-

gular leaf spot, anthracnose - Use 2½ pounds per acre in sufficient water for thorough coverage. Begin applications at first signs of infection and repeat at 5 to 7 day intervals or as needed to maintain control. (4 ¾ pounds)

TOMATO: Anthracnose, gray leaf spot, septoria leaf spot - Use 2½ to 3 ¾ pounds per acre in sufficient water for thorough coverage. Under severe infection conditions use 3 ¾ to 5 pounds per acre in sufficient water for thorough coverage. Apply at first signs of infection and repeat at 5 to 7 day intervals or as needed to maintain control. (6½ pounds)

BEST AVAILABLE COPY

PICA 50 CHARACTERS
ELITE - 60 CHARACTERS

O'K BY
[Barcode]

SHEET NO.

12-15

ADVERTISING AND SALES PROMOTION SERVICES

1408/16
ECP

FOR: AGWAY CAPTAN 80WP

DATE: 9/9/87

ORNAMENTALS

AZALEAS: Damping-off of cuttings - Use 2½ pounds. Dip cuttings before bedding. Petal blight - Use 1½ pounds. Apply to soil area around plants and spray flowers just before bloom. Repeat at 7 to 14 day intervals through bloom.


BEGONIAS (TUBEROUS): Damping-off, tuber rot - Use 2½ pounds. Dip tubers 30 minutes, drain and plant.

CAMELLIAS: Petal blight - Use 1 5/8 pounds. Apply to trench soil around plants beginning when flowers start to open. Repeat at 7 to 10 day intervals through bloom.

CARNATIONS: Alternaria leaf spot, rust - Use 1½ pounds. Begin at first sign of disease. Repeat at 7 to 10 day intervals. Shorten intervals during frequent rains and heavy dews. Damping-off of cuttings - Use 2½ pounds. Dip cuttings before bedding.

GLADIOLUS (CORMS): Corm rot and decay, damping-off - Use 1/3 to 1 pound per 10 gallons, dip corms 20 to 30 minutes. Drain and plant.

FIG. 10 SERIALS
10 11 - 1000-10000000



ADVERTISING AND SALES PROMOTION SERVICES

15 of 16
COPY

FOR: AGWAY CAPTAN BOWP

DATE: 9/9/87

ROSES: Black spot, Botrytis blossom blight - Use 1½ pounds. Begin at first growth or first sign of disease. Repeat at 7 to 14 day intervals, and more frequently during frequent rains and heavy dews.

CHRYSANTHEMUM: Botrytis flower blight, Septoria leaf spot- Use 1½ pounds. Apply at first sign of disease, repeat at 7 to 10 day intervals. Damping-off of cuttings - Use 2½ pounds. Dip cuttings before bedding.

SOIL AND GREENHOUSE BENCH TREATMENT: Preplanting for damping off and root rot diseases on seedlings or transplants of roses (and other shrubs, trees, flowers), Lawn seedbeds, vegetables (such as cucumbers & tomatoes) - Use 1½ pounds per 100 gallons water at rate of 15 gallons per 1,000 sq. ft. Cultivate into upper 3 to 4 inches of soil before planting.

Only the applicator is permitted to be in the greenhouse during application of captan. Open vents to greenhouse during application and for at least 1 hour after application.

Workers planting in captan-treated soil must wear chemical resistant gloves.

STORAGE & DISPOSAL (12 pt)

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

STORAGE: Store unused product in original container only. Store in a cool, dry area preferably locked and out of the reach of children and animals. Not for use or storage in or around the home. Protect from excess heat.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a Violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

PICA 50 CHARACTERS
ELITE - 60 CHARACTERS

U.S. BY

SHEET NO.

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY CAPTAN BOWP

DATE: 9/9/87

DISCLAIMER: Buyer assumes all risks of use,
 storage or handling not in strict accordance
 with directions given herewith.

NOTE: This pesticide is to be sold ONLY in this
 original unbroken package.

PICA - 50 CHARACTERS
 ELITE - 60 CHARACTERS

O'K BY