

# AGWAY PHALTAN GARDEN SPRAY

75% Folpet Wettable Powder fungicide for the control of diseases on fruits, roses and other flowers. For home and garden use only.

AGWAY INC. SYRACUSE, NY

75% Folpet Wettable Powder fungicide for the control of diseases on fruits, roses and other flowers.

INERT INGREDIENTS

TOTAL

\*Phaltan is the registered trademark for phthalimide. U.S. Pat. Nos. 2,550,776 & 2,550,777.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See additional precautionary statements on side panel.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

### STORAGE AND DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

Read the entire directions & disclaimer. Use strictly in accordance with label cautions, warnings & directions.

Always mix the recommended amount of Phaltan Garden Spray with a little water to form a thin, smooth paste before making final dilution.

### FLOWERS

Plant	Diseases	Recommendations
Roses	Black Spot Powdery Mildew	Use 1 tablespoon to one gallon of water. Apply when leaves first appear. Repeat at 7 to 10 day intervals or as necessary during growing season. Do not spray at temperatures in excess of 90° F. as injury may occur.
Chrysanthemums	Leaf Spot (Septoria)	Use 1 1/2 tablespoons to one gallon of water. Start treatments immediately after planting. Repeat at 7 to 10 day intervals or as necessary during growing season.
Iris	Leaf Spot (Dodymellina)	Use 1 1/2 tablespoons to one gallon of water. Start treatments immediately after planting. Repeat at 7 to 10 day intervals or as necessary during growing season.
Gladiolus Corms (Preplanting)	Rot	Use 10 tablespoons to one gallon of water. Dip bulbs in this diluted mixture for 20 to 30 min., drain and plant. Keep the mixture stirred.
Carnations	Rust	Use 1 1/2 tablespoons to one gallon of water. Begin application 2 weeks after planting. Continue bi-weekly applications for 3 to 4 applications.
Zinnias Marigolds	Alternaria Leaf Spot	Use 1 1/2 tablespoons to one gallon of water. Treat at least weekly during periods of rapid new growth and severe infection and twice weekly during periods of rapid new growth and severe infection caused by continuous rains or high humidity.
Chrysanthemums Asters Phlox	Powdery Mildew (Erysiph)	Use 1 1/2 tablespoons to one gallon of water. Start treatments immediately after planting. Repeat at 7 to 10 day intervals or as necessary during growing season.
Snapdragons	Anthraxnose Powdery Mildew	Use 1 1/2 tablespoons to one gallon of water. Start treatments immediately after planting. Repeat at 7 to 10 day intervals or as necessary during growing season.
Poinsettia (greenhouse use)	Root Rot (Pythium)	Mix 1/2 cup per 2 1/2 gallons of water and use 1 qt. per 8 sq. ft. of area. Make 1st application at time of planting. Repeat at 10 days.
Cuttings Azaleas	Leaf Spot (Septoria)	Use 22 tablespoons per one gallon of water. Allow to soak for 15 to 30 minutes before planting in rooting media.

### FRUITS & BERRIES

See recommendations on side panel.

Use 4 teaspoons per one gallon of water. Make first application before bloom & repeat at 7 to 14 days until harvest. Do not spray during the critical period just before bloom. Days after petal fall may cause fruit damage if spray is used as a direct fungicide. Red Delicious & Staygreen Apples.

Use 4 teaspoons per one gallon of water. Apply at petal fall & popcorn stage. Repeat in bloom and at 7 to 14 day intervals. Last application to foliage after harvest. Do not use on sweet corn as foliage injury may occur.

Use 4 teaspoons per one gallon of water. For dead arm disease application when new growth is 1 to 2 inches long. Repeat on new growths 4 to 6 inches. For black rot & powdery mildew, make first application just before bloom & repeat immediately after bloom. Depending on the susceptibility of the grape variety, from 3 to 5 sprays may be needed at 7 to 10 day intervals. For Powdery Mildew, make first application 2 to 3 weeks after bloom & repeat at 2 week intervals as needed. Caution: During extended periods of hot, dry weather, foliage injury to several varieties of grapes may occur.

Use 4 teaspoons per one gallon of water. Make first application when berries have set & repeat at 7 day intervals until harvest.

Use 4 teaspoons per one gallon of water. Make first application before first infection. Repeat at 7 day intervals until harvest.

Use 8 teaspoons per one gallon of water. Make first application before bloom period. Repeat at 7 to 10 day intervals in the post bloom period.

Use 4 teaspoons per one gallon of water. Make first application before bloom and repeat at 7 day intervals until after harvest.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS & DOMESTIC ANIMALS

**CAUTION:** May cause eye irritation. In case of contact with eyes, immediately flush with plenty of water for at least 15 minutes. Call a physician.

Avoid inhalation of dust or spray mist. Avoid contact with skin. In case of skin contact, wash with water.

Avoid contamination of feeds or feedstuffs.

**ENVIRONMENTAL HAZARDS:** This product is toxic to fish. Do not contaminate any body of water by cleaning equipment or disposal of wastes.

**PHYSICAL OR CHEMICAL HAZARDS:** This product may be combined with most commonly used insecticides and fungicides with the exception of strongly alkaline materials. Use 7 to 10 hydrated lime.

This product should not be used in combination with herbicides following sprays.

**NOTE:** This product is not to be used on citrus fruits. This is a water soluble product.

**DISCLAIMER:** Buyer assumes responsibility for the proper use of this product. Do not use with other fungicides.

AGWAY INC. SYRACUSE, NY

AGWAY INC. CHEMICAL DIVISION BOX 4933 SYRACUSE NY 13221

NET CONTENTS 1 LB.

A H R