

1056
Pm 12
8590-563
3030

ADVERTISING AND SALES PROMOTION SERVICES

FOR: AGWAY LIQUID FRUIT TREE SPRAY

DATE: 6/16/86

((FRONT PANEL))

Net Contents
8 oz., 1 pt.,
1 qt.

AGWAY

LIQUID FRUIT TREE SPRAY

A water soluble concentrate containing insecticide, fungicide, aphicide, miticide, scabicide and spreader-sticker.

ACTIVE INGREDIENTS:

Captan [N- trichloromethyl thio -4-cyclohexene-1,2-dicarboximide]	11.35%
Related Compounds	0.65%
Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate)	3.50%
*Methoxychlor, technical	12.00%
Carbaryl (1-naphthyl N-methylcarbamate)	3.00%
Xylene range aromatic solvent	62.33%

INERT INGREDIENTS:

7.17%

Total

100.00%

*Equivalent to 10.56% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 1.44% of other isomers and related compounds.

KEEP OUT OF REACH OF CHILDREN (12 pt)

DANGER (18 pt)

Causes irreversible eye damage.

See side panel for additional precautionary statements.

AGWAY INC.
P.O. BOX 4741
SYRACUSE, NEW YORK
13221

EPA Reg. No. 8590-563


EPA Est. No. 8590-PA-1

ACCEPTED
SEP. 12 1986
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 8590-563

*amendments
w. del in ref*

OK BY

ADVERTISING AND SALES PROMOTION SERVICES



R: AGWAY LIQUID FRUIT TREE SPRAY

DATE: 6/16/86

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & (DOMESTIC ANIMALS)

DAWLER
CAUTION: Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin & clothing. Remove & separately launder clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, Call a physician or Poison Control Center immediately. Gastric lavage is indicated if material was taken internally. **DO NOT INDUCE VOMITING** as vomiting may cause aspiration pneumonia.

If on skin, remove contaminated clothing and wash affected areas with soap and water.

If in eyes, flush eyes with plenty of water. Call a physician if irritation persists.

Wear Natural or synthetic rubber, vinyl or plastic gloves impermeable to Captan when using indoors or outdoors. When using outdoors wear long pants & long sleeved shirt and apply with the wind to your back. Wash Nondisposable gloves thoroughly with soap & water before removing. Clothing worn while handling this product must be laundered separately from other clothing before reusing.

Remove residue on edible fruit by wiping or washing before canning or eating. Avoid contamination of foodstuffs, feed or feeding troughs and water receptacles or pets or animals, outdoor

OK BY

ADVERTISING AND SALES PROMOTION SERVICES

3056

NR: AGWAY LIQUID FRUIT TREE SPRAY

DATE: 6/16/86

fish ponds, streams and water supplies.

FIGA 30 CHARACTERS
ELITE 40 CHARACTERS

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Drift & runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL/CHEMICAL HAZARDS
Do not use or store near heat or open flame.

EASY TO USE: Mixes with water instantly; no plugged nozzles; no messy powders to handle, measure or mix; no pre-mixing or straining necessary. Designed especially for home orchards to protect apples, cherries, grapes, peaches, pears, plums and prunes from the ravages of insects and diseases listed on spray chart below.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

SHAKE PRODUCT THOROUGHLY BEFORE USING. Contains micronic particles which settle upon standing and require reblending by agitation. Choose a

OK BY

ADVERTISING AND SALES PROMOTION SERVICES

NR: AGWAY LIQUID FRUIT TREE SPRAY

DATE: 6/16/86

cool, calm period, preferably early morning or evening, when trees will not be subjected to hot direct rays of the sun. Do not apply if rain is expected before leaf surfaces dry. It is easier to prevent disease and scale damage than to cure same, therefore, a preventative 10-14 day spray schedule is recommended. When insects are present, use maximum dilution. When active disease spores are present (warm, humid days) shorten spray intervals to 5-7 days at maximum dilution.

PICA - 30 CHARACTERS
ELITE - 30 CHARACTERS

Food utensils (Teaspoons, Tablespoons) should not be used for food purposes after use in measuring pesticides.

Apply spray to leaves, fruit branches, trunk and base of tree until lightly dripping. Shake sprayer occasionally or agitate to keep spray particles in suspension during application. Follow spray schedule given below in tablespoons per gallon of water. For larger quantities, use 1 gallon per 100 gallons of water. Observe caution statement and cautions for fruit to be treated.

STORAGE & DISPOSAL (12 pt)

Protect from freezing. Store in a cool, dry area, preferably locked and out of the reach of children.

Do not reuse empty container. Wrap container and put in trash collection.

ADVERTISING AND SALES PROMOTION SERVICES

5 of 6

30PV

FR: ACWAY LIQUID FRUIT TREE SPRAY

DATE: 6/16/86

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

PICA 30 CHARACTERS
ELITE 30 CHARACTERS

O'K'BY

60F6

SPRAY CHART

Fruit	Insects/Diseases	Dor- mant	Green Tip	Pre Bloom	Full Pink	Bloom	Petal Fall	1st Cover	
Apple:	apple maggots, codling moths, Jap beetles, leafhoppers, plum curcullos, aphids, bud moth, Farbes scale, eastern tent caterpillar, red banded leaf roller, mites, bitter rot, black rot, frog-eye leaf spot, black pox, botryosphaeria (white rot), botrytis rot, bullseye rot, Bruks fruit spot, flyspeck, cedar rust, quince rust, scab, sooty blotch	Dormant - absence of growth-fall, winter or spring. Dormant sprays are recommended. Use Agway Dormant Oil or Dormant Oil with Ethon up to 7 days before Green Tip application.	1 1/2 tbsp.	2 1/2 tbs.	2 1/2 tbs.		2 1/2 tbs.	2 1/2 tbs.	
	Including fruiting, flowering (edible) crabapple and ornamental crabapple: Scab.			Green Tip -- when branch tips show green tissue	Pre-bloom -- when blossom buds begin to show color	Full Pink -- when blossom buds show full color before opening		Petal Fall -- when last petals have fallen from blossom	1st Cover -- 7 - 10 days after petal fall
Cherry:	plum curcullos, cherry fruit flies, cherry fruit worms, black cherry aphid, fruit tree leaf roller, bud moth, Jap beetle, Botrytis rot, brown rot, leaf spot			1 1/2 tbs.	2 1/2 tbs.	2 1/2 tbs.		2 1/2 tbs.	2 1/2 tbs.
Grape:	flea beetles, Jap beetles, grape leaf skeletonizers, rose chafers, grape berry moths, spider mites, mealy bugs, leafhoppers, black rot, downy mildew (berries only), botrytis rot			1 1/2 tbs.	2 1/2 tbs.	2 1/2 tbs.		2 1/2 tbs.	2 1/2 tbs.
Peach:	plum curcullo, Jap beetle, spider mite, oriental fruit moth, green, peach, black cherry, black peach, rusty plum aphids, brown rot, rhizopus, scab, brown rot (blossom blight), coryneum blight			1 1/2 tbs.	1 1/2 tbs.	2 1/2 tbs.		2 1/2 tbs.	2 1/2 tbs.
Pear:	codling moth, pear slug, fruit tree leaf roller, pear psylla, plum curcullo, red banded leaf roller, mites, mealybugs, fruit spot, scab			1 1/2 tbs.	2 1/2 tbs.	2 1/2 tbs.		2 1/2 tbs.	2 1/2 tbs.
Plum, Prune:	plum curcullos, mealy plum aphid, bud moth, brown rot, russet or lacy scab		1 1/2 tbs.	1 1/2 tbs.	2 1/2 tbs.		2 1/2 tbs.	2 1/2 tbs.	

Bloom - When blossoms are open. Don't spray pollinators. Fungicide alone may be applied for disease