


AGWAY

LORSBAN* 50-SL

Wettable Powder Insecticide



For Use as a Preplant Slurry Treatment of Field Corn, Lima Bean, Snap Bean, and Sweet Corn Seeds to protect Germinating Seeds and Seedlings from Injury by Seed Corn Maggots

ACTIVE INGREDIENTS

Chlorpyrifos [0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate]

INERT INGREDIENTS

TOTAL

50
50
100

* Trademark of The Dow Chemical Company

KEEP OUT OF REACH OF CHILDREN.

WARNING

PRECAUTIONARY STATEMENTS

Hazards To Humans & Domestic Animals

MAY BE FATAL IF SWALLOWED

CAUSES EYE IRRITATION

WARNING: Do not take internally. Do not breathe dust. Do not get in eyes, on skin, or on clothing. Wear a NIOSH approved mask suitable for protection against inhalation of toxic dusts to prevent exposure during use. Wash thoroughly after handling and before eating or smoking. Remove and wash contaminated clothing before reuse. In case of contact, immediately flush eyes with plenty of water and get prompt medical attention if irritation develops or persists. Wash contaminated skin with soap and plenty of water.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water.

Do not contaminate water by cleaning of equipment or disposal of wastes. Use this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

General Information

LORSBAN 50 SL insecticide is a wettable powder formulation designed for slurry treatment of seed prior to planting to protect the germinating seeds and seedlings against injury by seed corn maggots. LORSBAN 50 SL may be applied in combination with captan, chloroneb or thiram fungicides if protection of the seed from fungi, which may cause disease or rot, is also desired.

Seed treated with LORSBAN 50 SL may be stored for up to 14 months without loss of insecticidal effectiveness.

Use and Dosage Recommendations

Field Corn, Lima Beans, Snap Beans, Sweet Corn (For control of seed corn maggots). Use 2 ounces of LORSBAN 50 SL per 100 pounds of seed by applying to the seed as a slurry treatment prior to planting.

Slurry Treatment

Disperse the required amount of LORSBAN 50 SL in a water solution containing 0.5% by weight of METHOCEL* A4C Premium methylcellulose or other acceptable sticking agent. Use 12 to 15 fluid ounces of solution for each 100 pounds of seed to be treated. Generally, rough surfaced seeds such as corn will require a greater volume of solution for complete coating than smooth surfaced seeds such as beans. If a fungicide is used, it should be added to the slurry in accordance with its label directions. Mix the slurry with the seed using a rotating blender or other suitable mixing equipment that will ensure each seed is uniformly coated. **DO NOT MIX THE SEED AND PRODUCT WITH BARE HANDS.**

Use Restrictions

Do not use treated seed for human consumption, as feed for livestock or poultry, or for oil purposes. Label treated seed as follows: **TREAT 1) SEED DO NOT USE FOR FOOD, FEED, OR OIL.**

STORAGE & DISPOSAL

Prohibitions: Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: Dispose of bags according to approved Federal, State or local procedures under the Resource Conservation and Recovery Act.

NOTE: This pesticide is to be sold ONLY in this original, unbroken package.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

© 1981 AGWAY INC.

EPA Est. No. 8537-PA-1



AGWAY INC., CHEMICAL DIVISION
BOX 4933, SYRACUSE, NY 13221

Net Weight 50 lbs.

A 7 81