FOR:	AGWAY BETASAN 7-G	DATE	:1/24/8
	(((FRONT PANEL)))	NET WEIGHT	e la filia
	A	GWAY	
	BETAS	AN® 7-G	
·	GRANULES - A Selective	Herbicide for Pre-Emer-	
4 1 '	gence Control of Crabg	cass and Annual Bluegrass	
	in Turf, Ornamentals,	and Ground Covers.	
	Contains 7 pounds Activ	ve Ingredient Per 100 Pounds	
	ACTIVE INGREDIENT:		
1	BENSULIDE:		
Underlined	letters S-(0,0-diisopropyl pho	osphorodithicate)	
hould be it	Ester of N-(2-mercapto	oethyl) benzene- 7.0%	
cized.	INERT INGREDIENTS:	93.0%	
		TOTAL 100.0%	
	Registered Trademark o	of Stauffer Chemical Co.	
	KEEP OUT OF REACH OF CH	ILDREN (12 pt.)	
	WARNING (18 pt.)		
	STATEMENT OF PRACTICAL	TREATMENT	, e
	Call a Physician Immedi	ately! If a known exposure	•••
	1	is suspected, do not wait	• ·
	for symptoms to develop	. Immediately initiate the	* * *
	recommended procedures	described below and simul-	• • •
	taneously contact a phy	sician, the nearest hospital	¢
	or a Poison Control Cer		
<u> </u>			
		SHE	ET NO

.

i

•

195

• •

,-

+-

÷

.

<u>FOR:</u>	AGWAY BETASAN 7-G	DATE: 1/24/83
	(((FRONT PANEL)))	(· · · · · · · · · · · · · · · · · · ·
	Inform the person contacted of the type and ex-	
	tent of exposure, describe the victim's symptoms	
	and follow the advice given.	
	NOTE: Be sure to advise the person contacted	
	that the active ingredient in this compound is	
	a cholinesterase inhibitor and follow the advice	
	given.	
	NOTE TO PHYSICIAN: Exposure to the active in-	
	gredient may cause cholinesterase inhibition.	
	Atropine by injection is antidotal. 2-PAM is	
	also antidotal when administered early and in	
	conjunction with atropine.	
	IF SWALLOWED: Call a physician or Poison Control	
	Center immediately. Immediately give large	
	amounts of water and induce vomiting by gagging	
	the victim with a finger placed on the back of the	е
	victim's tongue. Give water until vomitus is	· · ·
	clear. Do not induce vomiting or give anything	0 к зу
	by mouth to an unconscious person.	· . ·
		· · ·
	IF INHALED: Remove victim to fresh air. If not	
	breathing, give artific:al respiration, preferab	Y
	mouth-to-mouth. If breathing is difficult, give	
	oxygen, preferably with a physician's advice.	
		SHEET NO

-

1-

•

FOR:	AGWAY BETASAN 7-G	DATE: 1/24/83
	(((FRONT PANEL))) <u>IF ON SKIN</u> : Flush with plenty of water for at least 15 minutes. Remove contaminated clothing and shoes. Get medical attention if irritation occurs. Wash clothing before re-use.	
	IF IN EYES: Hold eyelids apart and flush eyes with large amounts of running water for at least 15 minutes. Get medical attention.	
	See side panel for additional precautionary statements.	
	AGWAY INC.	
	CHEMICAL DIV.	
	BOX 4933 EPA Reg. No. 8590-565	
	SYRACUSE, NY 13221 EPA Est. No. 8590-PA-1	
		С'К ЗҮ
		SHEET NO.

;

٠.

. ..

1

ADVERTISING AND SALES PROMOTION SERVICES

AGWAY BETASAN 7-G DATE:1/24/83 FOR: (((REAR PANEL))) Þ PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS WARNING: . Harmful if swallowed. . Avoid breathing dust. . Causes substantial but temporary eye injury. Do not get in eyes, on skin, or on clothing. Wear goggles or safety glasses when handling. . Wash thoroughly with soap and water after handling and before eating or smoking. . Mich all concuminated clothing with soap and hot water before re-use. ENVIRONMENTAL HAZARDS . This product is toxic to fish. Do not apply directly to any body of water. Do not apply where OK BY runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. . Do not apply when weather conditions favor drift from areas treated. SHEET NO

•

•....

ADVERTISING AND SALES PROMOTION SERVICES $\frac{1}{2}$

,-

ŀ

Þ

ŀ

4

1

۲

• •

٨

L,

. :

•

•

R:	AGWAY BETASAN 7-G	DATE:1/24/83
•	. Apply this product only as specified on this label.	
	PRODUCT INFORMATION	
	BETASAN 7G is recommended for use on home lawns,	
	parks and golf courses, on ornamental plants	
	and on ground covers for control of crabgrass,	
	annual bluegrass and other annual weeds listed.	
	Application of BETASAN 7-G must precede emergence	
	of weeds from the soil. Crabgrass will emerge	
	from the soil from four to six weeks before it is	
	visible above lawn sod. If weeds are visible	
	above the soil, it is too late to apply BETASAN 7-G.	
	Remove leaves, dead or tall grass, and other	
	debris before applying BETASAN 7-G to turf areas.	
	Any weeds present around ornamentals must be	
	removed or worked into the soil before applying	
	BETASAN 7-G.	ук ч ∨
		SHEET NO

-

FOR:	AGWAY BETASAN 7-G	DATE: 1/24/83
	WEEDS CONTROLLED	• · · · · ·
Underlined	Smooth Crabgrass (Digitaria ischaemum)	
words should	Hairy Crabgrass (<u>Digitaria sanguinalis</u>)	
be italicized.	Redroot Pigweed (Amaranthus retroflexus)	
	Barnyardgrass (Watergrass) (EchinochLoa_spp.)	
	Lambsquarters (Chenopodium album)	
	Goosegrass (Eleusine indica)	
	Annual Bluegrass (<u>Poa annua</u>)	
	Shepherdspurse (Capsella Bursa-pastoris)	
	Deadnettle (Henbit) (Lamium_spp.)	
	(NOTE: BETASAN 7G does not control established	
	weeds.)	
	STORAGE AND DISPOSAL (12 pt.)	
	Do not contaminate water, food or feed by stor-	
	age or disposal.	
	STORAGE: Store in a cool place.	
	PESTICIDE DISPOSAL: Pesticide that cannot be	
	used according to label instructions must be	О К <u>В</u> Ү
	disposed of according to applicable Federal, sta	te
	or local procedures.	•
	CONTAINER DISPOSAL: Completely empty bag by sha and tapping sides and bottom to loosen clinging	
	ticles. Empty residue into application equipmen Then dispose of bags in a sanitary landfill or b	t.
	incineration if allowed by State and local authorities.	SHEET NO
	1 www.exv**/**/**	4

••

. •

1

45

. : •2

. ±

ł

19-1

• •

•

.

•

•

۰.

•

	AGWAY BETASAN 7-G	DATE: 1/24/83
,	GENERAL USE PRECAUTIONS	
	. READ ALL DIRECTIONS BEFORE USING.	
	. BETASAN 7-G should be used only for recom-	
	mended uses and at recommended rates.	
	. Sweep up spills on walks or patios and hose	
	area.	
	. Do not allow children or pets on treated	
	lawns until BETASAN 7-G has been watered into	
	the soil and grass is completely dry.	
	. BETASAN 7-G is recommended for use on mineral	
	soils only.	
	. Do not apply peat moss to lawn, ground covers	
	or established ornamentals before treating	
	with BETASAN 7-G.	
	DIRECTIONS FOR USE	0'K 8Y
	It is a violation of federal law to use this	i
	product in a manner inconsistent with its label-	
	ing.	
		SHEET NO

FOR: DATE: 1/24/83 AGWAY BETASAN 7-G TIMING OF APPLICATION The most effective program for control of crabgrass, goosegrass, annual bluegrass and other annual weeds is one specifically designed for local conditions. Two or more applications per year may be necessary, using the recommended rates given below for a particular weed problem. For additional information regarding timing and rate of application, consult weed specialists of the State Agricultural Extension Service or the State Agricultural Experiment Station. GRASS LAWNS: BETASAN 7-G should be applied only to well established grass lawns, BETASAN 7-G can be used on the following established turf grasses: Bahia Pensacola Bahia Bentgrass spp. Perennial Ryegrass Bermudagrass spp. Poa Trivialis Perennial Bluegrass spp. Saint Augustine Centipede окву Red Top Fescue spp. Zoysia DICHONDRA LAWNS: BETASAN 7-G can be applied to dichondra lawns at the time of seeding or any time thereafter. SHEET NO

. .

,-

•

•

FOR:	AGWAY BETASAN 7-G		DATE: 1/24/83
	ORNAMENTALS AND GROUN	D COVERS: BETASAN 7-G	▲ 1 1 1 1 1 1
	should be applied only	y to the following well	
	established ornamenta	ls and ground covers.	
	Alyssum	Campanula	
	Aster	Gazania	
	Bachelor's Button	Coral Bell	
	Calendula	Daisy	
	Candy Tuft	Marigold	
	Daffodil	Pansy	
	Dahlia	Primrose	
	Freesia	Gladiolus	
	Abelia	Narcissus	
	Azalea	Juniper	
	Azara	Monterey Cypress	
	Boxwood	Monterey Pine	
	Daphne	Myrtle	
	Holly	Privet	
	Ajuga	Hypericum	
	Ice Plant	Tulip	•
	Ivy	Pyracantha	0.K 9.A
	Pachysandra	Sandankwa	
	Stock	Star Jasmine	
	Sweet Pea	Tobira	••
	Wall Flower	Xylosma	
	Zinnia	Periwinkle	
	Ranunculus	Sedum	
	Wild Strawberry		SHEET NO

FOR:	AGWAY BETASAN 7-G	DATE: 1/24/
	For control of Crabgrass, apply BETASAN 7-G at	∮
•	the recommended rate (see "RATES OF APPLICATION")
	any time from fall-September or October - throug	h
	early spring, for spring and summer control.	
	With a late winter or an early spring applicatio	1 ,
	the application must be made prior to crabgrass	
	emergence (see "DIRECTIONS FOR APPLICATION").	
	For control of Annual Bluegrass and other annual	
	weeds, apply BETASAN 7-G at the recommended rate	
	(see "RATES OF APPLICATION") before the weeds	
	emerge from the soil.	
	Where germination of crabgrass, annual bluegrass	
	and other weeds extends into the late summer and	
	early fall, a second application of BETASAN 7-G	
	at the recommended rate should be made four to	
	five months after the initial ppplication.	
	Consult weed specialists of the State Agricul-	-
	tural Extension Service or the State Agricultura	1 · 0'# 3Y
	Experiment Stations for additional information,	
	as the timing and rate of application needed wil	ע
	vary with local conditions.	· .
	BETASAN 7-G should be applied with a lawn spread	er
	or any other applicator that will give uniform	SHEET NO

•

1.

•

. :

•

-7

	1
coverage. A more uniform application can be	♦
made by spreading half of the required amount	
of BETASAN 7-G over the area to be treated,	
then applying the remainder at right angles	
or crosswise. Avoid misses by overlapping	
slightly. Sprinkle the area with water for 10	
to 15 minutes after application to wash the	
BETASAN 7-G down into the soil where the weed	
seeds are located.	
RATES OF APPLICATION	
Apply a maximum of 357 pounds of Betasan 7-G	
per acre (8.2 pounds per 1,000 sq. ft.) per	
year in two or more applications depending on	
the weed species and degree of infestation.	
CRABGRASS: Apply 107 to 143 pounds of BETASAN	
7-G uniformly on one acre (2.4 to 3.2 pounds	
per 1,000 sq. ft.)	
ANNUAL BLUEGRASS, GOOSEGRASS AND OTHER ANNUAL	OKJY
WEEDS: Apply 180 pounds of BETASAN 7-G uni-	
formly on one acre (10 pounds per 2,500 sq. ft.)	
Sprinkle the area with water for 10 to 15 minute	: S
after application to wash the BETASAN 7-G down	
into the soil where the weed seeds are located.	SHEET NO

,

٠.

2.

١

ł

۶

.

.

•

•

-7

•

OR:	AGWAY BETAS	AN 7-G			DATE: 1/24
•	When a weed	control program	n involves	two to three	• •
	applications	of BETASAN 7-0	for crabg	rass, goose-	
	grass and ar	nual bluegrass,	the rate	of applica-	
	tion for goo	segrass and ann	ual bluegr	ass may be	
	reduced to 1	.07 pounds to 14	13 pounds p	er acre,	
	using the hi	gher rate for h	neavier info	estations.	
	BETASAN 7	G RATE TABLE AN	ND SPREADER	SETTINGS	
Make of De	sired Rate	Desired Rate	Spreader	Settings	Spread
Spreader	Per Acre F	Per 1,000 Sq.Ft.	Once Over,	/Twice Over	Width
Cyclone (1)	107 lbs.	2.4 lbs.	3.55	3.25	5 ft.
	180 lbs	4.1 lbs.	3,95	3.45	5 ft.
Scott (1)	107 lbs.	2.4 lbs.	4.3	3.4	20 in.
Drop	180 lbs.	4.1 lbs.	5.6	4.0	20 in.
Scott	107 lbs.	2.4 lbs.	B-0.9-C	A-0.4-B	5 ft.
Centrifugal	180 lbs.	4.1 lbs.	G-0.8-D	B-0.3-C	5 ft.
(1)(2)(3)					
	(1) Based c	on 2 mile per ho	our rate of	travel	
	(2) Should	be used as a gu	ide only -	recalibrate	• О K ВY
	frequer	tly			
	(3) Numbers	; indicate how f	ar from fi	rst letter	
	toward	second letter e	ex. B-0.9-C	= 9 tenths	
	of dist	ance from B to	c.		
					SHEET NO

.

•

.

ι.,

.

•....

٠

<u>ه</u>د -

FOR:	AGWAY BETASAN 7-G	DATE: 1/24/83
	RESEEDING	4 ·
	Any material that provides effective control of	
	annual grasses may affect new lawn grass seedling	ngs
	which germinate after treatment. Proper fertili	
	zation and watering will encourage the existing	
	turf grass to fill in the thin or bare spots	
	formerly occupied by crabgrass.	
	The following procedures are suggested in order	
	to minimize any stunting of turf grasses which	
	may occur when reseeding into BETASAN 7G treated	
	areas:	
	1. Do not reseed within 4 months after applicat	ion.
	2. Renovate the area to be reseeded by raking	
	thoroughly to scarify the soil surface.	
	Where practical, mix peat in with the soil t	.0
	provide an optimum seed bed.	
	3. When turf diseases or physiological disorder	s
	necessitate reseeding sooner than 4 months	*
	after BETASAN 7-G application, apply powdere	d -
	activated charcoal prior to reseeding to	. O K 3Y
	deactivate the BETASAN 7-G.	
	Apply the activated charcoal at a rate of 7	-
	pounds per 1,000 square feet in 14 gallons o	£
	water (300 pounds per acre in 600 gallons of	
	water).	
		SHEET NO
	1	

1

•....

٦

FOR	AGWAY	BETASAN 7-G	DATE:	1/24/83
)	Following application, irrigate the turf with	1	· .
		sufficient water to wash the charcoal into the so:	11.	
		DISCLAIMER: Agway disclaims all implied warrantic	s,	
		including implied warranties or merchantability and	d	
		fitness for a particular purpose. Buyer shall be		
		solely responsible for any and all injury, loss of	ł	
		damage which results from the use of this product		
		in any manner which is inconsistent with the labe	1	
		directions, warnings or cautions. Any damages		
	i	shall be limited to direct damages, not to exceed		
		the purchase price of the product and shall not		
		include any indirect, incidental or consequential		
	1	damages.	ļ	
		NOTE: This pesticide is to be sold ONLY in this		
		original unbroken package.	Í	
				• • • •
			· · · ·	
			OKBY	• • • •
			ļ	
			SHEET	VO
]	
		•		

,

۲

÷.,

. . .

*

L.,

:

.

•.....

7

_1