

GALVINA FLOWERS

FOR AGRICULTURAL OR COMMERCIAL USE ONLY
Fungicide for Fruit, Vegetables and Seed Treatment

An aqueous suspension which contains 3.8 lbs. active ingredient per gallon.

ACCEPTED
JAN 20 1968
U.S. Dept. of Agriculture
Agricultural Chemicals Administration
Washington, D.C. 20250

ACTIVE INGREDIENTS	PERCENTAGE
Captan (2,4,5-trichlorophenyl methylcarbamate)	35.5%
Formalin (37% formaldehyde)	2.1%
Inert Ingredients	61.0%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN CAUTION
See additional precautionary statements on side panel

BEFORE USING MIX COMPONENTS THAT CONTAIN CAPTAN OR FORMALIN: Do not use these components on crops which are treated with formalin. Do not use formalin on crops which are treated with captan. Do not use captan on crops which are treated with formalin. Do not use captan on crops which are treated with formalin.

DIRECTIONS FOR USE

Use a solution of formalin to use this product in a manner consistent with its labeling.

PRECAUTIONS

Do not use this product on grapes, raisins, table grapes, figs, and other crops which are treated with formalin. Do not use formalin on crops which are treated with captan. Do not use captan on crops which are treated with formalin.

Do not use this product on crops which are treated with captan. Do not use captan on crops which are treated with formalin. Do not use formalin on crops which are treated with captan.

Do not use this product on crops which are treated with formalin. Do not use formalin on crops which are treated with captan. Do not use captan on crops which are treated with formalin.

Do not use this product on crops which are treated with captan. Do not use captan on crops which are treated with formalin. Do not use formalin on crops which are treated with captan.

Do not use this product on crops which are treated with formalin. Do not use formalin on crops which are treated with captan. Do not use captan on crops which are treated with formalin.

Do not use this product on crops which are treated with captan. Do not use captan on crops which are treated with formalin. Do not use formalin on crops which are treated with captan.

Do not use this product on crops which are treated with formalin. Do not use formalin on crops which are treated with captan. Do not use captan on crops which are treated with formalin.

is necessary for that crop, especially in the pre-pollination stage. Apply this product from pollination to harvest. Do not use this product on crops which are treated with captan. Do not use captan on crops which are treated with formalin.

APRICOTS: Brown rot, gray blotch, Jockey Rot. Use 1 qt. Apply at bloom, red bud stage and repeat at 75 percent petal fall.

CHERRIES: Brown rot, cherry leaf spot. Use 1 qt. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

CRUICIFEROUS: Downy mildew, black rot, stem rot. Use 1 qt. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

GRAPES: Black rot. Use 1 qt. Make two sprays, before bloom and just after bloom. Follow with at least 3 others, plus before the bunch is set.

LETTUCES, PEACHES: Downy mildew, peach leaf curl, brown rot. Use 1 qt. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

PEACHES: Brown rot, leaf curl. Use 1 qt. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

STRAWBERRIES: Downy mildew, leaf spot. Use 1 qt. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

ASPARAGUS PRE-PLANTING ROOT DIP (For Dipping of Potatoes): Use 1 qt. per 100 gal. water. Dip roots of potatoes for 24 hours.

BEANS: PEAS: Downy mildew, leaf spot. Use 1 qt. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

CANTALOUPES, CUCUMBERS, PUFFBALLS, SUMMER AND WINTER SQUASH, WATERMELONS: Angular Leaf Spot, Anthracnose, Downy Mildew. Use 1 qt. per 100 gal. water or apply 2

CAPSAICIN: Use 1 qt. per 100 gal. water. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

CAPSAICIN: Use 1 qt. per 100 gal. water. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

PEACHES: Brown rot, leaf curl. Use 1 qt. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

STRAWBERRIES: Downy mildew, leaf spot. Use 1 qt. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

PEACHES: Brown rot, leaf curl. Use 1 qt. Apply at green cluster, and repeat at 50 percent petal fall. Repeat at 75 percent petal fall.

POST HARVEST DIP: Use 1 qt. per 100 gal. water. Dip fruit for 24 hours before storage.

FOR USE AS A SEED TREATMENT: This product is used to protect seeds from rot and disease. Do not use on seeds which are treated with captan. Do not use captan on seeds which are treated with formalin.

TREATED SEED: Do not use treated seed on crops which are treated with captan. Do not use captan on crops which are treated with formalin.

ENVIRONMENTAL HAZARDS: This product is toxic to fish. Do not apply directly to water.

DISCLAIMER: Seller assumes no liability for use of this product on crops not listed on the label.

Crop	Rate	Crop	Rate
Apples	1 qt./100 gal.	Corn	1 qt./100 gal.
Apricots	1 qt./100 gal.	Cucumbers	1 qt./100 gal.
Asparagus	1 qt./100 gal.	Eggplants	1 qt./100 gal.
Capers	1 qt./100 gal.	Fig	1 qt./100 gal.
Cherries	1 qt./100 gal.	Garlic	1 qt./100 gal.
Citrus	1 qt./100 gal.	Green beans	1 qt./100 gal.
Crucifers	1 qt./100 gal.	Guava	1 qt./100 gal.
Cucumbers	1 qt./100 gal.	Lettuce	1 qt./100 gal.
Eggplants	1 qt./100 gal.	Peaches	1 qt./100 gal.
Fig	1 qt./100 gal.	Pears	1 qt./100 gal.
Garlic	1 qt./100 gal.	Plum	1 qt./100 gal.
Green beans	1 qt./100 gal.	Potatoes	1 qt./100 gal.
Guava	1 qt./100 gal.	Raspberries	1 qt./100 gal.
Lettuce	1 qt./100 gal.	Strawberries	1 qt./100 gal.
Peaches	1 qt./100 gal.	Summer squash	1 qt./100 gal.
Pears	1 qt./100 gal.	Watermelons	1 qt./100 gal.
Plum	1 qt./100 gal.	Winter squash	1 qt./100 gal.
Potatoes	1 qt./100 gal.		
Raspberries	1 qt./100 gal.		
Strawberries	1 qt./100 gal.		
Summer squash	1 qt./100 gal.		
Watermelons	1 qt./100 gal.		
Winter squash	1 qt./100 gal.		

1 qt./100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

Use 1 qt. per 100 gal. water for 24 hours before storage.

AGWAY INC. - CHEMICAL DIV.
PO BOX 4333
SYRACUSE, NY 13221

NET CONTENTS 1 GAL. A 6/82