

ADVERTISING AND SALES PROMOTION SERVICES

COPY

AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

((FRONT PANEL))

NET CONTENTS  
5 GAL.

PICA - 50 CHARACTERS  
ELITE - 60 CHARACTERS

AGWAY

CAPTAN 4 FLOWABLE

Agricultural Fungicide for Fruit & Vegetables.  
An aqueous suspension which contains 3.8 lbs.  
active ingredient per gallon.

ACTIVE INGREDIENTS:

\*Captan 38.13%

Related Derivatives 0.87%

INERT INGREDIENTS: 61.0%

8590-588  
TOTAL 100.0%

*Per self Cert. of  
DCT 19/87*

\*N-Trichloromethylthio-4-cyclohexene-1,2-  
dicarboximide

KEEP OUT OF REACH OF CHILDREN (12 pt.)

DANGER (18 pt) PELIGRO (18 pt)

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a  
physician.

If swallowed: Drink promptly a large quantity of  
milk, egg whites, gelatin solutions, or if these  
are not available, drink large quantities of  
water. Avoid alcohol.

If inhaled: Remove victim to fresh air. If not  
breathing give artificial respiration, preferably  
mouth-to-mouth. Get medical attention.

PRECAUCION AL USUARIO: Si Usted No lee ingles, No  
use este producto hasta que le etiqueta haya sido  
explicado ampliamente

O'K BY.

SHEET NO.

2012

ADVERTISING AND SALES PROMOTION SERVICES

COPY

AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

See additional precautionary statements below.

AGWAY INC.  
P.O. BOX 4741  
SYRACUSE, NEW YORK 13221

PLA - 50 CHARACTER.  
LITE - 10 CHARACTER.

EPA Reg. No. 8590-588

EPA Est. No. 8590-PA-1

O'K BY

SHEET NO.

1b-11

20612

AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

((SIDE PANEL))

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin & clothing. Remove & separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

This Pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Drift & runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

In the event that treated seeds are spilled & cannot be salvaged, completely cover them with soil.

PLA - NO CHARACTER  
ELITE - NO CHARACTER

078Y  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

SHEET NO.

40012

WAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

((REAR PANEL))

PLA - 10 CHARACTER  
LIFE - 10 CHARACTER

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse.

Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

Fieldworkers and harvesters must wear chemical resistant gloves. Leather or fabric gloves are not acceptable.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

A Chemical resistant apron must be worn when mixing and loading this product.

During application only, from a tractor with a completely enclosed cab or from an aircraft with an enclosed cockpit a long sleeved shirt and long legged pants must be worn. Chemically resistant gloves must be available in the cab or cockpit and worn while exiting.

RE-ENTRY

Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

O'K BY: [grid of dots]

SHEET NO.

50612

# ADVERTISING AND SALES PROMOTION SERVICES

# COPY

AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

### DANGER

(Insert area or field description) treated with Captan on (insert data of application). Do not enter treated areas within 4 days following application unless protective clothing is worn. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Captan and describe his condition. For further information see the "Statement of Practical Treatment".

Oral warnings must include the same information that is in the written warnings.

BEFORE USING MIX CONTENTS THOROUGHLY - STIR OR ROLL! Pour recommended amount of this material into nearly filled spray tank. Add balance of water. Maintain agitation during filling and spraying operations. Do not allow mixture to stand. Use a screen not finer than 50 mesh in entire system. Do not combine with emulsifiable liquids or wettable powders unless previous experience has proven them to be physically compatible and safe to plants.

PICA - 50 CHARACTERS  
ELITE - 60 CHARACTERS

O'K BY

SHEET NO.

609612

AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

Observe use limitations given on the label for specific crops.

In order that pesticidal residues on food and forage crops will not exceed tolerances established by the Federal Food and Drug Administration, use only at recommended rates and intervals. Do not apply or allow to drift onto adjoining food, fiber or pasture crops. The grower is responsible for residues on his crops as well as for damage caused by drift from his property to that of others.

Consult State Agricultural Experiment Stations or the State Agricultural Extension Service for additional information, as the timing of applications needed will vary with local conditions.

RECOMMENDATIONS

Unless otherwise stated, dosage rates on this label are given in terms of CAPTAN 4 FLOWABLE per 100 gals. of water.

ALMONDS: Brown rot, scab, shot hole - Apply 1-1½ qts. at popcorn, bloom and petal fall stages.

POCA - 50 CHARACTER  
ELITE - 50 CHARACTER

O'K BY.

SHEET NO.

5-11

70612

ADVERTISING AND SALES PROMOTION SERVICES

COPY

AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

APPLES - Apple Scab - Captan is most effective when used in a protective schedule where the spray is applied before infection occurs. Primary infection of apple scab may occur at any time during and after the green tip stage if over-wintering scab is mature and there is sufficient moisture to discharge spores.

Thorough coverage is necessary for best results, especially in the pre-pink and pink stage spray applications. Danger from primary infection may extend through the time of the first cover spray. In pre-blossom sprays, use 1 qt. per 100 gallons of water in high volume application. In low volume sprays, apply 2½ gals. per acre. In mid-summer, the dosage may be reduced to ½ qt. per 100 gal. in high volume application, or 1½ gals, per acre in low volume sprays. Bitter rot, black rot, Botryosphaeria, black pox, bull's eye rot: use 1 qt. per 100 gals. Make 1 or 2 applications with late cover sprays and 1 or 2 preharvest applications.

APRICOTS: Brown rot (twig blight), Jacket Rot - Use 1 qt. Apply at bloom, red bud stage and repeat at 75 percent petal fall.

CHERRIES - Brown rot, cherry leaf spot - Use 1 qt. Apply at pre-bloom, bloom and petal fall. For severe infection, use 2 qts. Repeat in cover sprays.

PICA 50 CHARACTERS  
LINE 10 CHARACTER

O'K BY: [Dotted grid]

For [Dotted grid]

12

# ADVERTISING AND SALES PROMOTION SERVICES

# COPY

AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

PGA - 50 CHARACTER  
TITLE - 40 CHARACTER

GRAPES - Black Rot - Use 1 qt. Make two sprays before bloom and just after bloom. Follow with at least 3 cover sprays before the bunches close.

NECTARINES, PEACHES - Coryneum Blight (peach blight), brown rot, scab - Use 1 qt. Apply at pink bud, full bloom and petal-fall stages. Repeat at 10-14 day intervals after petal-fall. Also, make late applications for Coryneum Blight.

PLUMS - Brown Rot - Use 1 qt. Apply at red bud or popcorn stage. Repeat at bloom and 75% petal-fall stages. Also repeat later in season as conditions warrant.

STRAWBERRIES - Botrytis Fruit Rot - Use 1-1½ qts. Apply 200 gals. of spray per acre or an equivalent of 2-3 qts. per acre in sufficient water for even coverage. Make first application when new growth starts in the spring before the fruit starts to form. Repeat at weekly intervals. Under severe infection conditions make con-

OKBY

SHEET NO.



AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

CANTALOUPES, CUCUMBERS,

Angular Leaf Spot,

Anthracnose, Downy Mildew - Use 1 qt. per 100 gals water, or apply 2 qts. per acre in sufficient water for good coverage. Apply at first sign of disease and repeat at 5 - 7 day intervals or as needed to maintain disease control.

TOMATOES: Anthracnose, Early and Late Blight,

Gray Leaf Spot, Septoria Leaf Spot - Use 1 qt.

per 100 gals. of water, or apply 2-3 qts. per acre in sufficient water to provide good coverage. Severe infection Conditions: Use 2 qts. per 100 gals. water, or apply 3-4 qts. per acre with adequate water. Apply at first sign of infestation and repeat at 5-7 day intervals or as needed to maintain control.

PLA - 0 CHARACTER  
LELE - 0 CHARACTER

OK BY

SHEET NO.

8-11

100012

# ADVERTISING AND SALES PROMOTION SERVICES

COPY

AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

FOR USE AS A SEED TREATMENT

This product is formulated for treatment of seeds for control of seed and soilborne diseases, including damping-off and seed decay organisms, and seedling blights.

IMPORTANT

The Federal Government and most states have seed laws or regulations requiring an appropriate caution statement be affixed to bags containing treated seeds, such as:

Treated Seed

Do Not Use for Food, Feed or Oil Purposes.

Mix thoroughly before use or use entire container at one time. Do not combine colloidal suspensions, wettable powder or emulsifiable concentrates unless previous experience assures compatibility. Store in a cool place. Do not store in direct sunlight. Protect from freezing temperatures. This product is to be used in liquid or slurry treaters.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues.

In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates.

OKBY

110612

33PV

# ADVERTISING AND SALES PROMOTION SERVICES

AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

## USE PRECAUTION

When using formulations that do not contain dye, to comply with 21CFR Chapter 1, Section 2.25, all seed treated with an economic poison must be colored to distinguish and prevent subsequent inadvertent use as a food for man, or feed for animals.

Application rates per 100 lbs. of seed of diluted product.

Dilute 1 gallon with 5 gallon of water then add amount listed after seed.

### SEED

|  |       |
|--|-------|
| CORN - Field   | 210cc |
| Sweet  | 440cc |
| WHEAT*   | 210cc |
| OATS*  | 320cc |
| BARLEY*  | 250cc |
| FLAX   | 410cc |
| PEAS   | 280cc |
| CUCUMBER, CANTALOUPE, TOMATO   | 320cc |
| SQUASH, PUMPKIN, WATERMELON and MUSKMELON  | 220cc |
| COTTON - Acid Delinted   | 320cc |
| Reginned   | 480cc |
| CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS,<br>COLLARDS, KALE, MUSTARD, TURNIP | 160cc |
| PEPPERS (California Wonder)  | 320cc |
| COWPEAS  | 320cc |
| RYE*   | 280cc |

\*For Seed and Seedling Rots, not for Bunt Smut.

Dilute 1 gallon with 3 gallons of water then add amount listed after seed.

|                                |       |
|--------------------------------|-------|
| BEANS**                        | 190cc |
| SOYBEANS                       | 180cc |
| SORGHUM - hulled (Kernel Smut) | 320cc |
| SUGAR BEETS - West             | 440cc |
| SPINACH                        | 440cc |

\*\*Do not use on Lima Beans.

# ADVERTISING AND SALES PROMOTION SERVICES

12 of 12  
COPY

FOR: AGWAY CAPTAN 4 FLOWABLE

DATE: 9/9/87

Dilute 1 gallon with 2 gallons of water then add amount listed after seed:

|  |        |
|--|--------|
| CLOVER, ALFALFA, LESPEDEZA, TREFOIL    | 450 cc |
| BEETS, SWISS CHARD, SUGAR BEETS - East | 650 cc |
| BLUE GRASS, GRASSES                    | 440 cc |

### STORAGE AND DISPOSAL (12 pt)

PROHIBITIONS: Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited.

STORAGE: Store unused product in original container only. Store in a cool, dry area out of reach of children and animals. Not for storage in or around the home. Protect from excess heat. Protect from freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a Violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

PICA - 50 CHARACTER  
ELITE - 60 CHARACTER

OK BY

SHEET NO.

11-11