

ADVERTISING AND SALES PROMOTION SERVICES

3037

FOR: AGWAY ZINEB-SEVIN [®] 4.5-5D

DATE: 1/10/83

((FRONT PANEL))

NET WT.
1#

AGWAY

ZINEB-SEVIN [®] 4.5-5D

To control insects and diseases of Vegetables.

ACCEPTED
with COMMENTS
in EPA Letter Dated:

FEB 11 1983

Under the Federal
Fungicide, Insecticide,
and Plant Growth Regulator
Act as amended,
registered under
8590-619

ACTIVE INGREDIENTS:

Zineb (Zinc ethylene bisdithiocarbamate) (metallic zinc content 1.10%)	4.5%
Carbaryl (1-Naphthyl N-Methylcarbamate)	5.0%

INERT INGREDIENTS:

90.5%

TOTAL 100.0%

[®] Trademark of Union Carbide Corp. for the active ingredient Carbaryl (1-Naphthyl N-Methylcarbamate)

KEEP OUT OF REACH OF CHILDREN (12 pt.)

CAUTION (18 pt.)

See additional precautionary statements on rear panel.

AGWAY INC.
CHEMICAL DIVISION
BOX 4933
SYRACUSE, NY 13221

EPA Reg. No. 8590-619
EPA Est. No. 8590-PA-1

O'K BY

SHEET NO 1 - 5

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY ZINEB-SEVIN^R 4.5-5D

DATE: 1/10/83

((REAR PANEL))

PICA - OPERATIONAL
ELECTRO-CHEMICAL

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed, inhaled or absorbed through skin. Avoid contact with skin, eyes or clothing. Do not breathe dust. Wash thoroughly after using.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

Wear protective clothing (long pants, long sleeve shirt, gloves, hat & boots) while mixing & loading.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not contaminate any body of water by cleaning of equipment or disposal of waste.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

OKAY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SHEET NO. 2 - 5

ADVERTISING AND SALES PROMOTION SERVICES

30PM

FOR: AGWAY ZINEB-SEVIN^R 4.5-5D

DATE: 1/10/83

STORAGE & DISPOSAL (12 pt.)

FOR CHARACTERS
LEFT CHARACTER

1# Do not reuse empty container. Wrap container and place in trash collection.

This product is formulated to control both insects and diseases. Best results can be obtained by applying before serious build-up occurs. Apply a uniform, thin coating of dust using at the rate of 1 lb. dust per 1,200 sq. ft. (except on cucumbers, squash, melons and pumpkins - use 1 lb. dust per 2,000 sq. ft.). Repeat at 7 to 10 day intervals throughout the growing season (except for beans - repeat at 4 to 10 day intervals). Number of days between treatment & harvest are indicated after crop in ().

BEANS (snap, string, lima) (7) - For control of Anthracnose, downy mildew, rust, cucumber beetles, flea beetles, Japanese beetles, leafhoppers and Mexican bean beetles. Begin application when disease threatens or after first bloom appears.

O'K BY

BROCCOLI (7), BRUSSELS SPROUTS (7), CABBAGE (7), CAULIFLOWER (7) - For control of downy mildew, leaf spots, small cabbage loopers, Diamondback moths, flea beetles, Harlequin bugs, imported cabbageworms and tarnished plant bugs.

SHEET NO. 3 - 5

ACCEPTED
with COMMENTS
by EPA Label Dept.

FEB 11 1983

RECEIVED
FEB 11 1983

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY ZINEB-SEVIN^R 4.5-5D

DATE: 1/10/83

CARROTS (7) - For control of early and late blights, flea beetles and leafhoppers.

PCA - CHARACTERIS
ELITE - CHARACTERIS

CUCUMBERS (5), MELONS (5), including cantaloupes, cassabas, crenshaws, honeydews, muskmelons, Persian melons and watermelons, SQUASH (5) including summer & winter varieties and PUMPKINS (0) - For control of alternaria and angular leaf spots, anthracnose, blossom blight, downy mildew, gummy

stem blight, cucumber beetles, flea beetles, leafhoppers, and squash bugs. Some leaf injury may occur if treatments are made when tender foliage is wet or in the presence of prolonged high humidity.

ENDIVE (10), LETTUCE (10) - For control of downy mildew, small cabbage loopers, flea beetles, Harlequin bugs, Imported Cabbageworms, and leafhoppers.

SPINACH (10) - For control of downy mildew, small cabbage loopers, flea beetles and leafhoppers.

SWEET CORN (0) - For control of Helminthosporium leaf blight, flea beetles, and Japanese beetles. For European Corn borers, dust every 5 to 7 days from the time eggs are about to hatch (for most varieties, this is when corn is knee high). For corn earworms, dust silks as soon as silks appear;

O'K BY

SHEET NO. 4 - 5

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FJR:

AGWAY ZINEB-SEVIN R 4.5-5D

DATE: 1/10/83

Make at least 3 applications at 2 to 3 day intervals. Application of this product to the tassel region of corn during the pollen seed period will seriously reduce bee population.

TOMATOES (5), EGG PLANTS (0), PEPPERS (0) - For control of anthracnose, leaf spots, early & late blight, downy mildew, Colorado potato beetles, flea beetles, leafhoppers, tomato fruitworms and tomato hornworms.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

AGWAY DISCLAIMS ALL IMPLIED WARRANTIES, INCLUDING IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE. BUYER SHALL BE SOLELY RESPONSIBLE FOR ANY AND ALL INJURY, LOSS OR DAMAGE WHICH RESULTS FROM THE USE OF THIS PRODUCT IN ANY MANNER WHICH IS INCONSISTENT WITH THE LABEL DIRECTIONS, WARNINGS OR CAUTIONS. ANY DAMAGES SHALL BE LIMITED TO DIRECT DAMAGES, NOT TO EXCEED THE PURCHASE PRICE OF THE PRODUCT AND SHALL NOT INCLUDE ANY INDIRECT, INCIDENTAL OR CONSEQUENTIAL DAMAGES.

ACCEPTED
AGREEMENTS
DATE: FEB 11 1983

FEB 11 1983

O'K BY

300 886-71
17 1

Agway, Inc.
Chemical Division
P. O. Box 4933
Syracuse, NY 13221

Attention: Mr. J. C. Prinzo

Gentlemen:


Subject: Zineb - Sevin 4.5 - 5D
EPA Registration No. 8590-619
Your Letter Dated January 10, 1983

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted.

1. Add "Do not apply directly to water" to Environmental Hazards Statement.
2. Make sure preharvest intervals are highlighted on the homeowner label.

A stamped copy of the label is enclosed for your records.

Sincerely yours,


Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

TS-767:HMJACOBY:DCR-03334:WANG-2361C:lwy:Raven:479-2013:2/2/83

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							

EPA Form 1320-1 (4-81) OFFICIAL FILE COPY