

ADVERTISING AND SALES PROMOTION SERVICES

FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

((FRONT PANEL))

NET CONTENTS

2 1/2 Gal.

PLATE CHARACTERISTICS  
EPA - SO CHARACTERISTICS

AGWAY  
FLOWABLE MANEB

**ACCEPTED**  
**APR 21 1986**  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for the pesticide  
registered under  
EPA Reg. No. 8590-624

4 Pounds of Maneb Fungicide per gallon.

ACTIVE INGREDIENT:

Maneb (Manganese Ethylenebisdithiocarbamate) 37.0%  
(Total manganese as metallic 7.6%)

INERT INGREDIENTS:

63.0%  
Total 100.0%

KEEP OUT OF REACH OF CHILDREN (12 pt)


CAUTION (18 pt)

See additional precautionary statements on side panel.

AGWAY INC.  
Chemical Division  
Box 4933  
Syracuse, New York 13221

EPA Reg. No. 8590-624  
EPA Est. No. 8590-PA-1

O'K BY



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

((REAR PANEL))

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with eyes, skin or clothing. In case of contact, flush with plenty of water. For eyes, get medical attention.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

If inhaled: remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

If on skin: remove by washing. Get medical attention if irritation persists.

If in eyes: Wash with plenty of water. Get medical attention.

Protective clothing (long pants, long sleeve shirt, gloves, hat and boots) should be worn during mixing and loading.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

PHYSICAL AND CHEMICAL HAZARDS

Keep away from fire and sparks.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL (12 pt)

**PROHIBITIONS:** Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Do not reuse empty container.

**STORAGE:** Store unused product in original container only. Store in a cool, dry area, preferably locked and out of

REGULATORY CHARACTERISTICS

OK BY

.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....  
.....

ADVERTISING AND SALES PROMOTION SERVICES

30611  
BOPV

FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

the reach of children and animals. Not for storage in or around the home. Protect from excess heat. Protect from freezing.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

RE-ENTRY

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written warnings must include the following information:

CAUTION

(Insert area or field description) treated with Maneb on (insert date of application). Do not enter treated areas without protective clothing until sprays have dried. In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Maneb and describe his condition. For further information see the "Statement of Practical Treatment".


Oral warnings must include the same information that is in the written warnings.

DIRECTIONS FOR USE AS A SPRAY

Flowable Maneb is approved for application in sufficient water to provide good coverage with available equipment in either dilute sprays or in concentrate ground or aerial sprays. Rates listed are based on 100 gallons of dilute spray unless otherwise noted. Rates for concentrate and aerial applications should be comparable to the rates for

DO NOT RE-ENTER TREATED AREAS

PKBY



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

dilute sprays on a per acre basis.


Flowable Maneb may be applied by aircraft at the rate shown below for control of certain crop disease; see product labeling for listing of specific crops, details of timing, use limitations, and other use instructions.

CROPS	Amount per Acre	
	Product in Qts.	Water Gals.
Tree Fruits & Nuts	4 to 5.0	10 to 15
Potatoes and most vegetables; Small Fruits; Field Crops	1.2 to 1.6	5 to 8
Tomatoes; Sweet Corn	1.2 to 2.4	5 to 8

Apply with properly calibrated aerial equipment, arrange nozzles so that spray delivery is uniform over the entire spray swath.

AGRICULTURAL FRUIT AND NUT CROP USES

ALMONDS: Brown Rot Blossom and Twig Blight, Leaf Blight, Scab - 1.2 to 1.6 quarts. Apply in popcorn, full bloom, and petal fall or every 7 to 10 days if bloom is staggered. Shotole - 1.2 to 1.6 quarts. Begin in dormant stage.



ADVERTISING AND SALES PROMOTION SERVICES

COPY 5 of 17

FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85


Repeat at 7 to 10 day intervals through petal fall. If applied after petal fall, do not feed hulls to dairy animals or animals being finished for slaughter. Do not apply later than 5 weeks after petal fall.

APPLES: Black Rot, Bitter Rot, Botryosphaeria (White Rot), Bullseye Rot, Cedarapple Rust, Flyspeck, Scab, and Sooty Blotch - 1.2 to 1.6 quarts plus suitable spreader-sticker. Begin in prebloom, calyx, or first cover stages. Repeat at 7 to 14 day intervals as necessary. Do not graze livestock in treated areas. Do not apply within 15 days of harvest in Arkansas, Delaware, Illinois, Indiana, Kansas, Kentucky, Maryland, Missouri, New Jersey, North Carolina, Ohio, Pennsylvania, South Carolina, Tennessee, Virginia and West Virginia. Do not apply within 30 days of harvest in other states.

APRICOTS: Brown Rot (Blossom, Twig and Fruit) - 1.2 to 1.6 quarts. Begin at red bud stage. Repeat in early bloom, full bloom, petal fall, and at 7 to 14 day intervals as necessary up to 2 weeks before harvest. Jacket Rot (Green Rot) 1.6 quarts at full bloom and petal fall.

PICA - 50 CHARACTERS  
ELITE - 60 CHARACTERS

QKBY



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85


Shothole - 1.6 quarts from dormant through petal fall stages in regular schedule. Do not apply later than 14 days before harvest.

CRANBERRIES: Fruit Rots (Massachusetts) - 1.6 to 2.4 quarts. Begin at midbloom. Repeat at 10 to 14 day intervals. Do not apply later than 4 weeks after midbloom. Twig Blight (Lophodermium), (Washington & Oregon) 1.6 quarts per 100 gallons of water. Apply 350 gallons per acre. Begin after blooming. Repeat between July 10 and 20, August 1 and 10, and August 10 and 25. Fruit Rots (Guignardia, Sporonema, and Glomerella) (New Jersey) - 1.6 quarts. Begin after midbloom. Repeat 10 to 14 days later and 4 weeks after midbloom. Do not apply within 30 days of harvest.

PICA - 50 CHARACTERS  
ELITE - 60 CHARACTERS

GRAPES: Black Rot (Bunch Rot) - Apply 1.2 to 4.0 Qt/acre by ground or air starting when new shoots are 8 to 10 inches long (pre-bloom). Repeat after bloom and again 10 days later. When fruit is present use maximum of 1.5 qts per acre. Do not apply within 7 days of harvest.

O'K.BY



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

NECTARINES: Brown Rot, Shothole - 1.2 to 1.6 quarts. Begin at red bud. Repeat in early bloom, full bloom, petal fall, and at 7 to 14 day intervals up to 2 weeks before harvest. Do not apply within 14 days of harvest.

PEACHES: Brown Rot, Scab, Shothole - See under Nectarines. Peach Leaf Curl - 1.2 to 1.6 quarts. Apply in spring dormant period. Do not apply within 2 days of harvest. If applied within 14 days of harvest, remove residues by brushing.

AGRICULTURAL VEGETABLE CROP USES


ASPARAGUS: Rust - 2.4 quarts/50 to 200 gallons/acre. Begin in established fields immediately after harvest. Repeat at 10 day intervals as long as necessary. In young fields which will not be harvested, begin when disease appears and repeat at 10 day intervals. Postharvest application only on established fields.

BEANS (Lima): Anthracnose, Downy Mildew - 1.2 to 1.6 quarts. Begin at first sign of disease. Repeat at 4 to 7 day intervals. Do not apply within 4 days of harvest of succulent beans.

BEANS (Green and Dry): Anthracnose, Downy Mildew,

PICA - 50 CHARACTERS  
ELITE - 60 CHARACTERS

OK BY



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

Rust - 1.2 to 1.6 quarts. Begin when plants are small. Repeat at 5 to 7 day intervals. Do not apply within 4 days of harvest of succulent beans.


BROCCOLI: Alternaria Leaf Spot, Downy Mildew - 1.2 to 1.6 quarts. Begin when diseases threaten. For downy mildew begin in plant bed and repeat at 3 day intervals. In field, repeat at 7 to 14 day intervals. If applied within 3 days of harvest, remove residues by washing or trimming.

BRUSSELS SPROUTS, CAULIFLOWER, AND KOHLRABI: Alternaria Leaf Spot, Downy Mildew - 1.2 quarts. Begin when diseases threaten. Repeat at 3 day intervals in plant beds and at 7 to 10 day intervals in field.

CABBAGE: Alternaria Leaf Spot, Downy Mildew - 1.2 to 1.6 quarts. Begin 7 to 10 days after planting. Repeat at 3 to 5 day intervals in plant bed and field. Do not apply within 7 days of harvest.

PICA - 50 CHARACTERS  
ELITE - 60 CHARACTERS

OKBY





FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

CANTALOUPE, CASSABAS, CRENSHAW, HONEYDEW  
 MELONS, HONEY BALLS, MUSKMELONS, PERSIAN MELONS,  
 AND WATERMELONS: Anthracnose, Cercospora Leaf  
 Spot, Downy Mildew, Gummy Stem Blight - 1.2 to  
 1.6 quarts. Begin at late bunching when vines  
 begin to run or when disease first appears. Re-  
 peat at 7 to 10 day intervals. Do not apply  
 within 5 days of harvest.

CARROTS: Alternaria and Cercospora Leaf Spots -  
 1.2 to 1.6 quarts. Begin when plants are 6  
 weeks old. Repeat at 7 to 10 day intervals.


CASSABA MELONS: See under Cantaloupes.

CAULIFLOWER: See under Brussels Sprouts.

CELERY: Early Blight (Cercospora), Late Blight  
 (Septoria) - 1.2 to 1.6 quarts. Begin when plants  
 emerge in plant beds. Repeat at 3 to 5 day  
 intervals in plant beds and at 7 to 10 day  
 intervals after plants are set in field.  
 Remove excess residues by stripping, trimming,  
 and washing. Do not apply within 14 days of  
 harvest.

PICA - 50 CHARACTERS  
 ELITE - 60 CHARACTERS

OKBY



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

COLLARDS, KALE, MUSTARD GREENS, TURNIPS: Alternaria Leaf Spot and Downy Mildew - Use .8 to 2.4 quarts per acre and start applications when disease first appears in area and repeat at 7 to 10 day intervals. Do not apply within 10 days of harvest. Remove excess residues by washing.

CORN (Sweet): Helminthosporium Blight - 1.2 quarts. Begin when plants are 6 inches high. Repeat at 4 to 7 day intervals until harvest. Do not feed treated forage to livestock.


CRENSHAW MELONS: See under Cantaloupes.

CUCUMBERS: Alternaria (Macrosporium) Leaf Spot, Angular Leaf Spot, Anthracnose, Downy Mildew, Pythium Fruit Rot - 1.2 to 1.6 quarts. Begin when diseases threaten or plants begin to run. Repeat at 7 to 10 day intervals or at 3 to 5 day intervals under severe disease conditions. Do not apply within 5 days of harvest.

EGGPLANT: Anthracnose, Early Blight (Alternaria), Phomopsis Blight or Fruit Rot - 1.2 to 1.6 quarts. Begin when fruit starts to form. Repeat at 7 to 10 day intervals.

PICA - 50 CHARACTERS  
ELITE - 60 CHARACTERS

OK BY



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

ENDIVE, LETTUCE: Downy Mildew - 1.2 to 1.6 quarts. Begin when disease appears. Repeat at 7 to 10 day intervals. Do not apply within 10 days of harvest. Remove residues from head lettuce by stripping and trimming, and from leaf lettuce and endive by washing or other effective means.

HONEY BALL MELONS: See under Cantaloupes.

HONEYDEW MELONS: See under Cantaloupes.

KOHLRABI: See under Brussels Sprouts.

LETTUCE: See under Endive.

MELONS: See under Cantaloupes.


MUSKMELONS: See under Cantaloupes.

ONIONS: Botrytis Leaf Blight (Blast), Downy Mildew, Purple Blotch - 1.2 to 1.6 quarts. Begin at first sign of disease. Repeat at 7 to 10 day intervals.

PEPPERS: Anthracnose, Cercospora Leaf Spot (Frogeye Spot), Ripe Rot - 1.2 to 1.6 quarts.

PICA - 50 CHARACTERS  
ELITE - 60 CHARACTERS

OKBY



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

Begin when disease threatens. Repeat at 7 to 10 day intervals.

PERSIAN MELONS: See under Cantaloupes.

POTATOES: Early and Late Blights - 1.2 to 1.6 quarts. Begin when plants are 2 to 6 inches high. Repeat at 5 to 10 day intervals, more often under severe late blight conditions.


POTATOES (Seed Pieces): Fusarium Seed Piece Decay - 1.0 quart/10 gallons. Dip whole or cut tubers. Spread in cool place if held before planting. Seed piece treatment only. Do not use excess seed pieces for food or feed.

PUMPKINS: Angular Leaf Spot, Downy Mildew - 2.4 quarts/acre in sufficient water to cover or 1.2 to 1.6 quarts/100 gallons. Begin when disease threatens. Repeat at 7 to 10 day intervals.

RHUBARB (Greenhouse (Michigan). Botrytis Leaf Rot - 1.6 quarts. Begin 4 weeks before harvest. Make 4 applications at 7 day intervals.

PICA - 50 CHARACTERS  
ELITE - 60 CHARACTERS

OKBY



ADVERTISING AND SALES PROMOTION SERVICES

COPY 130617

FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

SPINACH: Downy Mildew (Blue Mold), White Rust - 1.2 to 1.6 quarts. Begin when disease appears. Repeat at 7 to 10 day intervals. Do not apply within 10 days of harvest. Remove excess residues by washing.

SQUASH (Summer and Winter): Anthracnose, Downy Mildew - 1.2 to 1.6 quarts. Begin when plants start to run. Repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.

TOMATOES (Greenhouse and Field): Cladosporium Leaf Mold, Early and Late Blights, Gray Leaf Spot (Stemphylium), Septoria Leaf Spot - 1.2 to 1.6 quarts. Begin at first fruit cluster and repeat at 7 to 10 day intervals. Anthracnose - 2.4 quarts. Begin when first fruit forms. Repeat at 7 to 10 day intervals. Do not apply within 5 days of harvest.


WATERMELONS: See under Cantaloupes.

AGRICULTURAL FIELD CROP USES

GRASSES (Ornamental - Grown for Replanting Seed Purposes Only): Leaf, Stem, and Stripe Rust - 1.6 to 2.4 quarts/50 to 100 gallons per acre. Begin when rust pustules are first seen. Repeat

PICA 50 CHARACTER  
ELITE 60 CHARACTER

PKBY



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

at 7 to 14 day intervals. Do not feed treated grass to livestock.


SUGAR BEETS: Leaf Spot (Cercospora) - 1.2 to 2.4 quarts in sufficient water/acre. Begin when disease threatens. Repeat at 7 to 10 day intervals for 3 to 5 applications. Do not apply within 14 days of harvest. Do not feed treated tops to livestock.

TOBACCO: Blue Mold - 0.4 to 3.2 quarts. Use 2.5 to 5.0 gallons of diluted suspension/100 square yards of plant bed. Begin when plants are about the size of a dime or when blue mold is first reported in area. Repeat twice weekly until plants are set in field. Use lower rates in early applications increasing to 3.2 quarts as plants increase in size. For disease control under field conditions, use 1.4 to 2.5 quarts per acre. Repeat as necessary.

AGRICULTURAL ORNAMENTAL CROP USES

CARNATIONS: Alternaria Leaf Spot, Anthracnose, Botrytis Blight - 0.8 to 1.2 quarts. Begin when new growth starts. Repeat weekly.

PICA - 10 CHARACTERS  
ELITE - 10 CHARACTERS



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

▶ CHRYSANTHEMUMS: Ascochyta Ray Blight, Botrytis Petal Spot, Septoria Leaf Spot. Use 12.8 fluid ounces plus 3/4 pound CAPTAN 50W per 100 gallons of water. Apply twice weekly during the blooming season for petal spot and ray blight. For septoria leaf spot apply weekly throughout the season.

◀ PICA 50 CHARACTERS  
ELITE - 60 CHARACTERS


DAHLIAS: Alternaria Leaf Spot, Botrytis Blight - 0.8 to 1.2 quarts. Begin when new growth starts. Repeat weekly.

DOGWOOD: Anthracnose - 0.8 to 1.2 quarts. Begin when buds open. Repeat when bracts fall 4 weeks later and in late summer.

GLADIOLUS: Botrytis Blight, Curvularia and Stem-phylium Leaf Spots - 1.2 quarts. Begin when flower spikes are developing. Repeat 2 to 3 times at weekly intervals.

GRASSES (Nongrazed Areas): Brown Patch - 4.8 fluid ounces in sufficient water/1,000 square feet. Begin at first sign of disease. Repeat at 7 to 14 day intervals. Dollar Spot - 9.6 to 12.8 fluid ounces in sufficient water/1,000 square feet. Apply as for brown patch, above. Melting-Out - 4.8 to 6.4 fluid ounces in sufficient water/

OKAY



FOR: AGWAY FLOWABLE MANEB

DATE: 10/21/85

1,000 square feet. Apply as for brown patch, above. Leaf, Stem and Stripe Rusts - 2.4 quarts/100 gallons/acre. Begin when rust pustules are first seen. Repeat at 7 to 14 day intervals. Do not graze treated areas. Do not feed clippings to livestock.

LILIES: Botrytis Blight - 0.8 to 1.2 quarts. Begin with new growth. Repeat weekly.

PANSIES: Anthracnose - 0.8 to 1.2 quarts. Begin with new growth. Repeat weekly.


PEONIES: Alternaria Leaf Spot, Botrytis Blight, Phytophthora Blight - 0.8 to 1.2 quarts. Apply to foliage and soil in early spring and early fall and 7 to 10 day intervals during the growing season.

ROSES: Black Spot, Cercospora Leaf Spot (Texas) Rust (California) 0.8 to 1.2 quarts. Begin when first leaves unfold. Repeat at 7 to 10 day intervals.

SNAPDRAGONS: Rust - 0.8 to 1.2 quarts. Begin with emergence. Repeat weekly.

PICA - 50 CHARACTERS  
ELITE - 60 CHARACTERS

KEY





ADVERTISING AND SALES PROMOTION SERVICES

COPY 17/17

FOR:

AGWAY FLOWABLE MANEB

DATE: 10/21/85

ZINNIAS: Alternaria Blight, Leaf Spot, Botrytis Blight - 0.8 to 1.2 quarts. Begin with emergence. Repeat weekly.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

PICA 50 CHARACTERS  
ELITE - 60 CHARACTERS

OKBY

