

PM12

8590-625

DEC 16 1983

BEST AVAILABLE COPY

Mr. Julius C. Prinzo
Agway, Inc.
P.O. Box 4741
Syracuse, NY 13221

Dear Mr. Prinzo:

Subject: EFW P.R. Notice of September 23, 1983
EFW-4E
EPA Registration No. 8590-594
EFW-5E
EPA Registration No. 8590-608
EFW-4G
EPA Registration No. 8590-625 ✓

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

However, final printed labels must be revised as noted below, and submitted to us prior to your releasing the product for shipment.

Note, the label for EFW-4G must be revised in the "Environmental Hazards" section to read the same as stated in the Federal Register Notice and as stated on the enclosed accepted labels for EFW-5E and EFW-4E.

Sincerely yours,

Jay S. Ellenberger
Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

DCR-28447:RD-5;12/14/83;KIM:Del.12/26/83

CONCURRENCES

| | | | | | | | |
|-----------|--|--|--|--|--|--|--|
| SYMBOL ▶ | | | | | | | |
| SURNAME ▶ | | | | | | | |
| DATE ▶ | | | | | | | |

FOR:

AGWAY EPN 4G

DATE: 11/9/83

((FRONT PANEL))

RESTRICTED USE PESTICIDE

PICA 28 CHARACTERS
DATE - 10 CHARACTERS

FOR RETAIL SALE TO AND USE ONLY BY CERTIFIED APPLICATORS OR PERSONS UNDER THEIR DIRECT SUPERVISION AND ONLY FOR THOSE USES COVERED BY THE CERTIFIED APPLICATOR'S CERTIFICATION.

AGWAY

EPN 4G

4% Insecticide Granules for the control of European Corn Borer

ACTIVE INGREDIENT:

C-Ethyl (O-p-nitrophenyl) phenylphosphonothioate 4%

INERT INGREDIENTS:

96%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN



DANGER-POISON



PELIGRO: PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

HARMFUL IF SWALLOWED OR INHALED

DO NOT BREATHE DUST

DO NOT GET IN EYES, ON SKIN OR CLOTHING


ACCEPTED
with COMMENTS
In EPA Letter Dated:

DEC 10 1983

Under the Federal Insecticide,
Fungicide and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

2590-10

O'K BY



FOR:

AGWAY EPN 4G

DATE: 11/9/83

((FRONT PANEL))

STATEMENT OF PRACTICAL TREATMENT

If swallowed, call a physician or Poison Control Center.

Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled, remove from exposure. Have patient lie down and keep quiet. If patient is not breathing, start artificial respiration immediately.

If in eyes, wash with plenty of water, get medical attention.

If on skin, immediately remove all contaminated clothing and wash skin thoroughly with soap and warm water. Wash contaminated clothing before reuse.

AGWAY INC.
CHEMICAL DIVISION
P. O. BOX 4933
SYRACUSE, NEW YORK 13221

EPA Reg. No. 8590-625
EPA Est. No. 8590-PA-1

Net Weight 50#

100 - 100 CHARACTERS
100 - 100 CHARACTERS

100 - 100 CHARACTERS
100 - 100 CHARACTERS

.....
O'K BY
.....
.....
.....
.....
.....
.....
.....
.....

FOR:

AGWAY EPN 4G

DATE: 11/9/83

IMPORTANT

PROTECTIVE CLOTHING REQUIRED: Wear clean protective clothing, goggles, and respirator approved by NIOSH or the American National Standards Institute when applying or handling, or when reentering fields within 24 hours of treatment. The following protective clothing must be worn: lightweight, unlined, natural rubber gloves at least midforearm in length; a wide brimmed waterproof hat or waterproof hood; a protective suit or coveralls of a nonpermeable, non-cloth material covering the body from ankles to wrists; lightweight, unlined, natural rubber boots at least midcalf in length; full-face respirators are recommended; half-face respirators and goggles are required. Aerial applicators in positive pressure cockpits and other applicators in comparable ground equipment with appropriate filtered air intakes need not comply with these protective clothing requirements.

Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

THE REGISTRAR

O.K BY

SHEET NO. 3 - 7

SHEET NO. 3 - 7

FOR:

AGWAY EPN 4G

DATE: 11/9/83

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

 **DANGER-POISON** 

Do not breathe dust. Harmful if swallowed or inhaled. Wash hands, arms and face thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use. In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and warm water.


ANTIDOTE AND FIRST AID

ATROPINE IS AN ANTIDOTE. CONSULT PHYSICIAN FOR EMERGENCY SUPPLY OF 1/100 GRAIN ATROPINE TABLETS. CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED POISONING.

If warning symptoms appear (see WARNING SYMPTOMS below), before physician arrives, immediately swallow two atropine tablets (each 1/100 gr); thereafter, every 10 to 15 minutes take one atropine tablet (1/100 gr) until throat becomes dry and skin becomes dry and flushed. Take additional tablets as necessary to maintain a moderately dry throat and dry, flushed skin until physician arrives.

NOTE TO PHYSICIAN

WARNING SYMPTOMS - EPN poisoning produces effects associated with anticholinesterase activity which may include weakness, blurred vision, headache, nausea, abdominal cramps, discomfort in the chest, constriction of pupils, sweating

OK BY 

FOR:

AGWAY EPN 4G

DATE: 11/9/83

slow pulse, muscle tremors.

TREATMENT - Atropine sulfate should be used for treatment. Administer repeated doses, 1.2 to 2.0 mg intravenously every 10 to 30 minutes until full atropinization is achieved. Maintain atropinization until patient recovers. Artificial respiration or oxygen may be necessary. Allow no further exposure to any cholinesterase inhibitor until recovery is assured. 2-PAM may be used to supplement the atropine sulfate treatment. Do not use morphine.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to lakes, streams or ponds. Do not apply to areas where runoff will occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is toxic to bees exposed to direct treatment or residues on crops. Do not apply when bees are actively visiting the area. Protective information may be obtained from your local Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For aerial applications, flagging must be by fully automated mechanical means or by humans working in totally enclosed vehicles.

PICA - 30 CHARACTERS
ELITE - 40 CHARACTERS

.....
O.K. BY
.....
.....
.....
.....
.....
.....

FOR:

AGWAY EPN 4G

DATE:

Agway EPN 4G insecticide is recommended for use on corn (field and sweet) to control European corn borer. Apply at the rate of 5 to 12 1/2 lbs. per acre, using standard granular insecticide application equipment.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide or rinse water that cannot be used according to label instructions must be disposed of according to Federal or approved state procedures under Subtitle C of the Resource Conservation & Recovery Act.

Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

FIELD CORN

Apply 10 days after eggs first hatch or when 75% of the plants show evidence of feeding in the leaf whorls. Repeat 7 to 10 days later if new evidence of leaf feeding is present.

Do not apply within 14 days of harvest.

Do not enter treated fields within 24 hours following treatment.

PCA...
E-178-10 CHARACTER...

O.K. BY

.....
.....
.....
.....
.....
.....
.....
.....
.....
.....

FOR:

AGWAY EPN 4G

DATE: 11/9/83

SWEET CORN

PICA - 10 CHARACTER
ELITE - 8 CHARACTER

Make first application when 25% of the plants show evidence of leaf feeding and continue at 5-day intervals as long as leaf feeding is evident. From 1 to 3 treatments may be required. For control of second brood borers, apply when eggs begin to hatch.

Do not apply within 14 days of harvest.

Do not enter treated fields within 24 hours following treatment.

Do not apply this product when weather conditions favor drift from treated area.

Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

O'K BY
.....
.....
.....
.....
.....
.....
.....
.....
.....
.....