

AGWAY MANEB-SEVIN FLOWABLE

A Water Dispersible Fungicide-Insecticide Combination for Fruit and Vegetable Diseases and Insect Control

ACCEPTED
SEP 14 1982
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 8590-629

ACTIVE INGREDIENTS

Maneb (manganese ethylene bisdithiocarbamate)	37.0%
(total manganese as metallic 7.6%)	
Carbaryl (1-Naphthyl N-methylcarbamate)	24.4%
INERT INGREDIENTS	38.6%
TOTAL	100.0%

Each gallon of product contains 3.8 pounds of Maneb and 2.5 pounds of Sevin

Registered trademark of Union Carbide Corp. for the active ingredient 1-Naphthyl N-Methylcarbamate

STORAGE AND DISPOSAL

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, state or local procedures

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state and local authorities

All recommendations based on 100 gallons of spray unless otherwise indicated. Can be used up to day of harvest unless indicated otherwise

APPLES: Use 1 1/2 qt — For diseases, begin in prebloom, Calyx or first cover stages. Repeat at 7 to 14 day intervals. For insects, begin at first sign of insects or insects' feeding. Black rot, bitter rot, Botryosphaeria (White rot) bullseye rot, Cedar apple rust, flyspeck, scab, sooty blotch, Apple Mealybug, codling moth, green apple aphid, white apple leathopper, aphids (rosy apple, woolly apple), apple maggot, bagworm, eastern tent caterpillar, European apple sawfly, eye-spotted bud moth, fruit tree leaf roller, green fruitworm, Japanese beetle, mites (apple rust, pear leaf blister, pear rust), pear psylla, periodical cicada, plum curculio, red banded leaf roller, scales (forbes, lecanium, oystershell, San Jose), tarnished plant bug, tentiform leaf miner. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. When heavy aphid infestations are present, repeated applications may be necessary for optimum control. To control rosy apple aphids, apply before leaves are curled. For optimum scale control, apply when crawlers are present. Apply at 10 day intervals for apple maggot control. Applications within 30 days after full bloom may also provide apple thinning. To avoid this, delay use until at least 30 days after bloom. May cause foliar injury if used before second cover on York and McIntosh apples. Do not apply within 30 days of harvest

BEANS (snap or string): Use 1 1/2 qt — For anthracnose, downy mildew, Mexican bean beetle, Bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leathoppers, velvet bean caterpillar, western bean cutworm. Begin when plants are small. Repeat at 4 to 7 day intervals. Do not apply within 4 days of harvest

COLE CROPS (BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI): Use 1 1/2 qt — For alternaria leaf spot, downy mildew, flea beetles, harlequin bug, Armyworms, imported cabbage worms, diamondback larvae and loopers, corn earworm. Repeat at 5 to 10 day intervals. Do not apply within 3 days of harvest for broccoli, 7 days for cabbage.

CANTALOUPE, CASSABAS, CRENSHAW'S, HONEYDEW MELONS, HONEY BALLS, MUSKMELONS, PERSIAN MELONS, and WATER-MELONS: Use 1 1/2 qt — Anthracnose, Cercospora Leaf Spot, Downy Mildew, Gummy Stem Blight, Melonworm, pickleworm, Cucumber beetles, flea beetles, leathoppers, squash bugs. Avoid excessive application. To

avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Do not apply within 5 days of harvest

CUCUMBERS, PUMPKINS, SQUASH (Summer and Winter): Use 1 1/2 qt — Alternaria leaf spot, angular leaf spot, anthracnose, downy mildew, cucumber beetles, flea beetles, leathoppers, and squash bugs. Begin when diseases or insects appear or when plants start to run. Repeat at 7 to 10 day intervals or under severe pressure every 3 to 5 days. Do not apply within 5 days of harvest for cucumbers and squash

ENDIVE, LETTUCE: Use 1 1/2 qt — Downy mildew, flea beetles, Harlequin bugs, leathoppers (including six-spotted leathoppers (aster yellows vector)). Do not apply within 7 days of harvest. Remove residues from head lettuce by stripping and trimming, and from leaf lettuce and endive by washing. Do not apply within 10 days of harvest

POTATOES, TOMATOES, EGGPLANTS and PEPPERS: Use 1 1/2 qt — Anthracnose, Early and late blight, Septoria leaf spot, Colorado potato beetle, flea beetles, leathoppers, tomato hornworm. Begin when fruit starts to form or earlier if diseases or insects appear. Repeat at 5 to 10 day intervals, more frequently under severe blight conditions. Do not apply within 5 days of harvest for tomatoes

ORNAMENTAL PLANTS, FLOWERS, SHRUBS and SHADE TREES: ROSE, DAHLIA, LILIES, PANSIES, PEONIES, SNAPDRAGONS, CARNATIUM, GLADIOLUS, ZINNIA, CHRYSANTHEMUM, LILAC, ARBOR VITAE, JUNIPER HYDRANGEA, AZALEA, ELM, MAPLE, OAK, DOGWOOD, BIRCH and PINES (Do not use on Boston Ivy): Apply 1 qt per 100 gallons water when diseases, insects or their damage appear. Repeat weekly or as needed to control. Alternaria leaf spot, anthracnose, black spot, leaf blights, rust, Apple aphid, bagworm, birch leaf miner, blister beetles, boxelder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetles, lace bugs, leathoppers, leaf rollers, mealybugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, roseslug, scale insects, tent caterpillars, thorn bugs, thrips (exposed), willow leaf beetles.

NOTE: This pesticide is to be sold ONLY in this original unbroken package

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith

EPA Reg No 8590-629

EPA Est No. 8590-PA-1

AGWAY INC. — CHEMICAL DIVISION
PO BOX 4933
SYRACUSE, NY 13221

NET CONTENTS 2 1/2 GAL.

A 7/82

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person

If inhaled, remove victim to fresh air. Apply artificial respiration, if indicated

If on skin, remove contaminated clothing and wash affected areas with soap and water

If in eyes, flush eyes with plenty of water. Call a physician if irritation persists

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

BEE CAUTION: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to place hives beyond bee flight range until 1 week after application or to take other equally effective precautions

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

