

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY MANEB-SEVIN [®] FLOWABLE

DATE: 1/6/83

((FRONT PANEL))

NET CONTENTS

.Gal., 2½ Gal.

AGWAY
MANEB-SEVIN [®] FLOWABLE

A water dispersible fungicide-insecticide combination for fruit and vegetable diseases and insect control.

ACTIVE INGREDIENTS:

Maneb (manganese ethylene bisdithiocarbamate)

(total manganese as metallic 7.6%)

Carbaryl (1-Naphthyl N-methylcarbamate) 24.4%

INERT INGREDIENTS 38.6%

TOTAL 100.0%

Each gallon of product contains 3.8 pounds of Maneb and 2.5 pounds of Sevin.

[®]Registered trademark of Union Carbide Corp. for the active ingredient 1-Naphthyl N-Methylcarbamate.

KEEP OUT OF REACH OF CHILDREN (12 pt.)

CAUTION (18 pt.)

See additional precautionary statements on side panel.

AGWAY INC.
CHEMICAL DIVISION
BOX 4933
SYRACUSE, NY 13221

EPA Reg. No. 8590-629
EPA Est. No. 8590-PA-1

FEB 11 1983
8590-629

O'K BY

SHEET NO. 1 - 8

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY MANEB=SEVIN [®] FLOWABLE

DATE: 1/6/83

((REAR PANEL))

FOR: MANEB=SEVIN
DATE: 01/06/83

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, Drink 1 or 2 glasses of water & induce vomiting by touching back of throat with finger. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled, remove victim to fresh air. Apply artificial respiration, if indicated.

If on skin, remove contaminated clothing & wash affected areas with soap & water.

If in eyes, flush eyes with plenty of water. Call a physician if irritation persists.

Wear protective clothing (long pants, long sleeve shirt, gloves, hat & boots) during mixing & loading.

O'K BY

SHEET NO. 2 - 8

ADVERTISING AND SALES PROMOTION SERVICES COPY

FOR: AGWAY MANEB-SEVIN® FLOWABLE

DATE:1/6/83

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

BEE CAUTION: This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until 1 week after application or to take other equally effective precautions.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL (12 pt.)

2½ Gal:

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, state or local procedures.

Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by incineration if allowed by state & local authorities.

✓

OKBY

ADVERTISING AND SALES PROMOTION SERVICES

30PV

FOR: AGWAY MANEB-SEVIN [®] FLOWABLE

DATE: 1/6/83

Gal:

Do not reuse empty container. Wrap container and put in trash collection.

All recommendations based on 100 gallons of spray unless otherwise indicated. Can be used up to day of harvest unless indicated otherwise.

Apples: Use 1½ qt. for diseases, begin in pre-bloom, Calyx or first cover stages. Repeat at 7 to 14 day intervals. For insects, begin at first sign of insects or insects' feeding. Black rot, bitter rot, Botryosphaeria (White rot), bullseye rot, Cedar-apple rust, flyspeck, scab, sooty blotch, Apple Mealybug, codling moth, green apple aphid, white apple leafhopper, aphids (rosy apple, woolly apple), apple maggot, bagworm, eastern tent caterpillar, European apple sawfly, eye-spotted bud moth, fruit tree leaf roller, green fruitworm, Japanese beetle, mites (apple rust, pear leaf blister, pear rust), pear psylla, periodical cicada, plum curculio, red-banded leaf roller, scales (forbes, lecanium, oyster-shell, San Jose), tarnished plant bug, tentiform leaf miner. Apply at petal fall and every 10 to 14 days thereafter until control is achieved. When heavy aphid infestations are present, repeated

100-30000000

O'K BY

SHEET NO. 4 - 8

ADVERTISING AND SALES PROMOTION SERVICES

300PY

FOR:

AGWAY MANEB-SEVIN® FLOWABLE

DATE:1/6/83

applications may be necessary for optimum control. To control rosy apple aphids, apply before leaves are curled. For optimum scale control, apply when crawlers are present. Apply at 10 day intervals for apple maggot control. Applications within 30 days after full bloom may also provide apple thinning. To avoid this, delay use until at least 30 days after bloom. May cause foliar injury if used before second cover on York and McIntosh apples. Do not apply within 30 days of harvest.

Beans (snap or string): Use 1½ qt. - For Anthracnose, downy mildew, Mexican bean beetle, Bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvet bean caterpillar, western bean cutworm.

Begin when plants are small. Repeat at 4 to 7 day intervals. Do not apply within 4 days of harvest.

COLE CROPS (BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, KOHLRABI): Use 1½ qt. - For alternaria leaf spot, downy mildew, flea beetles, harlequin bug, Armyworms, imported cabbage worms, diamondback larvae and loopers, corn earworm. Repeat at 5 to 10 day intervals.

OK BY

SHEET NO. 5 - 8

ADVERTISING AND SALES PROMOTION SERVICES

30PV

FOR:

AGWAY MANEB-SEVIN® FLOWABLE

DATE: 1/6/83

Do not apply within 3 days of harvest for
broccoli, 7 days for cabbage.

CANTALOUPE, CASSABAS, CRENSHAW, HONEYDEW MELONS,
HONEY BALLS, MUSKMELONS, PERSIAN MELONS, AND

WATERMELONS: Use 1½ qt. - Anthracnose, Cercospora
Leaf Spot, Downy Mildew, Gummy Stem Blight, Melon-
worm, pickleworm, Cucumber beetles, flea beetles,
leafhoppers, squash bugs, Avoid excessive appli-
cation. To avoid possible injury on tender foliage,
do not apply when foliage is wet or when rain or
excessive humidity is expected during the next 2
days. Do not apply within 5 days of harvest.

CUCUMBERS, PUMPKINS, SQUASH (SUMMER & WINTER):
Use 1½ qt. - Alternaria leaf spot, angular leaf
spot, anthracnose, downy mildew, cucumber beetles,
flea beetles, leafhoppers, and squash bugs. Begin
when diseases or insects appear or when plants start to
run. Repeat at 7 to 10 day intervals or under severe
pressure every 3 to 5 days. Do not apply within
5 days of harvest for cucumbers & squash.

ENDIVE, LETTUCE: Use 1½ qt. - Downy mildew, flea
beetles, Harlequin bugs, leafhoppers (including six-
spotted leafhopper (aster yellows vector)). Do
not apply within 7 days of harvest. Remove residues
from head lettuce by stripping & trimming, and from
leaf lettuce & endive by washing. Do not apply
within 10 days of harvest.

OK BY

SHEET NO 6 - 8

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY MANEB-SEVIN [®] FLOWABLE

DATE: 1/6/83

POTATOES, TOMATOES, EGGPLANTS, & PEPPERS: Use 1½ qt. - Anthracnose, Early & late blight, Septoria leaf spot, Colorado potato beetle, flea beetles, leafhoppers, tomato hornworm. Begin when fruit starts to form or earlier if diseases or insects appear. Repeat at 5 to 10 day intervals, more frequently under severe blight conditions. Do not apply within 5 days of harvest for tomatoes.

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES: ROSE, DAHLIA, LILIES, PANSIES, PEONIES, SNAPDRAGONS, CARNATION, GLADIOLUS, ZINNIA, CHRYSANTHEMUM, LILAC, ARBOR VITAE, JUNIPER HYDRANGEA, AZALEA, ELM, MAPLE, OAK, DOGWOOD, BIRCH AND PINES. (Do not use on Boston Ivy) - Apply 1 qt. per 100 gallons water when diseases, insects or their damage appear. Repeat weekly or as needed to control: Alternaria leaf spot, anthracnose, black spot, leaf blights, rust, Apple aphid, bagworm, birch leaf miner, blister beetles, box-elder bug, boxwood leaf miner, elm leaf aphid, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf rollers, mealybugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, roseslug,

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY MANEB-SEVIN [®] FLOWABLE

DATE: 1/6/83

scale insects, tent caterpillars, thorn bugs,
thrips (exposed) willow leaf beetles.

2 CA. 10 CHARACTER.
10 TE. 10 CHARACTER.

NOTE: This pesticide is to be sold ONLY in this
original unbroken package.

DISCLAIMER: Buyer assumes all risks of use,
storage or handling not in strict accordance with
directions given herewith.

O'K BY

SHEET NO. 8 - 8

17 33633

Agway, Inc.
Chemical Division
P. O. Box 4933
Syracuse, NY 13221

Attention: Mr. J. C. Prinzo

Gentlemen:

Subject: Maneb - Sevin Flowable
EPA Registration No. 8590-629
Your letter dated January 10, 1983

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished labeling must be submitted.

1. For homeowner use, on container, the preharvest intervals must be highlighted.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure

TS-767:HMJACOBY:DCR-03334:WANG-2361C:lwy:Raven:479-2013:2/2/83

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							