

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR: AGWAY LAWN & GARDEN MALATHION 50

DATE: 7/19/82

((FRONT PANEL))

NET CONTENTS
8 oz., pt., qt.

ALL CHARACTERS
ELITE-40 CHARACTERS

AGWAY

LAWN & GARDEN MALATHION 50

FOR HOME GARDEN USE ONLY

Kills aphids, mealybugs and scale insects on
fruits, vegetables & ornamentals. Also kills
mosquitoes and ants.

ACTIVE INGREDIENTS:

*Malathion 50%

Xylene Range Aromatic Solvent 40%

INERT INGREDIENTS 10%

TOTAL 100%

*O,O-dimethyl dithiophosphate of diethyl mercap-
tosuccinate.

ACCEPTED

SEP 29 1982

Under Federal Insecticide,
Fungicide, and Plant Fertilizer Act
as amended for two pesticide
registrations under
EPA Reg. No. 8590-632

KEEP OUT OF REACH OF CHILDREN (12 pt.)

CAUTION (1B pt.)

See additional precautionary statements on rear
panel.

AGWAY INC.
CHEMICAL DIVISION
BOX 4933
SYRACUSE, NY 13221

EPA Reg. No. 8590
EPA Est. No. 8590-PA-1

OKBY

SHEET NO. 1 - 12

BEST DOCUMENT AVAILABLE

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY LAWN & GARDEN MALATHION 50

DATE:7/19/82

((REAR PANEL))

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: Harmful if swallowed. Do not breathe spray mist. Do not get in eyes or on skin.

STATEMENT OF PRACTICAL TREATMENT

If swallowed, drink 1 or 2 glasses of water & induce vomiting by touching back of throat with finger. Call a physician immediately. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled, remove victim to fresh air. Apply artificial respiration, if indicated.

If on skin, remove contaminated clothing & wash affected areas with soap & water.

If in eyes, flush eyes with plenty of water. Call a physician immediately.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

PICA 40 CHARACTERS
ELITE - 80 CHARACTERS

OK BY

SHEET NO. 2 - 12

BEST DOCUMENT AVAILABLE

ADVERTISING AND SALES PROMOTION SERVICES COPY

FOR: AGWAY LAWN & GARDEN MALATHION 50

DATE: 7/19/82

This product is highly toxic to bees exposed to direct treatment or residues on crop. Protective information may be obtained from your cooperative Agricultural Extension Service.

PCA - 28 CHARACTERS
SLITS - 28 CHARACTERS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL (12 pt.)

Do not reuse empty container. Wrap container and place in trash collection.

Apply as a spray using 2 teaspoons per gallon of water (unless indicated otherwise). Spray entire plant thoroughly, including lower leaf surfaces. Apply as insects appear; repeat as needed.

Ornamentals (flowers, shrubs, shade trees)
Oak Kermes, oyster shell scale, Fletcher Scale, Florida Red Scale, Juniper Scale, (apply when scale crawlers have settled on foliage); Azalea scale, birch leaf miner, boxwood leaf miner, Euonymus scale, European pine shoot moth, four-lined leaf bug, adult Japanese beetle, Magnolia scale, pine leaf scale, potato leafhopper, rose

OK BY

SHEET NO. 3 - 12

BEST DOCUMENT AVAILABLE

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY LAWN & GARDEN MALATHION 50

DATE: 7/19/82

leafhopper, scurfy scale, tarnished plant bug, thrips, tent caterpillar, aphids, mealybugs, spider mites, whitefly, bagworm, lace bug, black scale crawlers, Monterey pine scale, soft brown scale. For wax scale & pine needle scale use 4 teaspoons (apply in spring when crawlers are active. Repeat 1 or 2 full-coverage applications at 10 day intervals.

Injury may occur on ferns, hickory, Viburnum, Lantana, crassula & canariti juniper. Slight injury may occur on Boston, pteris & Maidenhair ferns, petunias, small-leaf spirea, white pine, maple & elm.

BERRIES

Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries; Japanese beetles, leafhoppers, mites, thrips; for aphids and rose scale use 3 teaspoons. Do not apply within 1 day of harvest.

Blueberries; Japanese beetles. Do not apply within 1 day of harvest.

WCA - MECHANICALS
RITE - MECHANICALS

OKBY

SHEET NO. 4 - 12

BEST DOCUMENT AVAILABLE

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY LAWN & GARDEN MALATHION 50

DATE: 7/19/82

Strawberries; aphids, spider mites, potato leafhoppers, strawberry leafroller, strawberry root weevil, whitefly. For field cricket, lygus bug, spittlebug and thrips use 3 teaspoons. Do not apply within 3 days of harvest.

PICA 50 CHARACTERS
ELITE 100 CHARACTERS

Melons, including cantaloupe, casabas, crenshaw, honeydew, muskmelons, persian melons (and hybrids) and watermelons (and hybrids); aphids, cucumber beetle, spider mites. Do not apply within 1 day of harvest.

FRUIT

Apples; bud moth, Woolly apple aphid, green apple aphid, rosy apple aphid, Forbes scale, mealybug, Codling moth, plum curculio, red-banded leaf folder, tent caterpillar. May injure McIntosh & Cortland Varieties. Do not apply within 3 days of harvest.

Apricots; aphids, codling moth, orange tortrix, soft brown scale, Terrapin scale. Do not apply within 7 days of harvest.

O.K. BY

Cherries; black cherry aphid, bud moth, cherry fruit Fly, Japanese beetle, fruit tree leaf roller. Do not apply within 3 days of harvest.

SHEET NO. 5 - 12

BEST DOCUMENT AVAILABLE

ADVERTISING AND SALES PROMOTION SERVICES COPY

FOR: AGWAY LAWN & GARDEN MALATHION 50

DATE: 7/19/82

Grapes; European fruit lecanium, mealybug, spider mites, Terrapin scale, leafhoppers. Do not apply within 3 days of harvest.

Nectarines; Mites, plum curculio. Product may cause fruit spotting. Do not apply within 7 days of harvest.

Peaches; black cherry aphid, black peach aphid, green-peach aphid, Japanese beetle, rusty plum aphids, cottony peach scale, Terrapin scale, Oriental fruit moth, plum curculio. Do not apply within 7 days of harvest.

Pears; Mealybug, mites, pear psylla, Codling Moth, fruit tree leaf roller, plum curculio, red banded leaf roller. May cause injury on Bos pears under certain conditions in the Northeast. Do not apply within 1 day of harvest.

Plums; Mealy plum aphid. Do not apply within 3 days of harvest.

Prunes; Mealy plum aphid, bud moth. Do not apply within 3 days of harvest.

PICK - 14 CHARACTERS
PLUTE - 14 CHARACTERS

O'K B'

SHEET NO. 6 - 12

BEST DOCUMENT AVAILABLE

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY LAWN & GARDEN MALATHION 50

DATE: 7/19/82

VEGETABLES

Asparagus; asparagus aphid, asparagus beetle, thrip. Do not apply within 1 day of harvest.

Beans (green, lima, navy, red kidney, snap, wax, cowpeas & black-eyed peas); aphids, cucumber beetles, Japanese beetle, mites, Mexican bean beetle. Do not apply within 1 day of harvest.

Beets (table & garden); aphids. Do not apply within 7 days of harvest.

Broccoli; aphids. Do not apply within 3 days of harvest.

Brussels sprouts; aphids. Do not apply within 7 days of harvest.

Cabbage; aphids, cabbage looper, imported cabbageworm. Do not apply within 7 days of harvest.

Carrots; aphids, leafhoppers. Do not apply within 7 days of harvest.

Cauliflower; caterpillars. Do not apply within 7 days of harvest.

100 CHARACTERS

ELITE 20 CHARACTERS

OK BY

SHEET NO. 7 - 12

BEST DOCUMENT AVAILABLE

ADVERTISING AND SALES PROMOTION SERVICES COPY

FOR: AGWAY LAWN & GARDEN MALATHION 50

DATE: 7/19/82

Celery; aphids, spider mites. Do not apply within 7 days of harvest.

PCB 32 CHARACTERS
FILE 40 CHARACTERS

Collards; aphids, Caterpillars, Harlequin cabbage bug. Do not apply within 7 days of harvest.

Corn (field, sweet & pop); aphids, armyworms, cereal leaf beetle, corn rootworm adults, grasshoppers, sap beetles, thrips. Do not apply within 5 days of harvest.

Cucumbers; aphids, pickleworm, spider mites. For cucumber beetles use 3 teaspoons. Do not apply within 1 day of harvest. Do not apply unless plants are dry.

Eggplant; aphids, spider mites. For lace bugs use 3 teaspoons. Do not apply within 3 days of harvest.

OK BY

Endive (escarole); aphids, mites. Do not apply within 7 days of harvest.

Garlic; aphids, thrips. Do not apply within 3 days of harvest.

SHEET NO. 8 - 12

BEST DOCUMENT AVAILABLE

ADVERTISING AND SALES PROMOTION SERVICES **COPY**

FOR: AGWAY LAWN & GARDEN MALATHION 50

DATE: 7/19/82

Parsley; aphids. Do not apply within 21 days of harvest.

Peas; aphids, pea weevil. Do not apply within 7 days of harvest if vines are used as animal feed, 3 days if vines are not used as animal feed.

Peppers; aphids, maggots. Do not apply within 3 days of harvest.

Potatoes; aphids, false chinch bugs, leafhoppers, mealybugs. Do not apply within 3 days of harvest.

Pumpkins; aphids, Mites. For squash vine borer use 3 teaspoons. Do not apply within 3 days of harvest. Do not apply unless plants are dry.

Rutabagas; aphids. Do not apply within 3 days of harvest.

Spinach; aphids. Do not apply within 7 days of harvest.

Squash; aphids, pickleworm, spider mites. For squash vine borer & cucumber beetles use 3 teaspoons. Do not apply within 1 day of harvest.

PCA - GENERAL USE
ELITE - CHARACTER

OKAY

SHEET NO. 10 - 12

BEST DOCUMENT AVAILABLE

ADVERTISING AND SALES PROMOTION SERVICES COPY

FOR:

AGWAY LAWN & GARDEN MALATHION 50

DATE:

Sweet potatoes; leafhoppers. For Morning-glory leafminers use 2 to 3 teaspoons. Do not apply within 3 days of harvest.

FIG. 10 CHARACTERS
LINE - 40 CHARACTERS

Tomatoes; aphids, Drosophila (fruit flies), spider mites. Do not apply within 1 day of harvest.

Turnips; aphids, cabbage looper, imported cabbage-worm. Do not apply within 3 days of harvest.

ANTS

Mix 2 tablespoons in 1 gallon of water & thoroughly saturate ant mounds & mix moist earth on top of hill. For other small ants in flower beds, lawns & around trees, spray the infested area lightly. Repeat in 10 to 15 days if ants return.

DOG & CAT PESTS

Fleas & lice; using a hand sprayer, wet the animal thoroughly with a spray composed of 1 oz. (2 tablespoons) of product in 1 gallon of water. If necessary, repeat treatment in 7 days.

OK BY

If the animal is confined to an area which is free of fleas, one treatment may suffice. If, however, the animal is allowed to run free, it may become

SHEET NO. 11 - 12

BEST DOCUMENT AVAILABLE

ADVERTISING AND SALES PROMOTION SERVICES

COPY

FOR:

AGWAY LAWN & GARDEN MALATHION 50

DATE: 7/19/82

re-infested with fleas. Animals that run free should be re-examined closely every three weeks or so and be retreated if fleas are present.

Animal quarters: Animals become reinfested unless areas such as kennels, pens, yards, under houses or porches & lawn areas are also treated. Remove manure or debris from these areas and spray using 5 oz. (10 tablespoons) product in 1 gallon of water & apply at the rate of 1 gallon of mixture per each 1,000 sq. ft. using a hand or garden sprayer. Do not repeat application for 7 days and then only if infestation is present.

NOTE: This pesticide is to be sold ONLY in this original unbroken package.

DISCLAIMER: Buyer assumes all risks of use, storage or handling not in strict accordance with directions given herewith.

ALL INFORMATION CONTAINED
HEREIN IS UNCLASSIFIED

OK BY

SHEET NO. 12 - 12

BEST DOCUMENT AVAILABLE