

8593-1

PM-15

1/4/70

13/1

DIRECTIONS: Read entire instructions and observe precautions before using. Apply a small amount of SCALEX with a cotton swab to the scaly and crusty areas on the bird's legs, feet and face. Repeat on non feathered areas every 5 to 7 days. On feathered areas use sparingly every 7-10 days. Do not soak feathers. Repeat treatments 6 times or as otherwise directed by your veterinarian.

To help prevent reinfestation use 2 or 3 times a month. **IMPORTANT** -- Be sure the perches are kept clean. Rub the perches with SCALES about twice a month to discourage the presence of mites. Do not use if bird chews perches.

CAUTION -- Keep SCALEX out of bird's eyes, mouth, nostrils and vent. SCALEX should only be used by adults experienced in handling and treating birds. Use under the supervision of a veterinarian. Do not use on sick or debilitated birds. Keep out of reach of children and pets. Harmful if swallowed. Avoid human skin contact. Use rubber gloves when applying product.

THE ARLIS COMPANY

SCALEX

For external use on Birds
for
Scaly Face and Scaly Leg Mites

Keep out of Reach of Children
and Pets

CAUTION

Harmful if Swallowed

CONTENTS: 28 ml. (0.95 FL.OZ.)

Manufactured by
The ARLIS Company
Division of RICH HEALTH
P.O. Box 18258, Irvine, CA 92713

ACTIVE INGREDIENTS:

Thymol (5-Methyl-2-isopropylphenol) 0.55%
Betanaphthol (Betahydroxynaphthalene) ... 0.45%
Naphthalene (Tar Camphor) 0.50%
INERT INGREDIENTS 98.50%

EPA Est. No. 43576-CA-01 EPA Reg. No. 8593-1

CAUTION: Flammable mixture. Do not use near heat or open flame. Keep container closed when not in use.

PRODUCT NO. 35095

SCALEX has been used since 1964. For use on Parakeets (Budgies), Canaries, Love Birds, Cockatiels, Finches.

Not Reviewed. Registrant claims to be in accordance with registration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

