



# FEED TREET<sup>T.M.</sup>

FOR THE SOLE CONTROL OF MOLD AND FUNGI GROWTH

**ACTIVE INGREDIENTS:** Propionic Acid

**INERT INGREDIENTS:** 92%

Sodium Chloride, Propyl p-Hydroxybenzoate, Lactic Acid, Ethyl Methylphenylglycidate, Calcium Silicate, Ethyl Orthoaminobenzoate, Ascorbyl Palmitate.

E.P.A. Registration No. 8596-9

NET CONTENTS \_\_\_\_\_ LBS.

## CAUTION

KEEP OUT OF REACH OF CHILDREN


See other precautions on the side panel

KEMIN INDUSTRIES  
DES MOINES, IOWA 50301 U.S.A.

ACCEPTED

*January 17 1973*

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTRATION  
ED UNDER NO. 8596-9



# FEED TREET<sup>T.M.</sup>

## DIRECTIONS

A mold inhibitor for mixed feeds. Controls mold, heat, bridging and setting up problems in all types of mixed feeds. Keeps all feed free flowing, improves palatability, nutrient retention, digestibility and color of all mixed feeds.

Use 5 pounds per ton of feed grain and mix thoroughly. May be premixed with 10 pounds of a suitable diluent, such as soybean meal or finely ground grain. Mix this 15 pound premix thoroughly in a ton of feed grain.

**Moisture levels:** Maximum moisture content of feed is 20%. If feed moisture is above 20% the feed must be dried to a safe moisture level.

**CAUTION:** Harmful if swallowed. Causes skin irritation. Avoid contact with skin and eyes. In case of contact, flush skin or eyes with plenty of water for at least 15 minutes. For eyes get medical attention. Avoid contamination of foods. Do not reuse container. Destroy when empty.

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the face hereof. All statements made concerning this product apply only when used as directed.

PRINTED IN U.S.A. ©KEMIN INDUSTRIES, INC. 1972 PRODUCT NO. 07-3877 720911

TAKE TIME



OBSERVE LABEL  
DIRECTIONS

KEMIN INDUSTRIES  
DES MOINES, IOWA 50301 U.S.A.