


KEM SAN™
PRODUCT 05006



ACTIVE INGREDIENT	
Propionic Acid	30.5%
INERT INGREDIENTS	31.5%
TOTAL	100.0%

For the sole control of bacteria in processed drinking water for livestock and poultry, and mold and fungi in poultry litter, feed, silage and grain storage.

KEEP OUT OF REACH OF CHILDREN
DANGER

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

Corrosive, causes irreversible eye damage. Do not get in eyes. Wear goggles, face shield, or safety glasses. Avoid contact with skin, eyes or clothing. May be harmful if swallowed or absorbed through the skin. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Environmental Hazards

Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment wash waters.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Immediately flush with plenty of water for at least 15 minutes. Call a physician. If on skin: Wash with plenty of soap and water. Get medical attention. Remove contaminated clothing and shoes at once; wash thoroughly before reuse. If swallowed: Do not induce vomiting. Drink large quantities of water. Obtain medical attention immediately.

WARRANTIES

The manufacturer makes no warranties, expressed or implied, concerning this product or its use, which extend beyond the description on the face hereof. All statements made concerning this product apply only when used as directed.

PRODUCT COLOR CHANGE WILL NOT AFFECT PERFORMANCE.

NOTE:

KEM SAN™ is not a substitute for good management.

EPA Registration No. 8596-

EPA Est. 8595-IA-01

NET WT. 50 lbs (22.68 kg)

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

Do not contaminate water or food by storage and disposal.

PESTICIDE STORAGE: Store in a warm, dry place above 40° F to prevent freezing.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DRINKING WATER: Use as an aid in the reduction of bacteria in processed drinking water for livestock and poultry. Use 1/3 to 1-1/2 fluid ounces per gallon of water. Use 1/3 fluid ounce under normal conditions. Use 1-1/2 fluid ounces when water contains high bacterial contamination.

POULTRY LITTER: Use 6 fluid ounces in one gallon of water. Apply this solution to 1000 square feet of litter.


FEED: Use 2 pounds per ton. When used in feed, do not dilute.

SILAGE STORAGE: Remove all old silage and dirt from all interior surfaces of the storage areas. Mix 6 fluid ounces in one gallon of water. Apply this solution to 1000 square feet of surface storage areas.

GRAIN STORAGE: Remove all old grain and dirt from all interior surfaces of the storage areas. Mix 6 fluid ounces in one gallon of water. Apply this solution to 1000 square feet of surface storage areas.

™Trademark Kemin Industries, Inc. U.S.A.

Printed in U.S.A. © Kemin Industries, Inc. U.S.A., 1990 900412



BEST AVAILABLE COPY