

ACCEPTED



DIRECTIONS FOR USE

APPLICATIONS TO THE SOIL SHOULD BE WORKED INTO THE TOP FEW INCHES IMMEDIATELY AFTER APPLICATION WITH A ROTARY HOE, DISK HARROW, CULTIVAR, HARRAW, OR OTHER SUITABLE MEANS FOR GREATEST EFFECTIVENESS.

FIELD AND FORAGE INSECTS

BEANS (PINTO) LESSER CORNSTALK BORERS: APPLY 7-14 LBS. PER ACRE IN A 6-INCH BAND CENTERED OVER THE ROW WHEN BEANS ARE 1/2-1 INCH HIGH.

PEANUTS: SOUTHERN CORN POOTWORMS: BROADCAST 28 LBS. PER ACRE OR THE EQUIVALENT IN A BAND OVER THE FRUITING ZONE, JUST PRIOR TO PEGGING. WORK INTO TOP FEW INCHES OF SOIL.

CUTWORMS: APPLY 14 LBS. PER ACRE IN A 12-INCH BAND OVER THE ROW WHEN CUTWORMS FIRST APPEAR. UP TO THREE ADDITIONAL APPLICATIONS AT THREE-WEEK INTERVALS MAY BE NECESSARY. DO NOT APPLY WITHIN 21 DAYS OF PEANUT HARVEST.

LESSER CORNSTALK BORERS: APPLY 14-21 LBS. PER ACRE IN A BAND OVER THE FRUITING ZONE JUST PRIOR TO PEGGING. COVER THE BAND WITH 1-2 INCHES OF SOIL.

NOTE: LIVESTOCK (1) MAY BE FED PEANUT HULLS (2) MAY BE FED PEANUT FORAGE 7 DAYS FOLLOWING TREATMENT AND (3) MAY BE FED PEANUT HAY 21 DAYS FOLLOWING TREATMENT.

SORGHUM: CUTWORMS: BROADCAST 14-28 LBS. PER ACRE JUST PRIOR TO PLANTING. WORK INTO SOIL 2-3 INCHES TO CONTROL SURFACE CUTWORMS; 3-6 INCHES TO CONTROL SUBTERRANEAN CUTWORMS.

LESSER CORNSTALK BORERS: APPLY 7-14 LBS. PER ACRE IN A 10-INCH BAND OVER THE ROW AT PLANTING TIME OR AS CROP EMERGES. COVER THE BANK WITH 1-2 INCHES OF SOIL.

DO NOT APPLY DIRECTLY TO THE SEED FURROW AS REDUCTION OF SORGHUM SEED GERMINATION MAY OCCUR.

SOYBEANS: CUTWORMS: BROADCAST 14-28 LBS. PER ACRE JUST PRIOR TO PLANTING. WORK INTO SOIL 2-3 INCHES TO CONTROL SURFACE CUTWORMS; 3-6 INCHES TO CONTROL SUBTERRANEAN CUTWORMS.

LESSER CORNSTALK BORERS: APPLY 7-14 LBS. PER ACRE IN A 6-INCH BAND OVER THE ROW JUST PRIOR TO PLANTING OR WHEN SOYBEANS ARE 1-2 INCHES HIGH. COVER THE BAND WITH 1-2 INCHES OF SOIL.

SUGAR BEETS: SUGAR BEET ROOT MAGGOTS: APPLY 7-14 LBS. PER ACRE IN A 3-7 INCH BAND ABOVE THE SEED, OR APPLY AS A FURROW TREATMENT AFTER THE SEED IS COVERED. MAKE APPLICATION AHEAD OF THE PRESS WHEEL. THIS TREATMENT WILL REDUCE DAMAGE AND LOSS OF GERMINATING SEEDLINGS DUE TO SUGAR-BEET ROOT MAGGOT.

SUGARCANE: SUGARCANE BORERS: BROADCAST 10 1/2-17 1/2 LBS. PER ACRE AFTER JOINTS HAVE BEGUN TO FORM AND WHEN AT LEAST 5% OF PLANTS SHOW BORERS FEEDING IN OR UNDER THE LEAF SHEATH BUT WHICH HAVE NOT BORED INTO THE STALKS. ADDITIONAL APPLICATIONS MAY BE NECESSARY TO ASSURE CONTINUING CONTROL.

NOTE: DO NOT MAKE MORE THAN FOUR APPLICATIONS PER GROWING SEASON. ALLOW 28 DAYS BETWEEN LAST APPLICATION AND HARVEST. AT THAT TIME BAGASSE AND FORAGE MAY BE FED TO LIVESTOCK.

WIREWORMS: APPLY 42 LBS. PER ACRE IN A 12-INCH BAND OVER THE OPEN FURROW, DIRECTLY TO SEED PIECES.

FORAGE CROPS (ALFALFA, CLOVER, CONFEAS, AND LESPEDEZA): CUTWORMS: BROADCAST 14-28 LBS. PER ACRE JUST PRIOR TO PLANTING. WORK INTO SOIL 2-3 INCHES TO CONTROL SURFACE CUTWORMS, 3-6 INCHES TO CONTROL SUBTERRANEAN CUTWORMS.

VEGETABLE CROP INSECTS

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, AND CAULIFLOWER: ROOT MAGGOTS: BROADCAST 14-21 LBS. PER ACRE BEFORE PLANTING OR TRANSPLANTING. WORK INTO SOIL 3-4 INCHES.

ONIONS: ONION MAGGOTS: APPLY 14-28 LBS. PER ACRE AS A BROADCAST SOIL TREATMENT JUST PRIOR TO PLANTING. THOROUGHLY MIX THE CHEMICAL INTO THE TOP 3-4 INCHES OF SOIL BY DISKING, HARRAWING OR OTHER SUITABLE MEANS.

RADISHES: ROOT MAGGOTS: APPLY 9 1/2-14 LBS. PER ACRE (2 1/2-4 1/2 OZ. PER 1,000 LINEAR FEET OF ROW) IN SEED FURROW AT PLANTING TIME.

TOMATOES: YINEGAP FLIES (DROSOPHILA SPP.): BROADCAST 7 LBS. PER ACRE WHEN FLIES FIRST APPEAR. REPEAT IN 7-10 DAY INTERVALS OR AS REQUIRED.

Diazinon 14G Granular Insecticide

FOR CONTROL OF SOIL INSECTS SUCH AS ROOTWORMS, WIREWORMS, CUTWORMS, ORNAMENTAL, TURF AND OTHER CROP INSECT PESTS

ACTIVE INGREDIENT:

Table with 2 columns: Ingredient name and percentage. Includes 0,0-DIETHYL O-(2-ISOPROPYL-6-METHYL-4-PYRIMIDINYL) PHOSPHOROTHIOATE (14.3%) and INERT INGREDIENTS (85.7%).

(P) DIAZINON IS A REG. TRADEMARK OF CIBA-GEIGY, LTD. PATENT NO. 2,750,114

CAUTION: Keep Out of Reach of Children

HARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FOOD AND WATER. DO NOT STORE NEAR FOOD OR FEED PRODUCTS. AVOID INHALATION OF DUST. AVOID CONTACT WITH SKIN, EYES, AND CLOTHING. WASH THOROUGHLY AFTER HANDLING. MAY BE ABSORBED THROUGH SKIN.

FISH AND WILDLIFE CAUTION

THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE. KEEP OUT OF LAKES, STREAMS, OR PONDS. BIRDS FEEDING ON TREATED AREAS MAY BE KILLED. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DO NOT REUSE CONTAINER. DESTROY IT BY BURYING OR BURNING. AVOID SMOKE OR FUMES.

DIRECTIONS FOR USE AND CONDITIONS OF SALES, WARRANTY AND LIMITATION OF DAMAGE PROVISION.

IMPORTANT: READ THESE ENTIRE DIRECTIONS, CONDITIONS OF SALE, WARRANTY AND LIMITATION OF DAMAGE PROVISION, BEFORE USING B & G DIAZINON 14 G.

CONDITIONS OF SALE: THESE DIRECTIONS FOR USE REFLECT THE OPINION OF EXPERTS BASED ON TESTS OF EFFECTIVENESS UNDER NORMAL CONDITIONS OF USE. HOWEVER, BECAUSE OF THE WIDE RANGE OF CONDITIONS UNDER WHICH THIS PRODUCT MAY BE USED, EVEN THOUGH LABEL DIRECTIONS ARE FOLLOWED, IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS ASSOCIATED WITH ITS USE BECAUSE OF ABNORMAL CONDITIONS BEYOND THE CONTROL OF THE SELLER. B & G COMPANY AND THE SELLER OFFER, AND THE BUYER USES, THIS PRODUCT SUBJECT TO THE UNDERSTANDING THAT THE BUYER ASSUMES ALL SUCH RISKS.

WARRANTY AND LIMITATION OF DAMAGES: B & G COMPANY WARRANTS THAT THIS MATERIAL CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL AND IS REASONABLY FIT FOR THE PURPOSES REFERRED TO IN THE DIRECTIONS, CONDITIONS OF SALE; SUBJECT TO RISKS REFERRED TO THEREIN. B & G COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY, OR ANY OTHER EXPRESS OR IMPLIED WARRANTY, AND NO AGENT OF B & G COMPANY IS AUTHORIZED TO DO SO EXCEPT IN WRITING, WITH A SPECIFIC REFERENCE TO THIS WARRANTY. IN NO CASE SHALL B & G COMPANY OR THE SELLER BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

FEA Reg. No. 8617-62

NET WEIGHT \_\_\_\_\_ POUNDS

EPA Reg. No. 1012-TX 1.2 PK-1



Handwritten notes and signatures in the top right corner.

GARDEN SYMPHYLANT CENTIPEDS

Table listing vegetable crops: CABBAGE, CARROTS, LETTUCE, PEAS, POLE BEANS, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, COLLARDS, ENDIVE (ESCAROLE), KALE, RADISHES, RED BEETS, SNAP BEANS, TOMATOES, TURNIPS.

PROCEED WITH CARE IN GARDENS. WORK INTO SOIL 4-6 INCHES OR PLANT TREATMENT WILL AUTOMATICALLY CONTROL CUTWORMS, TERRESTRIAN MITE, CRICKETS, ROOT MAGGOTS, AND WIREWORMS.

MOLE CRICKETS:

Table listing vegetable crops: BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, COLLARDS, ENDIVE (ESCAROLE), KALE, LETTUCE, PEPPERS, RADISHES, TOMATOES, TURNIPS.

APPLY 7 LBS. PER ACRE WHEN INSECTS ARE PRESENT. MAKE APPLICATION TO PLANT BEDS OR FIELDS 1-2 DAYS PRIOR TO SOWING OF SEED OR TRANSPLANTING AND WORK INTO SOIL 1-2 INCHES.

LESSER CORNSTALK BORERS:

BEANS (LIMA, PINTO, SNAP): APPLY 7-14 LBS. PER ACRE IN A 6 INCH BAND CENTERED OVER THE ROW. APPLY WHEN PLANTS ARE 1/2-1 INCH TALL.

WIREWORMS:

Table listing vegetable crops: BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CANTALOUPE, CARROTS, CAULIFLOWER, CELERY, COLLARDS, CUCUMBERS, ENDIVE (ESCAROLE), KALE, LIMA BEANS, MUSKMELONS, ONIONS, PARSLEY, PEAS, PEPPERS, POLE BEANS, SNAP BEANS, SPINACH, SUMMER SQUASH, SWEET POTATOES, SWISS CHARD, TOMATOES, WATERMELONS, WINTER SQUASH.

BROADCAST 21-28 LBS. PER ACRE JUST PRIOR TO PLANTING. WORK INTO SOIL 4-6 INCHES.

POTATOES: BROADCAST 21-42 LBS. PER ACRE BEFORE PLANTING. WORK INTO SOIL 4-6 INCHES OR APPLY 7-21 LBS. PER ACRE AT PLANTING, PLACING DIAZINON IN BANDS 3-6 INCHES DEEP ALONG BOTH SIDES OF ROW.

SWEET POTATOES (SOUTHEASTERN U.S.): BROADCAST 21 LBS. PER ACRE ON THE TOPS OF VINES WHEN ROOTS BEGIN TO ENLARGE.

SMALL FRUIT INSECTS

STRAWBERRIES: MOLE CRICKETS: APPLY 7 LBS. PER ACRE WHEN INSECTS ARE PRESENT. MAKE APPLICATION TO PLANT BEDS OR FIELDS 1-2 DAYS PRIOR TO TRANSPLANTING AND WORK INTO SOIL 1-2 INCHES.

LAWN INSECTS

LAWN CHINCH BUGS: APPLY 1/2-1 LB. PER 1,000 SQ. FT. (2 1/2-5 LBS. PER 5,000 SQ. FT.) AND BRUSH LAWN TO WORK GRANULES DOWN TO THE SOIL SURFACE. WATER LAWN THOROUGHLY AFTER APPLICATION. REPEAT APPLICATION IN 7-10 DAYS IF NECESSARY. FOR LONGER RESIDUAL CONTROL AND ON DENSE GRASS TYPE SUCH AS ST. AUGUSTINE GRASS, USE 1 1/2-2 LBS. PER 1,000 SQ. FT. (3-7 1/2 LBS. PER 5,000 SQ. FT.).

TEXAS ONLY

Handwritten notes and signatures in the bottom right corner.