

**DIRECTIONS FOR USE, CONDITIONS OF SALES, WARRANTY  
and LIMITATION OF DAMAGE PROVISION**

**IMPORTANT:** Read these entire Directions, Conditions of Sale, Warranty, and Limitation of Damage Provision before using BGS 5 Dust Insecticide.

**CONDITIONS OF SALE:** These directions for use reflect the opinion of experts based on tests of effectiveness under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though the label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. B&G Company and the seller offer, and the buyer uses, this product subject to the understanding that the buyer assumes all such risks.

**WARRANTY AND LIMITATION OF DAMAGES:** B&G Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purpose referred to in the directions, conditions of sale; subject to risks referred to therein. B&G Company makes no other express or implied warranty of fitness or of merchantability or any other express or implied warranty, and no agent of B&G Company is authorized to do so except in writing, with a specific reference to this warranty. In no case shall B&G Company or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

**CAUTIONS**

**IMPORTANT PREHARVEST AND GRAZING USE  
INFORMATION & LIMITATIONS**

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Sevin residues when Sevin is used in accordance with label directions. Preharvest and grazing use limitations are listed at the end of the directions for each crop use. Where no preharvest or grazing limitations are stated, application can be made to the day of harvest, and crops can be grazed without time delay if directions for use on the crop are followed. If Sevin insecticide is used in accordance with label directions, forage and feed crops - including bean vines, carrot tops, cowpea hay, peanut hay, pea vines, rice straw, soybean hay and sugar beet tops - may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat.

**GENERAL CAUTIONS**

1. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.
2. DOES NOT control Spider Mites but is compatible with common miticides.

**DIRECTIONS FOR USE**

**IMPORTANT** The directions on this label are based on tests and field experience relating to (A) effectiveness, (B) possible injury to plants and animals, and (C) residues in food, feed, meat and milk. Follow directions for most effective and safest use.

**VEGETABLE INSECT CONTROL**

Suggested dosages refer to pounds of BGS 101 DUST per acre. Where a dosage range is indicated, use lower rate on young plants, higher rate on mature plants. Use sufficient dust to obtain full coverage. Apply when insects or their damage appear. Repeat at 7 to 10 day intervals or as necessary, unless shorter interval is specified below.

**BEANS** Green Beans, Lima Beans, Snap Beans, Cowpeas & Blackeyed Peas Use 5 lbs. for Mexican Bean Beetle; 10 lbs. for Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Velvet-Bean Caterpillar and Western Bean Cutworm.  
Use 10 to 15 lbs. for Armyworms, Corn Earworm, Stink Bug and Tarnished Plant Bug.

**IN CALIFORNIA** Use 20 lbs. for Corn Earworm, Lima Bean Pod Borer, Lygus and Stink Bugs.

**CORN** Use 10 to 20 lbs. for Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworms, Flea Beetles, Japanese Beetle, Leaf Beetles and Leaf Hoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Application of Sevin to the tassel region of corn during the pollen shed period will seriously reduce bee populations.

**CUCUMBERS, MELONS, PUMPKIN AND SQUASH** Use 5 to 10 lbs. for Pickworm and Melonworm; 10 lbs. for Cucumber Beetles, Flea Beetles, Leafhoppers and Squash Bugs. Avoid excessive applications. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Sevin injures watermelons in Florida.

**GARDEN PEAS** Use 10 lbs. for Leafhoppers, and Colorado Potato Beetle. For Colorado Potato Beetle control, direct application to weeds in fields or garden peas. This application is intended to help prevent insect contamination in processed peas.  
Use 10 to 15 lbs. for Armyworms and Grasshoppers.

**POTATO, TOMATO, EGGPLANT AND PEPPER** Use 5 to 10 lbs. for Colorado Potato Beetle, Flea Beetle and Leafhoppers. Use 10 to 20 lbs. for European Corn Borer, Fall Armyworm, Lacebugs, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug, and Stinkbugs.

**OKRA:** Use 10 to 20 lbs. for Corn Earworms and Stinkbugs. Repeat at 5 to 7 day intervals, or as necessary.

**SUBSEQUENT PLANTING OF FOOD AND FEED CROPS IN TREATED SOILS MUST BE MADE WITH CROPS ON THIS OR OTHER CARBARYL LABEL(S).**



# BGS 5 DUST INSECTICIDE

CONTAINS SEVIN® CARBARYL INSECTICIDE  
FOR CONTROL OF INSECT PESTS

ACTIVE INGREDIENT	Carbaryl (1-Naphthyl Methylcarbamate) ..	50	by wt.
INERT INGREDIENTS	.....	950	by wt.
	TOTAL	1000	

\*Sevin is the Registered Trademark of Union Carbide Corp. for Carbaryl Insecticide.

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS**  
CAUTION. May be harmful if swallowed. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Do not use this product in commercial food areas of food handling establishments, restaurants, or other places where food is prepared or processed.

Avoid contamination of food, feed, water supplies, streams and ponds.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

**READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.**

NET WEIGHT 50 POUNDS

MANUFACTURED FOR



DALLAS, TEXAS 75220

EPA REG. NO. 0612-75  
EPA EST. NO. 0612-TX-1

DIRECTIONS FOR USE (continued)

**ORNAMENTAL INSECT CONTROL**

Sufficient dust should be used on each plant so that the upper and lower leaf surfaces are thoroughly covered. Sevin 10 DUST can be used to control insects on ornamental plants. Specific plants and insects are listed below. Repeat 7 to 10 day intervals or as necessary unless shorter interval is specified below. Recommendations below are for treatment of individual plants.

**ROSES** Apply dust to upper and lower leaf surface for control of Rose Aphid, Rose Slug, Thrips and Plant Bug.

**ARBOR VITAE** Just to upper and lower leaf surface for control of Bag and Sawflies.

**ELM** Apply dust to upper and lower leaf surface for control of Elm Leaf Aphid, Elm Leaf Beetle, Leaf Roller and Perio Clebs.

**MIMOSA** Apply dust to upper and lower leaf surface for control of Mimosa Webworm.

**OAK** Apply dust to upper and lower leaf surface for control of Oak Leaf Miner, Sciss Insects and Leaf Roller.

**BIRCH** Apply dust to upper and lower leaf surface for control of Birch Leaf Miner.

**PYRACANTHA** Apply dust to upper and lower leaf surface for control of Lacebugs.

**NOTE** Sevin does not control spider mites. Where infestations are encountered, use a recommended miticide.

Sevin injures Boston Ivy.

**HOUSEHOLD INSECT CONTROL**

**GENERAL DIRECTIONS FOR USE:** Do not apply where children are likely to come in frequent contact with treated surfaces. Do not contaminate food, food containers or containers. Do not use in edible products areas of food processing plants, restaurants or other areas where food is commercially prepared or processed. Do not use in serving areas while food is being prepared.

Apply dust only in a sufficient quantity to produce a fine coating. If applied too heavily, excess dust will tend to repel insect pests.

**FOR CONTROL OF ANTS:** Apply BGS 5 DUST to ant locations such as door sills, around window frames and at pipe into structure entrances, etc. Dust ant trails and holes. Repeat as necessary.

**FOR CONTROL OF FLEAS AND BROWN DOG TICKS ON PREMISES:** Thoroughly dust all infested areas with BGS 5 DUST including cracks and crevices in woodwork, floor and floor coverings, and animal sleeping quarters. Repeat as necessary.