

**IMPORTANT.** Read these entire Directions, Conditions of Sale, Warranty, and Limitation of Damage Provision before using BGS 10 Dust Insecticide.

**CONDITIONS OF SALE** These directions for use reflect the opinion of experts based on tests of effectiveness under normal conditions of use. However, because of the wide range of conditions under which this product may be used, even though label directions are followed, it is impossible to eliminate all risks associated with its use because of abnormal conditions beyond the control of the seller. B&G Company and the seller offer, and the buyer uses, this product subject to the understanding that the buyer assumes all such risks. **WARRANTY AND LIMITATION OF DAMAGES** B&G Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions, conditions of sale, subject to risks referred to therein. B&G Company makes no other express or implied warranty of fitness or of merchantability, or any other express or implied warranty, and no agent of B&G Company is authorized to do so except in writing, with a specific reference to this warranty. In no case shall B&G Company or the seller be liable for consequential, special or indirect damages resulting from the use or handling of this product.

#### CAUTIONS

#### IMPORTANT PREHARVEST AND GRAZING USE INFORMATION & LIMITATIONS

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Sevin residues when Sevin is used in accordance with label directions. Preharvest and grazing use limitations are listed at the end of the directions for each crop use. Where no preharvest or grazing limitations are stated, application can be made to the day of harvest, and crops can be grazed without time delay if directions for use on the crop are followed. If Sevin Insecticide is used in accordance with label directions, forage and feed crops - including bean vines, carrot tops, cowpea hay, peanut hay, pea vines, rice straw, soybean hay and sugar beet tops - may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat.

#### GENERAL CAUTIONS

- To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days.
- DOES NOT control Spider Mites but is compatible with common miticides.

#### DIRECTIONS FOR USE

**IMPORTANT.** The directions on this label are based on tests and field experience relating to (A) effectiveness, (B) possible injury to plants or animals, and (C) residues in food, feed, meat and milk. Follow directions for most effective and safest use.

Suggested dosages refer to pounds of BGS 10 DUST per acre. Where a dosage range is indicated, use lower rate on young plants, higher rate on mature plants. Use sufficient dust to obtain full coverage. Apply when insects or their damage appear. Repeat at 7 to 10 day intervals or as necessary, unless shorter interval is specified below.

**BEANS** Green Beans, Lima Beans, Snap Beans, Cowpeas & Black-eyed Peas Use 5 lbs. for Mexican Bean Beetle; 10 lbs. for Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Velvet-Bean Caterpillar and Western Bean Cutworm. Use 10 to 15 lbs. for Armyworms, Corn Earworm, Stink Bug and Tarnished Plant Bug.

**IN CALIFORNIA:** Use 20 lbs. for Corn Earworm, Lima Bean Pod Borer, Lygus and Stink Bug.

**CORN.** Use 10 to 20 lbs. for Corn Earworm, Corn Rootworm Adults, European Corn Borer, Fall Armyworm, Flea Beetles, Japanese Beetle, Sap Beetles and Leaf Hoppers. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. Application of Sevin to the tassal region of corn during the pollen shed period will seriously reduce bee populations.

**CUCUMBERS, MELONS, PUMPKIN AND SQUASH** Use 5 to 10 lbs. for Pickleworm and Melonworm; 10 lbs. for Cucumber Beetle, Flea Beetles, Leafhoppers and Squash Bug. Avoid excessive applications. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Sevin injures watermelons in Florida.

**GARDEN PEAS:** Use 10 lbs. for Leafhoppers, and Colorado Potato Beetle. For Colorado Potato Beetle control, direct application to weeds in fields or garden peas. This application is intended to help prevent insect contamination in processed peas.

Use 10 to 15 lbs. for Armyworms and Grasshoppers.

**POTATO, TOMATO, EGGPLANT AND PEPPER** Use 5 to 10 lbs. for Colorado Potato Beetle, Flea Beetle and Leafhoppers. Use 10 to 20 lbs. for European Corn Borer, Fall Armyworm, Lacebugs, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bug, and Stinkbug.

**OKRA** Use 10 to 20 lbs. for Corn Earworm and Stinkbug. Repeat at 5 to 7 day intervals, or as necessary.

SUBSEQUENT PLANTING OF FOOD AND FEED CROPS IN TREATED SOILS MUST BE MADE WITH CROPS ON THIS OR OTHER CARBARYL LABEL(S).

# B&G

# BGS

# 10 DUST

## INSECTICIDE

CONTAINS SEVIN® CARBARYL INSECTICIDE FOR CONTROL OF INSECT PESTS

#### ACTIVE INGREDIENT

Carbaryl (1-Naphthyl Methylcarbamate) ... 10% by wt.  
INERT INGREDIENTS: ..... 90% by wt.  
TOTAL 100%

\*Sevin is the Registered Trademark of Union Carbide Corp. for Carbaryl Insecticide.

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS

CAUTION May be harmful if swallowed. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Do not use this product in commercial food areas of food handling establishments, restaurants, or other places where food is prepared or processed.

Avoid contamination of food, feed, water supplies, streams and ponds.

NOTE TO PHYSICIAN Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

MANUFACTURED FOR

EPA REG. NO. 8612-00  
EPA EST. NO. 8612 TX-1

**B&G**  
**COMPANY**

NET WEIGHT 50 POUNDS

DALLAS, TEXAS 75220

Sufficient dust should be used on each plant upper and lower leaf surfaces are thoroughly Sevin 10 DUST can be used to control insect plants. Specific plants and insects are listed 7 to 10 day intervals or as necessary unless is specified below. Recommendations below of individual plants.

**ROSES.** Apply dust to upper and lower leaf control of Rose Aphids, Rose Slug, Thrips

**ARBOR VITAE** Apply dust to upper and lower leaf control of Bagworm and Sawflies.

**ELM** Apply dust to upper and lower leaf of Elm Leaf Aphid, Elm Leaf Beetle, Leaf Roll Ckade.

**MIMOSA:** Apply dust to upper and lower leaf control of Mimosid Weevils.

**OAK.** Apply dust to upper and lower leaf of Oak Leaf Miner, Scale Insects and Leaf Rollers.

**BIRCH** Apply dust to upper and lower leaf control of Birch Leaf Miner.

**PYRACANTHA.** Apply dust to upper and lower leaf control of Lacebugs.

NOTE: Sevin does not control spider mites. are encountered, use a recommended miticide.

Sevin Injures Boston Ivy.

#### HOUSEHOLD INSECT CONTROL

**GENERAL DIRECTIONS FOR USE.** Do not let children are likely to come in frequent contact surfaces. Do not contaminate food, food containers, utensils. Do not use in edible products areas of food restaurants or other areas where food is consumed or processed. Do not use in serving areas.

Apply dust only in a sufficient quantity to provide light coating. If applied too heavily, excess will repel insect pests.

**FOR CONTROL OF ANTS** Apply BGS 10 DUST to locations such as door sills, around window pipe into structure entrances, etc. Dust on Repeat as necessary.

**FOR CONTROL OF FLEAS AND BROWN DOG PREMISES** Thoroughly dust all infested areas 10 DUST including cracks and crevices in walls and floor coverings, and animal sleeping quarters as necessary.