

171851
17/03
300

JUL 1 1986

Mr. K.H. Miller
B&G Company
10539 Maybank
P.O. Box 20372
Dallas, TX 75220

Dear Mr. Miller:

Subject: General Revision of Label
BGS 10 Dust
EPA Registration No. 8612-80
Your Application Dated March 31, 1986

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

In your Environmental Hazards section the paragraph "This product is toxic to fish and wildlife. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on the label" can be deleted.

Sincerely yours,

Lawrence J. Schnaubelt ^{LJS}
Acting Product Manager (12)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

Enclosure

92964:Schnaubelt:E-9:KENCO:6/6/86:6/12/86:eg:vo

CONCURRENCES

SYMBOL	SURNAME	DATE						

PRECAUTIONARY STATEMENTS - Hazards to Humans
CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Do not use this product in commercial food areas of food handling establishments, restaurants, or other places where food is prepared or processed. Avoid contamination of food, feed, water supplies, streams and ponds. Read entire label. Use in strict accordance with label directions and cautions.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

IMPORTANT PREHARVEST AND GRAZING USE INFORMATION & LIMITATIONS: Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Carbaryl insecticide residues when Sevin® Brand Carbaryl Insecticide is used in accordance with label directions. Preharvest and grazing use limitations are listed at the end of directions for each crop use. Where no preharvest or grazing limitations are stated, application can be made to the day of harvest, and crops can be grazed without time delay if directions for use on the crop are followed. If this product is used in accordance with label directions, forage and feed crops (including bean vines, carrot tops, cowpea hay, peanut hay, pea vines, rice straw, soybean hay and sugar beet tops) may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat.

GENERAL CAUTIONS: (1) To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next two days. (2) DOES NOT control Spider Mites but is compatible with common miticides.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and wildlife. **KEEP OUT OF LAKES, STREAMS OR PONDS.** Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

IMPORTANT: The directions on this label are based on tests and field experience relating to (a) effectiveness, (b) possible injury to plants and animals, and (c) residues in food, feed, meat and milk. Follow directions for most effective and safest use.

VEGETABLE INSECT CONTROL

Suggested dosages refer to pounds of BGS 10 DUST per acre. Where a dosage range is indicated, use lower rate on young plants, higher rate on mature plants. Use sufficient dust to obtain full coverage. Apply when insects or their damage appear. Repeat at 7 to 10 day intervals, or as necessary, unless shorter interval is specified below.

BEANS. Green Beans, Lima Beans, Snap Beans, Cowpeas & Blackeyed Peas. Use 5 lbs. for Mexican Bean Beetle, 10 lbs. for Bean Leaf Beetle, Cucumber Beetles, Flea Beetles, Japanese Beetle, Leafhoppers, Velvet bean Caterpillar, and Western Bean Cutworm. Use 10 to 15 lbs. for Armyworms, Corn Earworm, Stink Bugs and Tarnished Plant Bug. **IN CALIFORNIA:** Use 20 lbs. for Corn Earworm, Lima Bean Pod Borer, Lygus and Stink Bug.

CUCUMBERS, MELONS, PUMPKIN AND SQUASH: Use 5 to 10 lbs. for Pickleworm and Mealy Bug, 10 lbs. for Cucumber Beetles, Flea Beetles, Leafhoppers and Squash Bugs. Avoid excessive applications. To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days. Carbaryl insecticide injures watermelons in Florida.

POTATO, TOMATO, EGGPLANT AND PEPPER: Use 5 to 10 lbs. for Colorado Potato Beetle, Flea Beetle, and Leafhoppers. Use 10 to 20 lbs. for European Corn Borer, Fall Armyworm, Lacebugs, Tomato Fruitworm, Tomato Hornworm, Tarnished Plant Bugs and Stinkbugs.

BGS 10

CONTAINS SEVIN FOR CONT

ACTIVE INGREDIENT
Carbaryl (1-naphthyl)
INERT INGREDIENTS

Sevin is the Registered T

KEEP OUT OF CA

STATEMENT OF

IF SWALLOWED: Call a physician if possible, vomiting should be induced. Drink one or two glasses of water. If necessary, drink anything by mouth to dilute the poison.
IF INHALED: Remove victim to fresh air. If indicated, remove clothing.
IF ON SKIN: Remove clothing. Wash with soap and water.
IF IN EYES: Flush eyes immediately.

EPA REG. NO. 8612 90
EPA EST. NO. 34874 TX 1

LABEL # 0718751 06

DIRECTIONS FOR USE (continued)

OKRA: Use 10 to 20 lbs. for Corn Earworms and Stinkbugs. Repeat at 5 to 7 day intervals, or as necessary.

SUBSEQUENT PLANTING OF FOOD AND FEED CROPS IN TREATED SOILS MUST BE MADE WITH CROPS ON THIS OR OTHER CARBARYL LABEL(S)

ORNAMENTAL INSECT CONTROL

Sufficient dust should be used on each plant so that the upper and lower leaf surfaces are thoroughly covered. This product can be used to control insects on ornamental plants. Specific plants and insects are listed below. Repeat at 7 to 10 day intervals or as necessary unless shorter intervals are specified below. Recommendations below are for treatment of individual plants.

ROSES: Apply dust to upper and lower leaf surface for control of Rose Aphid, Rose Slug, Thrips and Plant Bugs.

PYRACANTHA: Apply dust to upper and lower leaf surface for control of Lacebugs.

NOTE: Carbaryl insecticide does not control spider mites. Where infestations are encountered, use a recommended miticide. Carbaryl insecticide injures BOSTON IVY.

HOUSEHOLD INSECT CONTROL

GENERAL DIRECTIONS FOR USE: Do not apply where children are likely to come in frequent contact with treated surfaces. Do not contaminate food, food containers or cooking utensils. Do not use in edible products areas of food processing plants, restaurants, or other areas where food is commercially prepared or processed. Do not use in serving areas while food is exposed. Apply dust only in sufficient quantity to produce a fine light coating. If applied too heavily, excess dust will tend to repel insect pests.

FOR CONTROL OF ANTS: Apply BGS 10 DUST to ant entrance locations such as door sills, around window frames and sash, pipe into structure entrances, etc. Dust ant trails and hills. Repeat as necessary.

FOR CONTROL OF FLEAS AND BROWN DOG TICKS ON PREMISES: Thoroughly dust all infested areas with BGS 10 DUST including cracks and crevices in woodwork, floors and floor coverings, and animal sleeping quarters. Repeat as necessary.

PET INSECT CONTROL

ON DOGS AND CATS: For Ticks and Fleas, dust liberally over the entire animal, avoiding the eyes, and rub thoroughly into the skin. Comb out dead ticks in a few hours after treatment. Repeat at weekly intervals if needed. Avoid treatment of kittens less than four weeks of age. DO NOT USE ON PREGNANT DOGS.

IN OR AROUND DOG AND CAT QUARTERS: For Ticks and Fleas, dust liberally in and around sleeping quarters, kennels and dog houses by means of a shaker can, hand or power duster. Force dust into cracks and crevices. Repeat as necessary.

STORAGE AND DISPOSAL

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL

Plastic Container: Do not reuse container. Rinse thoroughly before discarding in trash.
Paper Bag: Do not reuse bag. Discard bag in trash.

PESTICIDE STORAGE: Store in a cool, dry place, inaccessible to children and pets.

IMPORTANT

Seller's guarantee shall be limited to the terms of this label, and subject thereto, the buyer assumes any risks to persons or property rising out of the use or handling and accepts the product on these conditions.

