PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

MARMFUL IF SWALLOWED. AVOID CONTAMINATION OF FOOD OR FEED. DO NOT STORE NEAR FOOD OR FEED PRODUCTS. AVOID INHALATION OF DUST. AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WASH THOROUGHLY AFTER HANDLING. MAY BE ABSORBED THROUGH SKIN. KEEP CHILDREN AND PETS OFF TREATED AREAS UNTIL THIS HATERIAL IS WASHED INTO THE SOIL AND GRASS IS DRY.

ENVIRONMENTAL HAZARDS: THIS PRODUCT IS TOXIC TO FISH, BIRDS AND OTHER WILDLIFE. KEEP OUT OF LAKES, STREAMS OR PONDS. BIRDS FEEDING ON TREATED AREAS MAY BE KILLED. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

DO NOT ALLOW CHILDREN ON TREATED AREAS UNTIL MATERIAL IS WASHED INTO THE SOIL AND DRY.

DISPOSAL

DO NOT USE EMPTY CONTAINER

WRAP CONTAINER AND PUT IN TRASH CO. - SCTION

OVERALL LAWN APPLICATION: DIAZINON 5% GRANULES CONTROLS THE FOLLOWING LAWN INSECTS ON BOTH GRASSES AND DICHONDRA LAWNS; CHINCH BUGS, EARWIGS, ANTS, LAWN MOTHS (SOD WEBWORMS), BERMUDA MITES, CRICKETS, ARMYWORMS, BROWN DOG TICKS, CHIGGERS, CLOVER MITES, CUTWORMS, FLEAS, LEAFHOPPERS, SOWBUGS, MILLEPEDES, SPRINGTAILS (COLLEMBOLA).

APPLY DIAZINON 5% GRANULES TO YOUR LAWN WITH A FERTILIZER SPREADER. APPLICATION SHOULD BE MADE WHEN GRASS IS DRY. WATER LAWN LIGHTLY AFTER APPLICATION. REPEAT AS NECESSARY. SEE FOLLOWING PANEL FOR SPREADER SETTINGS.

FOR APPLICATION WITH OTHER TYPES OF EQUIPMENT, OR BY MAND, APPLY AT THE RATE OF 2 LBS. PER 1000 SQ.FT. OR 3.2 OZ. PER 100 SQ.FT. FOR HEAVY LAWNS SUCH AS ST. AUGUSTINE LAWNS, APPLICATION SHOULD BE INCREASED TO 3 LBS. PER 1000 SQ.FT. OR 4.8 OZ. PER 100 JQ. FT.

OR 4.8 OZ, PER 100 JQ. FT.

BILLBUGS AND WHITE GRUBS OF JAPANESE BEETLE, EUROPEAN CHAFER

AND SOUTHERN CHAFER. APPLY 2.5 LBS, PER 1000 SQ.FT. OF LAWN.

MATER GRASS THOROUGHLY AFTER APPLICATION. FOR GRUB CONTROL



FOR CONTROL OF CHINCH BUGS, SOD WEBWORMS, LAWN MOTHS, EARWIGS, MILLEPEDES, BROWN DOG TICKS, FLEAS, GRUBS AND

OTHER INSECTS LISTED HEREIN

DIAZINON IS A REGISTERED TRADEMARK OF CIBA GEIGY.

READY TO USE. CONTAINS 5% DIAZINON INSECTICIDE.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED-DRINK ONE OR TWO GLASSES OF WATER AND INDUCE

VOMITING BY TOUCHING THE BACK OF THE THROAT WITH

FINGER. REPEAT UNTIL VOMIT FLUID IS CLEAR. CALL

A PHYSICIAN IMMEDIATELY. DO NOT INDUCE VOMITING

OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

IF INHALED
REMOVE VICTIM TO FRESH AIR. APPLY RESPIRATION IF

INDICATED.

IF ON SKIN
REMOVE CONTAMINATED CLOTHING AND WASH AFFECTED AREAS
WITH SOAP AND WATER.

1F IN EYES- FLUSH EYES FOR AT LEAST 15 MINUTES WITH WATER. CALL PHYSICIAN IF IRRITATION PERSISTS.

Keep Out of Reach of Children CAUTION

SEE SIDE PANELS FOR ADDITIONAL CAUTIONS.

EPA REG, No. 8612-100 EPA Est. No. 8612-TX-1,2;0K-1 MANUFACTURED BY

BOMPANY

NET CONTENTS 50 POUNDS LABEL # 012861

16

DALLAS, TEXAS 75220



FOR CONTROL OF CHINCH BUGS, SOD WEBVORMS, LAWN MOTHS, EARWIGS, MILLEPEDES, BROWN DOG TICKS, FLEAS, GRUDS AND

OTHER INSECTS LISTED HEREIN

ACTIVE INGREDIENTS:
0,0-DIETHYL 0-(2-150PROPYL-6-METHYL-4-PYRIMIDINYL)
PHOSPHOROTHIOATE..... TOTAL......95.00% INERT INGREDIENTS: R

DIAZINON IS A REGISTERED TRADEMARK OF CIBA GEIGY.

READY TO USE. CONTAINS 5% DIAZINON INSECTICIDE.

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED-DRINK ONE OR TWO GLASSES OF WATER AND INDUCE
VOMITING BY TOUCHING THE BACK OF THE THROAT WITH
FINGER. REPEAT UNTIL VOMIT FLUID IS CLEAR. CALL A PHYSICIAN IMMEDIATELY. DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. IF INHALED-REMOVE VICTIM TO FRESH AIR. APPLY RESPIRATION IF INDICATED. IF ON SKIN-REMOVE CONTAMINATED CLOTHING AND WASH AFFECTED AREAS WITH SOAP AND WATER.
FLUSH EYES FOR AT LEAST 15 MINUTES WITH WATER. CALL
PHYSICIAN IF IRRITATION PERSISTS. IF IN EYES-

Keep Out of Reach of Children CAUTION

SEE SIDE PANELS FOR ADDITIONAL CAUTIONS.

12-100 12-TX-1,2;0K-1



NET CONTENTS 50 Pounds LABEL # 012801

DALLAS, TEXAS 75220

DIRECTIONS FOR USE

CONTINUED

MAKE APPLICATION ANY TIME BETWEEN LATE JULY AND EARLY OCTO-BER. FOR BILLBUG CONTROL APPLY WHEN ACTIVITY 19 FIRST OB-SERVED OR WHEN CHEWED OR BROWN GRASS NED CATES DAMAGE FROM THESE INSECTS. REPEAT AS NECESSARY.

> SPREADER SETTING NO. (PER 1000 SQ. FT.)

	WER 1000	34, 11,7	•• • • •
SPREADER	INSECTS LISTED BELOW" .		WHITE GRUBS
MAKE & MODEL	REGULAR OR DICHONDRA LAWN 2_LBS.	ST. AUGUSTINE LAWN 3 LBS.	AND BILLBUGS
SCOTT	5-1/8	8-1/1	**£-574
JACKSON 18-26 (18")	5	5-573*	5-1/2
SEARS ROEBUCK CROSS COUNTRY		• • • •	
(19")	6-1/4	6-3/4	6-1/2
SALEM MICRO SET (22")	9-1/4	8-5/8	8-7/8
CYCLONE MODELS 10,15,20 MODEL B-1	4-3/4	5-2/4 5	5-1/4 5-1/2
LAWN BEAUTY	6	6-7/8	6-1/2
GREENFIELD MODEL 103	1 4	5	5-1/2
MODEL 105	7	7-1/2	8_

*CHINCH BUGS, EARWIGS, ANTS, LAWN MOTHS (SOD WEBWORMS) BERMUDA MITES, CRICKETS, ARMYWORMS, BROWN DOG TICKS, CHIGGERS,
CLOVER MITES, CUTWORMS, FLEAS, LEAFHOPPERS, SOWBUGS, MILLEPEDES, SPRINGTAILS (COLLEMBOLA)
NOTE: ABOVE SPREADER SETTINGS ARE APPROXIMATE. ACCURATELY
WEIGH A FEW POUNOS INTO YOUR SPREADER AND APPLY TO A MEASURED
AREA. THEN ADJUST THE SPREADER SETTING IF NECESSARY IN OR-

DER TO ASSURE CORRECT RATE OF APPLICATION.

SPOT TREATMENT: ANTS: APPLY 1-1/2 TEASPOONS OF GRANULES OVER AND AROUND EACH INDIVIDUAL ANT HILL.

DIGGER WASPS: APPLY 1-1/2 TEASPOONS OF GRANULES IN AND AROUND EACH INDIVIDUAL GROUND NEST OPENING. MAKE APPLICATION ABOUT DUSK WHEN THE WASPS ARE NOT ACTIVE.

BAND TREATMENT AROUND FOUNDATION: TO CONTROL ANTS, BROWN DOG TICKS, CLOVER MITES, CRICKETS, EARWIGS, FLEAS, MILLE-PEDES, SOWBUGS AND SPRINGTAILS; APPLY A FIVE-FOOT BAND OF GRANULES NEXT TO THE FOUNDATION OF THE HOUSE. DISTRIBUTE THE GRANULES UNIFORMLY USING A RATE OF 3.2 OZ. FOR EACH, FIVE FOOT BY TWENTY FOOT AREA TO BE TREATED.

IMPORTANT

SELLER'S GUARANTEE SHALL BE LIMITED TO THE TERMS OF THIS A
LABEL AND, SUBJECT THERETO, THE BUYER ASSUMES ANY RISKS
TO PERSONS OR PROPERTY RISING OUT OF THE USE OR HANDLING
AND ACCEPTS THE PRODUCT ON THESE CONDITIONS.