



Diazinon® 25E

LAWN & GARDEN INSECT CONTROL

F-12-101

ACTIVE INGREDIENTS:
 O,O-DIETHYL O-(2-ISOPROPYL-6-METHYL-5-PYRIDINYL) PHOSPHOROTHIOATE..... 27.01
 AROMATIC PETROLEUM DERIVATIVE SOLVENT..... 50.44
 INERT INGREDIENTS..... 20.54
 TOTAL 100.00

CONTAINS: DIAZINON INSECTICIDE SPRAY CONCENTRATE
 ®
 DIAZINON IS A REGISTERED TRADEMARK OF CIBA-GEIGY, INC.

KEEP OUT OF REACH OF CHILDREN
WARNING
 SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED-CALL A PHYSICIAN IMMEDIATELY. GASTRIC LAVAGE IS INDICATED IF MATERIAL HAS BEEN TAKEN INTERNALLY. DO NOT INDUCE VOMITING UNLESS OTHER TREATMENT IS NOT AVAILABLE. VOMITING MAY CAUSE ASPIRATION PNEUMONIA. IF IT IS NECESSARY TO INDUCE VOMITING, GIVE VICTIM ONE OR TWO GLASSES OF WATER AND INSERT FINGER IN BACK OF THROAT. REPEAT UNTIL VOMIT IS CLEAR. DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. REMOVE VICTIM TO FRESH AIR. APPLY RESPIRATION IF INDICATED.
 IF INHALED-
 IF ON SKIN-
 IF IN EYES-
 WASH CONTAMINATED SKIN WITH SOAP AND WATER.
 FLUSH EYES WITH PLENTY OF WATER FOR AT LEAST 15 MINUTES. GET MEDICAL ATTENTION IMMEDIATELY.

EPA REG. NO. 8612-101
 NET CONTENTS 1 GALLON
 EPA EST. NO. 8612-T-1.20K-1
 LABEL NO. 012802



10539 MAYBANK
 DALLAS, TEXAS 75220

36

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING
 MAY BE FATAL IF SWALLOWED OR ABSORBED THROUGH SKIN.
 AVOID BREATHING SPRAY MIST. AVOID CONTACT WITH EYES,
 SKIN AND CLOTHING. WASH THOROUGHLY IF ON SKIN AND
 AFTER HANDLING. AVOID CONTAMINATION OF FEED AND FOOD-
 STUFFS. DO NOT USE ON HOUSEHOLD PETS OR HUMANS. DO
 NOT PERMIT CHILDREN OR PETS TO GO ONTO SPRAYED GRASS
 UNTIL SPRAY HAS COMPLETELY DRIED. DO NOT USE IN THE
 HOME. DO NOT CONTAMINATE ORNAMENTAL FISH PONDS.
ENVIRONMENTAL HAZARDS
 THIS PRODUCT IS TOXIC TO FISH, SHRIMP, CRAB, RIDS AND
 OTHER WILDLIFE. BIRDS FEEDING ON TREATED AREA MAY BE
 KILLED. KEEP OUT OF ANY BODY OF WATER. DO NOT APPLY
 WHEN WEATHER CONDITIONS FAVOR DRIFT FROM TREATED AREAS.
 DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT OR
 DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED
 ON THIS LABEL.
 THIS PESTICIDE IS HIGHLY TOXIC TO BEES EXPOSED TO DIRECT
 TREATMENT OR TO RESIDUES REMAINING ON THE TREATED AREA.
 DO NOT APPLY WHEN BEES ARE ACTIVELY VISITING THE CROP,
 CROP COVER, OR NEEDS BLOOMING IN THE TREATMENT AREA.
 APPLICATIONS SHOULD BE TIMED TO PROVIDE THE MAXIMUM
 POSSIBLE INTERVAL BETWEEN TREATMENT AND THE NEXT PERIOD
 OF BEE ACTIVITY.
PHYSICAL OR CHEMICAL HAZARDS
 DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS FOR USE

IS A VIOLATION OF FEDERAL LAW TO USE
 IN A MANNER INCONSISTENT WITH ITS LABEL.

STORAGE & DISPOSAL

- 1. PROHIBITIONS: DO NOT CONTAMINATE WATER, FOOD, OR FEED BY STORAGE OR DISPOSAL. OPEN DUMPING IS PROHIBITED.
- 2. PESTICIDE DISPOSAL: PESTICIDE, SPRAY, MIXTURE, OR RINSATE THAT CANNOT BE USED OR CHEMICALLY PROCESSED SHOULD BE DISPOSED IN A LANDFILL APPROVED FOR PESTICIDES OR BURIED IN A SAFE PLACE AWAY FROM WATER SUPPLIES.
- 3. CONTAINER DISPOSAL: TRIPLE RINSE (OR EQUIVALENT) AND OFFER FOR RECYCLING, RECONDITIONING, OR DISPOSAL IN APPROVED LANDFILL OR BURY IN A SAFE PLACE.
- 4. GENERAL: CONSULT FEDERAL, STATE OR LOCAL DISPOSAL AUTHORITIES FOR APPROVED ALTERNATIVE PROCEDURES.

DIRECTIONS FOR USE

HUANSE PESTS IN OUTSIDE AREAS: ANTS*, BOXELDER BUGS,
 BROWN DOG TICKS*, CHIGGERS*, CLOVER MITES*, COCKROACHES,
 CRICKETS*, CARMES*, FLEAS*, FLIES*, MILLIPEDES*,
 SONBUGS (PILLBUGS)*, SPIDERS, SPRINGTAILS (COLLEMBOLA)*, 4.
 8 FL. OZ. IN 3 GALS. OF WATER.
 THOROUGHLY SPRAY WINDOW AND DOOR SCREENS, SILLS, FOUNDATIONS,
 PORCH AND PATIO WALLS AND CEILINGS, ENTRANCEWAYS, WALKS,
 GARBAGE CANS, TREE TRUNKS, AND INTO CRACKS AND OTHER PLACES
 WHERE INSECTS CAN HIDE.
 *DO NOT TREAT PLANTS OR ANIMALS.
 RE-SPRAY A 5 FT. BAND OF SOIL AROUND THE HOUSE NEXT TO THE
 FOUNDATION AND SPRAY THE FOUNDATION OF THE HOUSE TO A HEIGHT
 OF 1 FT.

ADVICE: RECOMMENDATIONS FOR OTHER CONTROL MEASURES FOR
 PESTS.

FIRES ANTS: TO AID IN THE CONTROL OF FIRE ANTS IN LAWNS AND
 OTHER RECREATION AREAS, APPLY 1 FL. OZ. (2 TSP.) IN A MINIMUM
 OF 1 GAL. OF WATER AS A DRENCH. APPLY OVER AND 6 INCHES APART
 EACH MOUND. APPLY GENTLY TO AVOID DISTURBING ANTS. REAPPLY AS
 NECESSARY. LARGER QUANTITIES OF SOLUTION CAN BE PREPARED FOR
 MULTIPLE MOUND TREATMENTS. USE EQUIPMENT CAPABLE OF DELIVERING
 THE DILUTED SPRAY AS A GENTLE RAIN AT THE RATE OF 1 GAL./PER
 MOUND. HIGH PRESSURE SPRAYS MAY DISTURB THE ANTS AND CAUSE
 MIGRATION, REDUCING PRODUCT EFFECTIVENESS.

NOTE: FOR BEST RESULTS, APPLY IN COOL WEATHER, 65-68°F, OR IN
 THE MORNING OR LATE EVENING IN SUMMER. TREAT NEW MOUNDS AS THEY

DIRECTIONS FOR USE (CONTINUED)
 LAWN: BENT, BERMUDA, BLUEGRASS, BUCHONIA, FESCUE, FINGER GRASS, ST. AUGUSTINE, ANTS, ARMYWORMS, BERMUDA GRASS THRIP, DAY DOG TICKS, CHIGGERS, CRICKETS, COTTERWORMS, EARWIGS, FLIES, FRUIT FLIES, GARDEN MOTHS (CODELLOTHORACIS), LEAFHOPERS, MITES (BERMUDA, CLOVER), SONBUGS (PILLBUGS), SPRINGTAILS (COLLEMBOLA)

1 FL. OZ. IN 3 GALS. OF WATER 1,000 SQ. FT.

THOROUGHLY WET DOWN GRASS A FEW HOURS BEFORE APPLYING.
 BROWN DOG TICKS. DO NOT APPLY TO ANIMALS. IN PARTICULAR,
 SPRAY UNDER SHRUBBRY AND ON THE GRASS NEXT TO HOUSE.

XXLAWN BILLBUGS: WATER THE LAWN THOROUGHLY IMMEDIATELY
 AFTER TREATMENT.

*IN THE NORTH, IF CHINCH BUGS ARE THE ONLY LAWN INSECT
 PROBLEM, A LOWER RATE OF APPLICATION CAN BE USED: 1/2 FL. OZ.
 IN 3 GALS. OF WATER COVERS 125 SQ. FT.

MILLIPEDES: 1 FL. OZ. IN 3 GALS. OF WATER COVERS 125 SQ. FT.
 THOROUGHLY WET GRASS A FEW HOURS BEFORE APPLYING.

RHODESGRASS SCALES: 1 FL. OZ. IN 3 GALS. OF WATER COVERS 100 SQ. FT. APPLY WHEN SCALE CRAWLERS FIRST EMERGE.

ANTS, DIGGER WASPS: 1 FL. OZ. IN 3 GALS. OF WATER COVERS 125 SQ. FT. SPOT SPRAY ANT HILLS AND WASP GROUND NEST TO OPENINGS.

MAKE THE APPLICATION WHEN THE WASPS ARE NOT ACTIVE.

BILLBUGS, WHITE GRUBS OF JAPANESE BEETLE, EUROPEAN CHAFER, SOUTHERN CHAFER: 1 FL. OZ. IN 3 GALS. OF WATER COVERS 125 SQ. FT. WATER GRASS THOROUGHLY AFTER APPLICATION.

TRILLING CONTROL: APPLY WHEN ACTIVITY IS FIRST OBSERVED OR WHEN
 CHINED OR BROWN GRASS IS PRESENT, INDICATING DAMAGE FROM THE
 INSECT.

GRUB CONTROL: MAKE APPLICATION ANY TIME BETWEEN LATE JULY AND
 EARLY OCTOBER.

HYPEDOES NEVEIL (NEW YORK ONLY): 3/8 FL. OZ. IN 3 GALS. OF
 WATER COVERS 125 SQ. FT. WATER GRASS THOROUGHLY AFTER APPLICATION.
 TREAT PROBLEM AREAS IN MID-APRIL AND AGAIN IN MID-JUNE.

TREES & SHRUBS: ARBORVITAE, AZALEA, BIRCH, BOUDOIS, CAMELLIA,
 DOUGLAS FIR, ELM, HAWTHORN, HOLLY, JUNIPER, LILAC, LOCUST, MAPLE,
 MINDA, GINKGO, PINE, PLUM, POPLAR, RHODODENDRON, SPRUCE, WILLOW.
 WET LEAVES AND BRANCHES TO THE DРИPPING POINT. TRY TO PENETRATE
 DENSE FOLIAGE.

APHIDS, BAGWORMS, EUROPEAN PINE SHOOT MOTH, HOLLY BUD MOTHS,
 DIPTEROUS LEAFROLLERS (ARBOVITAE, AZALEA, BIRCH, BOUDOIS,
 HOLLY, GINKGO), LEAFHOPPER, MEALYBUGS, MITES (COTTON, CYCLAMEN,
 EUROPEAN RED, PRIVET, TWOSPOTTED SPIDER), GELSEME ROOT
 MAGGOTS, CITRUS LEAFROLLERS, SCALE CRAWLERS (COTTON, CUSHION,
 EUONYMUS, LEGUMINOSA, PINE NEEDLE, PIT SCALE, SHILOH, SOFT).

1 FL. OZ. IN 3 GALS. OF WATER. APPLY AS A THOROUGH FOLIAR
 SPRAY.

MEALYBUGS (CITRUS, SHILOH), WHITEFLIES: 1 FL. OZ. IN 3 GALS.
 OF WATER. APPLY AS A THOROUGH FOLIAR SPRAY.

PHONSA MEALYBUGS: APPLY AS A THOROUGH FOLIAR SPRAY WHEN ADULTS
 FIRST APPEAR. MAKE TWO ADDITIONAL APPLICATIONS AT 4-5 WEEK
 INTERVALS.

APPLE-AND-THREE SKELETONIZERS, HEMLOCK CHERRIES, SPA LICHENS,
 OBLIQUE-BANDED LEAFROLLERS, PEAR SLUGS, TENT CATERPILLARS,
 HERBADS (COTONEASTER, FALL): 3 FL. OZ. IN 3 GALS. OF
 WATER. APPLY AS A THOROUGH FOLIAR SPRAY.

ROSES & FLOWERS: AZALEAS, CAMELLIAS, DYESMANTHEMUM, CARNATION,
 GLADIOLI, LILAC, RHODODENDRON, WISTERIA. WET THE PLANTS TO THE
 DРИPPING POINT. TRY TO PENETRATE LEAVES AND PENETRATE
 DENSE FOLIAGE.

APRICOT, CHERRY SCALE CRAWLERS, PEA FEELETTES, STUPENDA, LEAF
 MINERS, LEAFHOPPER, MEALYBUGS, MITES (COTTON, BUD, CITRUS,
 SPANISH, CYCLAMEN, EUROPEAN RED, TWOSPOTTED SPIDER), WHITEFLIES,
 1 FL. OZ. IN 3 GALS. OF WATER. APPLY AS A

THOROUGH FOLIAR SPRAY.

NOTE: DO NOT USE ON FEIERS, FOINSETTEA, MITES, PINEAPPLE,
 PAPAYA, PILEA, GARDENIA, AND ORCHIDS. SINGLE INJURY
 MAY OCCUR.

FRUITFUL, JEWEL, RIBES, RIBES, RIBES, RIBES, WALNUTS, LAMP
 SEPARATING AFTER DILUTION. APPLICATION SHOULD BE MADE ACCORDING
 TO INSTRUCTIONS FOR HIGH APHIDS, MITES, AND CRAWLER STAGES OF
 SCALE INSECTS. ALL STAGES KILLED. IN PESTICIDES INSECTICIDE, REPEAT
 APPLICATION AT 7-10 DAY INTERVALS.

APHIDS, NEAT COTTONSEED (PEAS), SWEET COTTONSEED, SPIDER MITES
 1/2 FL. OZ. IN 3 GALS. OF WATER. APPLY AS A FOLIAR SPRAY.
 DO NOT APPLY AFTER SUNSET OR ON SPLEEN.

NOTE: SEE ALSO FOR FRUITFUL IN NEXT COLUMN.



Diazinon® 25E

LAWN & GARDEN INSECT CONTROL

ACTIVE INGREDIENTS:
O,O-DIETHYL O-(2-ISOPROPYL-6-METHYL-4-PYRIMIDINYL) PHOSPHOROTHIOATE..... 25.0%
AROMATIC PETROLEUM DERIVATIVE SOLVENT..... 55.4%
INERT INGREDIENTS:..... 2.6%
TOTAL 100.0%

CONTAINS DIAZINON INSECTICIDE SPRAY CONCENTRATE
DIAZINON IS A REGISTERED TRADEMARK OF CIBA-GEIGY.

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED-CALL A PHYSICIAN IMMEDIATELY. GASTRIC LAVAGE IS INDICATED IF MATERIAL WAS TAKEN INTERNALLY. DO NOT INDUCE VOMITING UNLESS OTHER TREATMENT IS NOT AVAILABLE. VOMITING MAY CAUSE ASPIRATION PNEUMONIA. IF IT IS NECESSARY TO INDUCE VOMITING, GIVE VICTIM ONE OR TWO GLASSES OF WATER AND INSERT FINGER IN BACK OF THROAT. REPEAT UNTIL VOMIT IS CLEAR. DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.
IF INHALED- REMOVE VICTIM TO FRESH AIR. APPLY RESPIRATION IF INDICATED.
IF ON SKIN- WASH CONTAMINATED SKIN WITH SOAP AND WATER.
IF IN EYES- FLUSH EYES WITH PLENTY OF WATER FOR AT LEAST 15 MINUTES. GET MEDICAL ATTENTION IMMEDIATELY.

EPA REG. NO. 8612-101 NET CONTENTS 1 GALLON
EPA EST. NO. 8612-TX-1,2;OK-1 LABEL NO. 012802



10539 MAYBANK
DALLAS, TEXAS 75220

DIRECTIONS FOR USE (CONTINUED)

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.
APPLIES TO: APPLES, CHERRIES, CRANBERRIES, GRAPEFRUIT, LEMON, LIME, MELON, PEACHES, PLUM, RASPBERRIES, STRAWBERRIES, TURNIPS, TURNIP RAPERS, TURNIP TOPS, TURNIPS, TURNIPS (CROWN + EUROPEAN RED).
1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.
BERRIES (BLACKBERRIES, BOYSBERRIES, CRANBERRIES, LOGAN-BERRIES, RASPBERRIES); APHIDS, LEAF-HOPPERS, RASPBERRY FRUITWORMS, TWO-SPOOTTED MITES, THrips.
1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.
CHERRIES, APRICOTS, PLUM, MEALY PLUM, THISTLE, MELON (CROWN + TURNIP + EUROPEAN RED).
1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.
CURRANTS, APHIDS, BLUEBERRY MAGGOT, CHERRY FRUITWORMS, CHERRY FRUITWORMS, CHERRY FRUITWORMS (MAGGOTS), TWO-SPOOTTED MITES, THrips.

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

STRAWBERRIES: APHIDS, TWO-SPOOTTED MITES, STRAWBERRY LEAFROLLERS

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

GRAPE: APHIDS, GRAPE BERRY MOTH, GRAPE LEAFWORMS, LEAFWORMS, SCALE CRABSPOTS.

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

GRAPEFRUIT, LEMONS, GRAPES, CITRUS THRIPS, SCALE CRABSPOTS (CITRUS SPIDER, CITRUS SPIDER)

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

PEACHES, STRAWBERRIES, PLUMS, APRICOTS, CHERRIES, GRAPEFRUIT, LEMONS, MELON, PEACHES, PLUMS, PRUNES, RASPBERRIES, STRAWBERRIES, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

VEGETABLES: CANTALOUPE, MUSKELONG, WINTER SQUASH, WATERMELON, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.

BEANS, CARROTS, CORN, PEAS, POLE BEANS, RED BEETS, TURNIPS, TURNIP RAPERS, TURNIPS, TURNIPS (CROWN + TURNIP + EUROPEAN RED).

1 FL. OZ. IN 3 GALLONS OF WATER. APPLY AS A FULL COVER SPRAY.