

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

May be fatal if swallowed or absorbed through skin. Avoid breathing spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly if on skin and after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto sprayed grass until spray has completely dried. Do not use in the home. Do not contaminate ornamental fish ponds.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, shrimp, crab, birds and other wildlife. Birds feeding on treated area may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, crop cover, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Store in a cool, dry place, inaccessible to children and pets. PROMISSIONS: Do not contaminate water, food, or feed by storage or disposal. PESTICIDE DISPOSAL: Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility. CONTAINER DISPOSAL: Triple rinse (or equivalent), then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state or local authorities, by burning. If burned, stay out of smoke. DO NOT STORE NEAR HEAT OR OPEN FLAME.

IMPORTANT

Seller's guarantee shall be limited to the terms of this label and, subject thereto, the buyer assumes any risks to persons or property arising out of the use or handling and accepts the product on these conditions.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

NUISANCE PESTS IN OUTSIDE AREAS: Ants, Boxelder bugs, Brown Dog ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (Collembola). 8 fl. oz. in 3 gals. of water.

Thoroughly spray window and door screens, sills, foundations, porch and patio walls and ceilings, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.

DO NOT TREAT PLANTS OR ANIMALS. Spray a 5 ft. band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 ft. See lawn recommendations for other control measures for these pests.

FIRE ANTS: To aid in the control of Fire Ants in lawns and other recreation areas, apply 1 fl. oz. (2 tbsp.) in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted spray as a gentle rain at the rate of 1 gal. per mound. High pressure spray may disturb the ants and cause migration, reducing product effectiveness.

NOTE: For best results, apply in cool weather, 65-80°F. or in early morning or late evening hours. Treat new mounds as they appear.

LAWNS: Bent, Bermuda, Bluegrass, Dichondra, Fescue, Meron, St. Augustine, Ants, Armyworms, Bermudagrass Thrips, Brown Dog Ticks, Chiggers, Crickets, Cutworms, Earwigs, Fleas, Frit Flies, Lawn Moths (Sod Webworms), Leafhoppers, Mites (Bermuda Clover), Sowbugs (Pillbugs), Springtails (Collembola). 1 fl. oz. in 3 gals. of water covers 1,000 sq. ft.

Thoroughly wet down grass a few hours before applying. Brown Dog Ticks: Do not apply to animals. In particular, spray under shrubbery and the grass next to house.

LAWN BILLBUGS: Water the lawn thoroughly immediately after treatment.

In the north, if Chinch Bugs are the only lawn insect problem a lower rate of application can be used: 1 fl. oz. in 3 gal. of water covers 125 sq. ft.

MILLIPEDES: 2 fl. oz. in 3 gals. of water covers 125 sq. ft. Thoroughly wet grass a few hours before applying.

ANTS, DIGGER WASPS: 1 fl. oz. in 3 gals. of water covers 125 sq. ft. Spot spray ant hills and wasp ground next to openings. Make the application when the wasps are not active.

BILLBUGS, WHITE GRUBS, JAPANESE BEETLE, EUROPEAN CHAFER, SOUTHERN CHAFER: 1 fl. oz. in 3 gals. of water covers 125 sq. ft. Water grass thoroughly after application.

GRUB CONTROL: Make application any time between late July and early October.

Aphids, Bagworms, European Pine Shoot Moths, Holly Bud Moths, Dicotylous Leafminers (Arborvitae, Aralia, Birch, Boxwood, Holly, Oak), Leafhopper, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Obscure Root Weevils, Omnivorous Leafhoppers, Scale Crawlers (Cottony-cushion, Euonymus, Lecanium, Pine Needle, Pit Scale, San Jose, Soft).

1 fl. oz. in 3 gals. water. Apply as a thorough foliar spray. WEBWORMS (JUNIPER, MIMOSA), WHITEFLIES: 1 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray.

MIMOSA WEBWORMS: Apply as a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals. Apple and Thorn Skeletonizers, Hemlock Chermes, Oak Looper, Oblique-banded Leafroller, Pear Slugs, Tent Caterpillars, Webworms (Cottonweaver, Fall): 3 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray.



THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS

Diazinon 25E

Lawn & Garden Insect Control

Table with 2 columns: ACTIVE INGREDIENTS and INERT INGREDIENTS. Includes Diazinon, Pyridinal, and Phosphorothioate.

Contains Diazinon insecticide spray concentrate

Diazinon is a registered trademark of CIBA GEIGY.

WARNING

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician immediately. Gastric lavage is indicated if material was taken internally. Do not induce vomiting unless other treatment is not available. Vomiting may cause aspiration pneumonia. If it is necessary to induce vomiting, give victim one or two glasses of water and insert finger in back of throat. Repeat until vomit is clear. Do not induce vomiting or give anything by mouth to an unconscious person. Remove victim to fresh air. Apply respiration if indicated. If On Skin: Wash contaminated skin with soap and water. If in Eyes: Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

Net Contents 1 Gallon

Label # 923002



COMPANY DALLAS, TEXAS 75220

DIRECTIONS FOR USE (Continued)

ROSES AND FLWERS: Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilacs, Rhododendrons, Roses. Wet the plants to the dripping point. Try to hit underside of leaves and penetrate dense foliage. Aphid, Euonymus Scale Crawlers, Flea Beetles, Dipterous Leafminers, Leafhopper, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies. 1 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray. NOTE: Do not use on Ferns, Poinsettia, Hibiscus, Papyrus, Pilea, Gardenia, and Orchids since injury may occur.

FRUITS AND NUTS: Almonds, Filberts, Pecans, Walnuts. Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted. If infestations warrant, repeat applications at 7-10 day intervals. Aphids, Nut Casemakers (Pecans), Scale Crawlers, Spider Mites: 1 fl. oz. in 3 gals. of water. Apply as a full cover spray. Do not apply after husks open or split. APPLES, PEARS: Aphids, Codling Moths, Eyespotted Bud Moths, Fruit Tree Leafrollers, Leafhoppers, Tentiform Leafminers (pears), Maggots (Apples), Mealybugs, Mites (European Red, Clover (brown), Pacific, Two-spotted spider), Psylla (pears), Scale Crawlers (Cottony-cushion, San Jose, Forbes, Soft). 1 fl. oz. in 3 gals. water. Apply as a full cover spray. C. Apples, slight ruseting may occur on some varieties such as Golden Delicious.

STRAWBERRIES: Aphids, Two-spotted Mites, Strawberry Leafrollers. 1 fl. oz. in 3 gals. water. Apply as a full cover spray. GRAPES: Aphids, Grape Berry Moths, Grape Leafrollers, Leafhoppers, Scale Crawlers. 1 fl. oz. in 3 gals. water. Apply as a full cover spray. GRAPEFRUIT, LEMONS, ORANGES: Citrus Thrips, Scale Crawlers (Citrus Snow, Cottony-cushion). 1 fl. oz. in 3 gals. of water. Apply as a full cover spray. *12 fl. oz. in 3 gals. water. Apply as a full cover spray after blooming and again in the summer. Thorough spray coverage of tree trunk, limbs and twigs is essential.

AFTER SPRAYING, DO NOT PICK FRUIT WITHIN

- 5 days strawberries, figs
7 days blueberries, cranberries
10 days apricots, cherries, grapes, nectarines, plums, prunes
14 days apples, pears
21 days grapefruit, lemons, oranges, peaches

VEGETABLES: Foliar application. Beans, Peas. Thoroughly spray all plant parts. Generally, applications at 7-10 day intervals may be required. Do not feed treated vines to livestock. Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Mites. 1 fl. oz. in 3 gals. water covers approx. 1,000 sq. ft. of garden. PEPPER, POTATOES, TOMATOES: Aphids, Colorado Potato Beetles, Fall Armyworm (Tomato), Flea Beetles, Leafhoppers, Southern Armyworms (Potato and Tomato), Vinegar Flies (Drosophila Spp Tomato). 1 fl. oz. in 3 gals. water covers approx. 1,000 sq. ft. of garden. CANTALOUPE, CUCUMBERS, MUSKMELON, SQUASH (SUMMER, WINTER), WATERMELON: Aphids, Thrips, Mites. 1 fl. oz. in 3 gals. of water covers approx. 1,000 sq. ft. of garden.

ONIONS: Onion Thrips. 1 fl. oz. in 3 gals. of water covers approx. 1,000 sq. ft. of garden.

SWEET CORN: Aphids, Flea Beetles, Grasshoppers, Mites. 1 fl. oz. in 3 gals. of water covers approx. 1,000 sq. ft. of garden.

VEGETABLES: PREPLANT SOIL APPLICATION. corn, lettuce, peas, pole beans, radish, tomatoes, turnips. Cutworms (surface). 6 fl. oz. per 1,000 sq. ft. in sufficient coverage. Apply broadcast just before planting. 2-3 inches for surface cutworms, 3-6 inches for cutworms. GARDEN SYMPHALANS (GARDEN CEE): 15 fl. oz. per 1,000 sq. ft. in sufficient coverage. Apply broadcast just before planting. 4-8 inches.

MOLT CRICKETS: Carrots, corn, peas and snap beans only. 1.5 fl. oz. per 1,000 sq. ft. in sufficient coverage. Apply broadcast just before planting. Sow or transplanting. Immediately.

ROOT MAGGOTS: Cabbage only. 4.5 fl. oz. per 1,000 sq. ft. in sufficient coverage. Apply broadcast just before planting. 3-4 inches before seeding, either in soil or transplanting.

WIREWORMS: Cabbage, carrots, corn, and tomatoes only. 6 fl. oz. per 1,000 sq. ft. in sufficient coverage. Apply broadcast just before planting. 4-8 inches.

AFTER SPRAYING, DO NOT PICK VEGETABLES WITHIN

- 1 day peas, tomatoes, sweet corn
3 days cantaloupe, muskmelon, winter squash, cauliflower, paprika, beans, cabbage, cucumbers,
7 days carrots, celery, collards, onions, parsnips, radishes,
10 days potatoes

HOW TO MIX AND APPLY

STEP 1: Determine the size of spray; vegetables, measure the area to be sprayed. applications, apply as a thorough coverage when insects first appear. Repeat application in the specific site and pest direction harvest intervals for fruits and vegetables.

STEP 2: Use clean sprayer. Carefully amount of B & C Diazinon 25E and water chart.

STEP 3: Spray as directed in rate of sprays for lawn, soil, and other surface mist sprays for trees and plants. important. Flush sprayer with clean water. 1 fl. oz. 6 teaspoons or 2 tablespoons. 1 fl. oz. in 3 gals. of water. 2 teasp.