



MULTI-PURPOSE INSECTICIDE S.E.C.

Combining Two Highly Active Synthetic Pyrethroid Compounds

A WATER EMULSIFIABLE INSECTICIDE CONCENTRATE

For use in Bakeries, Bottling Plants, Canneries, Flour Mills,
Food Processing Plants, Hospitals, Hotels, Restaurants, Schools, Horse Barns,
and on Ornamental Plants in Homes, Gardens and Greenhouses

ACTIVE INGREDIENTS

*Tetramethrin	2.50%
Related Compounds	0.34%
Resmethrin [(5-phenylmethyl)-3-furanyl] methyl-2,2- dimethyl-3-(2-methyl-1-propenyl) cyclopropanecarboxylate	2.50%
Related Compounds	0.46%
Petroleum Distillate	104.00%
INERT INGREDIENTS	86.20%
Total	100.00%

*1-cyclohexene-1, 2-dicarboximidomethyl 2,2-dimethyl-3-
(2-methylpropenyl) cyclopropanecarboxylate

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Side Panel for Additional Precautions

EPA REG. NO. 8612-117
EPA EST. NO. 8612-TX-1

Manufactured by



COMPANY
DALLAS, TEXAS 75220

NET CONTENT

GALLONS

Label No. MISEC-0287-01

DIRECTIONS FOR USE (continued)

TO CONTROL houseflies, mosquitoes, gnats and flying moths. Close doors and windows and shut off ventilating systems. Apply as a fog or fine mist directing the spray toward the upper portions of the room and toward windows and other light sources which attract these insects. Thoroughly fill the area with mist using 1/2 to 1 ounce per 1000 cubic feet of space. Keep area closed for at least 10 minutes. Vacate after treatment and ventilate before reoccupying. Repeat treatment as needed.

ORNAMENTAL PLANTS

Dilute this concentrate at the rate of 1 part material to 79 parts water.

TO CONTROL aphids, two-spotted spider mites and adult whiteflies on ornamental house and garden plants (African violet, azalea, Christmas cherry, chrysanthemum, ciperaria, diffenbachia, fuschia, gloxinia, ivy, lantana, Neanthe bella palm, petunia, rose and snapdragon and on outdoor plants such as azalea, chrysanthemum, Quonymus, ivy, marigold, petunia, rose, snapdragon and zinnia) Apply as a fog or fine mist, spraying plants with a sweeping motion so that the mist drifts into the plants. Direct spray so that the upper and lower portions of leaves are contacted. Caution on new growth and tender foliage. When using outdoors, spray when air is still. Repeat as necessary.

TO CONTROL mealybugs and leafminer: Dilute at the rate of 1 part concentrate to 39 parts water and apply in the same manner as given above.

TO CONTROL gypsy moths and caterpillars, forest and tent caterpillars, canker worms, garden webworms, and the above pests on ornamental trees: Spray as above. Do not spray bearing fruit trees.

IN HORSE BARN AND ON HORSES

TO PROVIDE quick relief from annoying flies, mosquitoes and gnats: Dilute 1 part concentrate with 19 parts water and apply as a fine mist spray to stable area, and over and around stabled horses. May also be sprayed directly on horses. DO NOT USE ON HORSES INTENDED FOR HUMAN CONSUMPTION.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store upright at room temperature. Avoid exposure to extreme temperatures. In case of spill or leakage, soak up with an absorbent material such as sand, sawdust, earth, fuller's earth, etc. Dispose of with chemical waste.

PESTICIDE DISPOSAL: Waste resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent) then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other approved state and local procedures.

NOTICE

Sellers guarantee shall be limited to the terms of this label, and subject thereto, the buyer assumes any risks to persons or property arising out of the use or handling and accepts the product on these conditions.

BEST AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. The use of this product in food processing plants should be confined to time periods when the plant is not in operation. Foods should be removed or covered during treatment and thoroughly cleaned and rinsed with potable water before using. Wash all contaminated clothing with soap and hot water before reuse. Remove pets, birds and cover fish aquaria before spraying. When used in hospitals, patients must be removed from the room and the room aired for two hours prior to re-entry.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Contact a physician or Poison Control Center immediately. Do not induce vomiting. Vomiting may cause aspiration pneumonia.

IF INHALED: Remove affected person to fresh air. Apply artificial respiration if indicated.

IF ON SKIN: Remove contaminated clothing and wash affected areas of skin with soap and water. Contact a physician if irritation persists.

IF IN EYES: Flush with plenty of water. Contact a physician if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply outdoors when weather conditions favor drift from area treated. This product is toxic to fish and birds. Do not apply directly to lakes, ponds or streams. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Water dilutions should be prepared at room temperature. Exposure to low temperatures may cause cloudiness or crystallization.

SURFACE SPRAY

This concentrate should be diluted with water before use, at a rate of 1 part concentrate to 9 parts water. The diluted spray should be applied in conventional, mechanical or compressed air spraying or fogging equipment. It is not recommended for use in thermal generating equipment.

TO CONTROL cockroaches, ants, spiders, silverfish, crickets, centipedes, earwigs, sowbugs and grain mites: Apply as a coarse droplet spray directed into cracks, crevices and other hiding places, along baseboards, around the bases of machinery and under pallets. Treat all returned bottle cartons and cases. Spray thoroughly, contacting insects directly whenever possible. Spray badly infested areas daily. In other areas, repeat treatment twice weekly or when reinfestation occurs.

SPACE SPRAY

This concentrate should be diluted with water before use, at a rate of 1 part concentrate to 19 parts water. The diluted spray should be applied in conventional, mechanical or compressed air spraying or fogging equipment. It is not recommended for use in thermal generating equipment.