


8622-13
 8622-13

RESTRICTED USE PESTICIDE
 For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



67-33

PREPLANT SOIL FUMIGANT

ACTIVE INGREDIENTS:	By W
Methyl bromide	67
Chloropicrin	33
Total	100

For control of nematodes, wireworms, weed seed, fungi diseases, grubs, ants and soil-borne insects. (Example: white grubs)

DANGER
KEEP OUT OF REACH OF CHILDREN
POISON

SEE SIDE OF CAN FOR ANTIDOTE & WARNINGS ON SIDE PA

NET WEIGHT _____ LBS.

EPA Reg No 8622-13

LOT NO _____

AMERIBROM INC
 1250 Broadway
 New York, N.Y. 10001

Many pesticidal chemicals are poisonous and may leave a toxic residue on the plants to which they are applied. The U. S. Food and Drug Administration has established maximum amounts of such pesticidal chemicals that may remain on raw agricultural products at harvest, and it is the user's responsibility to see that there is no residue on such crops at harvest in excess of these amounts. The "Fumigation Directions" are based on the best available information, and if followed carefully should not leave excessive residues at harvest. However, AMERIBROM INC. assumes no responsibility as to their accuracy nor for any loss due to excessive residues.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

Empty cylinders should be returned freight collect to AMERIBROM INC. location from which the same was purchased. Consult local sales office if questions arise.

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.

67-33

PREPLANT SOIL FUMIGANT

ACTIVE INGREDIENTS:	By Wt.
Methyl bromide	67%
Chloropicrin	33%
Total	100%

For control of nematodes, wireworms, weed seed, fungus diseases, grubs, ants and soil-borne insects. (Example: white grubs)



DANGER
KEEP OUT OF REACH OF CHILDREN
POISON

DO NOT INHALE VAPORS • SEE ANTIDOTE & WARNING SECTION

NET WEIGHT _____ LBS

EPA Reg. No. 552-2-13

LOT NO. _____

AMERIBROM, INC.
125 W. Broadway
New York, N.Y. 10038

67-33 FUMIGATION DIRECTIONS RECOMMENDED APPLICATION RATES

Soil treatment for Strawberries 225 to 350 lbs Acre

Expose to fumigation for 48 hours. Aerate 2 weeks before setting transplants in treated areas

Soil treatment for Tomatoes 225 to 325 lbs Acre

Expose to fumigation for 48 hours. Aerate 2 weeks before setting transplants in treated areas

CITRUS SOIL FOR CONTROL OF PHYTOPHTHORA IN FLORIDA SANDY SOILS

This is a preplant or replant treatment. Trees which are planted in this treated soil will not have harvestable fruit for a period of at least 24 months. Use a minimum of 1 1/2 lb/100 sq. ft. Expose to fumigation for 96 hours covering treated area with a 4 mil tarpaulin. Will control disease to a depth of 4 feet. Aerate 2 weeks before setting transplants in treated area

Apply with chisels spaced 12 inches apart to a depth of 8 inches. Seal fumigant with a drag or cult plow immediately following chisels. Cover with a gas proof tarpaulin. Remove tarpaulin after 96 hours and fumigant in period and aerate soil for 2 weeks before planting. This is persistent in the soil for 24 months.

Do not apply to soil with high water content or high organic matter.

Do not apply to soil with high water content or high organic matter.

Do not apply to soil with high water content or high organic matter.

POISON
DANGER

HARMFUL TO HUMANS AND DOMESTIC ANIMALS. See Antidote & Warning Section. POISON

See Antidote & Warning Section.

NOTICE
Following soil fumigation, undesirable concentrations of chloropicrin may drift to nearby areas. If this occurs, immediately cover treated area with a plastic tarpaulin. Leave tarpaulin on overnight and remove during the daytime. Recover the area if eye irritation indicates that the undesirable concentration is continuing. These concentrations are most likely to occur in the evening or at night when the air is most likely to be static in movement. Do not make application of 67-33 when there is little or no air movement or there is an inversion condition in the atmosphere.

NOTE CAREFULLY Following fumigation, the level of soluble salts and ammonia nitrogen may be raised. This is most likely to occur when heavy rates of fumigant and fertilizers are applied to soils that are either acid, wet, cold, or high in organic matter. Fertilizers containing ammonium salts should not be used. Apply only fertilizers containing nitrates until after the crop is well established and the soil temperature is above 65° F. in order to avoid ammonia injury, nitrate starvation, or both. After fumigation, fertilize as indicated by soil test to protect against injury to plant roots. It is best that highly acid soils be limed before application to promote soil fertility and reduce the possibility of ammonia toxicity. 048