

8622-68

7/9/2008

10/18



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
Washington, D.C. 20460

July 9, 2008

OFFICE OF
PREVENTION, PESTICIDES
AND TOXIC SUBSTANCES

Judy A. Smith
N. American Regulatory Affairs Manager
AmeriBorn, Inc.
95 MacCorkle Ave., SW
South Charleston, WV 25303

Subject: Product Name: Calirus 150
EPA Reg. No: 8622-68
Application for Label Notification Dated April 30, 2008 to Change Company
Name to ICL-IP America, Inc. Effective July 1, 2008

Dear Ms. Smith:

The Biopesticides and Pollution Prevention Division is in receipt of your application for Notification under 98-10 dated above. A preliminary screen of this request has been conducted for its applicability under PRN 98-10 and it has been determined that the action(s) requested falls within the scope of PRN 98-10. Our records have been duly noted, and the label submitted with this application has been stamped "Notification, received and reviewed" and will be placed accordingly in our records.

Questions concerning this action should be directed to Ms. Diana Hudson at (703) 308-8713 or email at HUDSON.DIANA@EPA.GOV.

Sincerely,

Linda Hollis

Linda Hollis, Chief
Biochemical Pesticides Branch
Biopesticides and Pollution Prevention
Division (7511P)

20f18

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060



United States
Environmental Protection Agency
Washington, DC 20460

Registration
 Amendment
 Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 8622-68	2. EPA Product Manager	3. Proposed Classification <input type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) CALIRUS 150	PM# 32	
5. Name and Address of Applicant (Include ZIP Code) AmeriBrom, Inc. 95 MacCorkle Ave., SW South Charleston, WV 25303 <input type="checkbox"/> Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3)(b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application. NOTIFICATION
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below

Date Reviewed: 7-9-08

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Reviewed By: D. Hudson

Notification of Company Name Change (EPA Reg. No. 8622) to ICL-IP America, Inc. per PR Notice 98-10.

Section - III

1. Material This Product Will Be Packaged In:				2. Type of Container	
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No		<input type="checkbox"/> Metal	<input type="checkbox"/> Plastic
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container	<input type="checkbox"/> Glass	<input type="checkbox"/> Paper
				<input type="checkbox"/> Other (Specify) _____	
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product		<input type="checkbox"/> Lithograph <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled		<input type="checkbox"/> Other _____	

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name Judy A. Smith	Title NA Regulatory Affairs Mgr	Telephone No. (Include Area Code) 304.720.3929
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.		6. Date Application Received (Stamped)
2. Signature 	3. Title NA Regulatory Affairs Mgr.	
4. Typed Name Judy A. Smith	5. Date April 30, 2008	



AMERIBROM INC. 2115 Linwood Avenue, Suite 200 Fort Lee, New Jersey 07024-5004, USA
Tel: (1) 2012426560, Fax: (1) 2012426561; www.icl-industrial.com, info@ameribrom.icl-ip.com

30418

May 1, 2008

U.S. Environmental Protection Agency
Office of Pesticide Programs
Document Processing Desk (COADR)
1200 Pennsylvania Avenue N.W
Ariel Rios Building
Washington DC 20460-0001
ATTENTION: Kathleen O'Malley

Certified Mail
7008 0150 0002 9518 4732
FAX 703-305-7670

Dear Ms. O'Malley:

**SUBJECT: Notice of Change of Company Name for Ameribrom Inc.
(EPA Company No. 8622 and 15298)**


Effective July 1, 2008, Ameribrom Inc. will change the corporate name to **ICL-IP America Inc.** The current corporate address and contact information will remain the same, i.e., 95 MacCorkle Avenue SW, South Charleston WV 25303-1411. No exchange of assets has occurred and no company-to-different-company registration transfers are to be done as a consequence of this name change; per our conversation on April 4, 2008, this is a simple corporate name change. This company name change will impact all Ameribrom federal registrations possessing a company number of **8622** or **15298**. We will be filing updated product labels to reflect the corporate name change from "Ameribrom Inc." to "**ICL-IP America Inc.**" in May 2008.

The telephone number for Ameribrom Inc. will remain as 304-720-3950. The Ameribrom fax number at the South Charleston location is 304-746-3101.

Please update the files at EPA to reflect the name change and notify the appropriate EPA product teams of this change.

If you have questions regarding this notice of change of corporate name, please call me at 304-720-3929.

Sincerely yours,



Judy A. Smith, Ph.D.
Ameribrom Inc.

ICL-IP America Inc. (effective 7/1/2008)
North American Regulatory Affairs Manager



Caring for your future today



8622-68

06/30/2005

Notification - made using 4 of 18
last stamped, EPA-
approved label. 1/15

CALIRUS 150 revised two sections to EPA 05 June 1

CALIRUS 150

A FUNGICIDE FOR THE CONTROL OF *PYTHIUM*, *PLASMODIOPHORA*, DOWNY MILDEWS, *ALTERNARIA* AND *PHYTOPHTHORA* ON AGRONOMIC CROPS.

ACTIVE INGREDIENTS

Mono- and dipotassium salts of Phosphorous Acid*.....	10.30%
Copper Sulfate.....	0.15%
OTHER INGREDIENTS.....	<u>89.55%</u>


TOTAL..... 100%

*Contains 1.02 lbs/gallon of active ingredients, mono- and dipotassium salts of Phosphorous Acid. Equivalent to 0.58 lbs. Phosphorous Acid/gallon.

KEEP OUT OF REACH OF CHILDREN

CAUTION

CUIDADO



Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

FIRST AID	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.
<p>HOT LINE NUMBER</p> <p>Have the product container or label with you when calling a poison control center or doctor, or going for treatment.</p> <p>YOU MAY ALSO CONTACT 1-800-420-9236 FOR EMERGENCY MEDICAL TREATMENT INFORMATION.</p>	

See side panels for additional precautionary statements.

50118
2/15

CALIRUS 150 revised two sections to EPA 05 June 1

Manufactured for:

MADE IN ISRAEL

~~Ameribrom, Inc.~~
~~2115 Linwood Ave., Suite 200~~
~~Fort. Lee, NJ 07024-5004~~
~~Telephone: (201) 242-6560~~
~~Fax: (201) 242-6561~~



EPA Reg. No. 8622-68

EPA Est. No. 57091-IS-001

IN CASE OF EMERGENCY CONTACT:

CHEMTREC - (800) 424-9300 or (703) 527-3887 (Collect calls accepted) [24 Hr. Number]

or

Ameribrom / Clearon Emergency Response Number - (304) 746-3000 [24 Hr. Number]

or

Ameribrom West - (805) 988-9719 [Business Hours ONLY]

5419 Santa Clara Ave.

Camarillo, CA 93010

NET CONTENTS: _____ gallons (_____ liters)

AB120-XX-Rev.0

05-05-2004

00118
3/15

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Applicators and other handlers must wear long-sleeved shirt, long pants, and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining personal protective equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d) (4-6)], the handler PPE requirements reduced or modified as specified in the WPS.

USER SAFETY RECOMMENDATIONS

Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.

Users should remove PPE clothing immediately if pesticide gets inside.

Then wash thoroughly and put on clean clothing.

Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

For terrestrial uses; do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe, consult the State or Tribal agency responsible for pesticide regulation.

Read entire label before using this product.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers shall be in the area during application. For any requirements specific to your State and Tribe, consult the State/Tribal agency responsible for pesticide regulation.

70418
4/15

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170.

This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), notification to workers, and restricted-entry interval. The requirements in this box apply to uses of this product that are covered by the Workers Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 4 hours.

PPE required for early entry into treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:

- coveralls over short-sleeved shirt and short pants,
- socks and chemical resistant footwear, and
- waterproof gloves.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

Do not enter treated areas until sprays have dried.

GENERAL APPLICATION INSTRUCTIONS

CALIRUS 150 is applied by various application methods, including foliar sprays, soil drench, application into planting holes, seed and bulb dip, seed coating and pelleting, and bare root dip. For foliar sprays, apply CALIRUS 150 with sufficient water for adequate coverage of foliage, according to crop and growth stage. To insure good coverage, spray to wet.

MIXING INSTRUCTIONS

CALIRUS 150 dilution must be made immediately before application.

1. Fill the tank with 1/2 of the required volume of water.
2. Add CALIRUS 150 slowly to the tank and agitate by hydraulic or mechanical means
3. Continue to fill the tank with water to the desired volume while agitating.
4. Continue agitation when applying.

80918
5/15

APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

Apply this product through microjet, drip, solid set and center pivot sprinkler irrigation systems. Do not apply this product through any other type of irrigation system.

PREPARATION OF INJECTION EQUIPMENT: Remove scale, pesticide residues and other foreign matter from the chemical tank and entire injector system. Flush with clean water.

APPLICATION INSTRUCTIONS: Fill tank with 1/2 to 3/4 of desired amount of water. Start mechanical or hydraulic agitation. Add the required amount of CALIRUS 150, then the remaining volume of water. Set sprinkler to deliver 0.1 to 0.3 inch of water per acre. Start sprinkler and uniformly inject the solution of CALIRUS 150 into the irrigation water line so as to deliver the desired rate per acre. Inject the CALIRUS 150 solution with a positive displacement pump into the main line ahead of a right angle turn to insure adequate mixing. For other questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

NOTE: For Microjet and Drip Irrigation Systems: Avoid further irrigation of the treated area for 24 to 48 hours after treatment with CALIRUS 150 has been completed. For Solid Set and Corner Pivot Irrigation Systems: Avoid further irrigation of the CALIRUS 150 treated area until after the foliage has completely dried. This will help to prevent washing the chemical off the crop.

GENERAL PRECAUTIONS FOR APPLICATIONS THROUGH SPRINKLER IRRIGATION SYSTEMS

The irrigation system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from back flow. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump. The pesticide injection pipeline must also contain a functional, normally closed solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected. Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Maintain continuous agitation in mix tank during mixing and application to assure uniformity of solution. Do not apply when wind speed favors drift beyond the area intended for treatment, when system connection or fittings leak, when nozzles do not provide uniform distribution, or when lines containing the product must be dismantled or drained. Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of chemigation water.

90F18
6/15

CALIRUS 150 revised two sections to EPA 05 June 1

Allow sufficient time for pesticide to be flushed through all lines and all nozzles before turning off irrigation water. A person knowledgeable of the chemigation system and responsible for its operation shall shut down and adjust the system as needed.

Do not connect an irrigation system (including greenhouse system) used for pesticide application to a public water system unless the label-prescribed safety devices for public water supplies are in place.

COMPATIBILITY

CALIRUS 150 is compatible with most products used in agriculture; however, crop sensitivity to these mixtures can vary. In particular, mixtures of CALIRUS 150 with some foliar fertilizers and copper products are incompatible and phytotoxicity to certain plants can result. To avoid undesirable phytotoxicity, do not apply or mix CALIRUS 150 and copper based compounds or fertilizers.

CALIRUS 150 use with spray adjuvants (i.e. stickers, spreaders, wetting agents) is limited to those adjuvants that have been tested before use and shown to be compatible with this product. Do not use acidifying type compatibility agents.

To determine the compatibility of CALIRUS 150 with other products, use a jar compatibility test. Add the correct proportions of each product and the appropriate quantity of water to clean container, thoroughly mix, then let stand for 3-5 minutes. If the mixture remains in solution or if the products remixed readily, the products are considered compatible. To determine if a combination is phytotoxic to a specific crop, spray a few plants/trees and evaluate those plants/trees 3- 7 days later for visual symptoms.

100 of 19
7/15

GENERAL INSTRUCTIONS FOR APPLICATIONS

Food Crops

CALIRUS 150 fungicide is used to control *Pythium*, *Plasmodiophora*, Downy Mildews, *Alternaria* and *Phytophthora* diseases on agronomic crops.

Apply CALIRUS 150 to food crop plants including:

Cereals: Millet, Oat, Rye, Sorghum, Corn, Wheat, Rice.

Field crops: Alfalfa, Clover, Cotton, Peanuts, Sugar Cane, Sunflower, Tobacco, Vetch.

Legumes: Dry beans, Fresh beans, Peas, Soybeans.

Roots & tubers: Beet, Carrot, Potato, Sugar Beets.

Bulb Vegetables: Garlic, Onion.

Fruit and Nut Trees:

Stone fruits: Apricot, Cherry, Nectarine, Peach, Plum, Prune.

Nut trees: Almond, Chestnut, Macadamia, Walnut.

Pome Fruit: Apple, Pear.

Tropical/Subtropical Trees (except Citrus): Avocado, Cinchona, Cocoa, Coconut, Coffee, Mango, Olive, Papaya

Citrus: Citrus spp. and Fortunella spp. including calamondin, citrus citron, citrus hybrids, grapefruit, kumquat, lemon, lime, mandarin (tangerine), sour orange, sweet orange, pomelo, Satsuma mandarin.

Small Fruits and Berries: Blueberry, Cranberry, Raspberry, Strawberry, Grape.

Fruiting Vegetables (except cucurbits): Chilies, Peppers, Tomato, Eggplant

Cucurbits: Chayote, Chinese waxgourd, citron melon, cucumber, gherkin, edible gourd, Momordica spp., muskmelon, pumpkin, summer squash, winter squash, watermelon.

Non-Brassica Leafy Greens: Celery, Lettuce (head and leaf), Rhubarb, Spinach

Brassica (Cole crops) Vegetables: broccoli, Chinese broccoli, broccoli raab, Brussels sprouts, cabbage, Chinese cabbage (bok choy, napa, gai choy), cauliflower, cavalo broccoli, collards, kale, kohlrabi, mizuna, mustard greens, mustard spinach, rape greens.

Herbs: angelica, balm, basil, borage, burnet, chamomile, catnip, chervil, chive, Chinese chive, Cinnamon, clary, coriander (leaf), costmary, cilantro (leaf), curry (leaf), dillweed, horehound, hyssop, lavender, lemongrass, lovage (leaf), marigold, marjoram (*Origanum* spp.), Mint, nasturtium, nutmeg, parsley, pennyroyal, rosemary, rue, sage, summer and winter savory, sweet bay, tansy, tarragon, thyme, wintergreen, woodruff and wormwood.

Miscellaneous: Artichoke, Asparagus, Hops, Kiwi, Pineapple ,

Apply product prior to disease development and/or upon disease appearance, in conjunction with good agricultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the indicated application rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed.

Disease	Crop	Application method	Rate	Application program
<i>Pythium</i> Damping-Off and Root Rot	All crops except Cucurbits and Lettuce	Soil drench	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank. Apply 2.5 to 5 gal. (10 - 20 L) CALIRUS 150 per acre	Apply diluted CALIRUS 150 with normal irrigation systems (such as drip irrigation system). Apply shortly before, or after sowing/planting, and at 2-week intervals within 2 months from time of planting/sowing.
<i>Pythium</i> Damping-Off and Root Rot	Cucurbits, Lettuce	Soil drench	Apply 1 gal. CALIRUS 150 per acre. Dilute in a volume of 30-100 gal. of water per acre	Apply diluted CALIRUS 150 with normal irrigation systems, shortly before, or after sowing/planting. Apply at 1-2-week intervals within 2 months from time of planting

11 of 18
8/15

CALIRUS 150 revised two sections to EPA 05 June 1

<i>Phytophthora</i> Root Rot of annuals	All crops	Soil drench	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank. Apply 2.5 to 5 gal. (10 - 20 L) CALIRUS 150 per acre	Apply diluted CALIRUS 150 with normal irrigation systems (such as drip irrigation system) upon appearance of disease
<i>Phytophthora</i> Root Rot of perennials	Fruit and nut trees except Citrus	Application into planting holes	Dilute CALIRUS 150 1:15 (e.g., add 1 gal. to 14 gal.; add 1 L to 14 L) in water in fertilizer tank. Apply about 7 fl. oz. (200 ml) CALIRUS 150 /planting hole	Apply diluted CALIRUS 150 to planting hole of existing mature tree upon appearance of disease. Repeat as needed.
<i>Phytophthora</i> Root Rot	Citrus	Application into planting holes	Soak the ground around the tree with water and immediately apply 7 fl. oz. CALIRUS 150 per tree into planting hole.	Apply into planting hole of existing mature tree upon appearance of disease. Repeat as needed.
<i>Plasmodiophora</i> <i>Brassica</i> (Clubroot)	Brassica	Soil drench	Apply 0.75%-1.5% CALIRUS 150. (e.g., dilute 0.75-1.5 gal. CALIRUS 150 in 100 gal. of water.) Use a volume of 30-100 gal. of water per acre	Apply diluted CALIRUS 150 with normal irrigation systems, shortly before, or after sowing/planting. Apply at 1-2-week intervals within 3 months from time of sowing/planting. Use the higher rate when disease severity is high.
<i>Pythium</i> Damping-Off and <i>Phytophthora</i> Root Rot	All crops	Seed coating and pelleting	Use 10 fl. oz (300 ml) of CALIRUS 150 for 2.2 lb (1 kg) seeds	Apply undiluted CALIRUS 150 to conventional equipment for seed coating and pelleting
<i>Pythium</i> Damping -Off	All crops	Seed and Bulb dip	Dip the desired amount of seeds or bulbs into undiluted CALIRUS 150 solution for 10 - 20 minutes	Apply immediately before sowing/planting. Allow complete air-drying of the seeds or bulbs before sowing. Shortly after sowing/planting, apply with normal irrigation systems at 1-2-week intervals within 2 months from time of planting
<i>Phytophthora</i> Late Blight	All crops except potatoes	Foliar spray	Dilute CALIRUS 150 1:27 (e.g., add 3 qt. to 80 qt. ; 3 L to 80 L) in water in sprayer tank. Apply about 6 qt. (6 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 to foliage with conventional sprayers at 2-3 week intervals, and repeat as needed
<i>Phytophthora</i> Late Blight	Potatoes	Foliar spray	Apply 2.5 to 6.0 gal CALIRUS 150/acre. Dilute in a volume of 30-100 gal water/acre	Upon disease appearance apply diluted CALIRUS 150 to foliage with conventional sprayers at 7-10 day intervals, and repeat as needed. Use the higher rate when disease is severe.
Downy Mildew	All crops except Cucurbits, <i>Brassica</i> , Lettuce, Hops, Coriander, Celery, Parsley, Grapes	Foliar spray	Dilute CALIRUS 150 1:27 (e.g., add 3 qt. to 80 qt. ; 3 L to 80 L) in water in sprayer tank. Apply about 6 qt. (6 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 to foliage with conventional sprayers at 2-3 week intervals, and repeat as needed
Downy Mildew	Cucurbits, <i>Brassica</i> , Lettuce, Hops, Coriander, Celery, Parsley	Foliar spray	Apply 1 gal. CALIRUS 150 per acre. Dilute in a volume of 30-100 gal. of water per acre	Apply diluted CALIRUS 150 to foliage with conventional sprayers at 2-3 week intervals, and repeat as needed

120118
9/15

CALIRUS 150 revised two sections to EPA 05 June 1

Downy Mildew	Grapes	Foliar spray	Apply 1.0% CALIRUS 150. Dilute 1.0 gal. CALIRUS 150 in 100 gal. of water. Use a volume of 80-100 gal. water per acre.	Spray up to a full coverage of foliage (leaves and fruits) at 2-3 week intervals. On Spring disease appearance, start spray at stage of Sprouting to Set-Up. On Autumn disease appearance, start spray upon disease appearance on leaves.
<i>Alternaria</i> spp.	Citrus	Foliar spray	Apply 1.0% CALIRUS 150. Dilute 1.0 gal. CALIRUS 150 in 100 gal. of water.	Spray up to a full coverage of foliage (leaves and fruits). Start spraying upon appearance of disease (Spring and Autumn appearances).
<i>Phytophthora</i> Brown Rot	Citrus	Foliar spray	Apply 1.0% CALIRUS 150. Dilute 1.0 gal. CALIRUS 150 in 100 gal. of water. Use a volume of 100 gal. of water per acre	Spray the lower parts of the trees (about 3-5 feet high above the ground) with diluted CALIRUS 150.
<i>Pythium</i> and <i>Phytophthora</i> Root Rot	All crops	Foliar spray	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. To 149 qt.; Add 1 L to 149 L) in water in sprayer tank. Apply 2.5 to 5 gal. (10 - 20 L) CALIRUS 150 per acre	Apply diluted CALIRUS 150 with sprayer on day of sowing/planting, and at 2-week intervals within 2 months from time of planting/sowing.

13018
19/15

Ornamentals

CALIRUS 150 fungicide is used for the control of *Pythium* Damping-Off, *Pythium* Root Rot, *Phytophthora* Root Rot, Downy Mildew and *Phytophthora* Blight diseases of ornamentals grown in commercial nurseries, greenhouses, parks and commercial landscapes.

Apply CALIRUS 150 to plants including: Aglaonema, Aphelandra, Arborvitae, Azaleas, Bougainvilleas, Boxwood, Cattleya skinneri, Ceznothus, Cotoneaster, Cissus, Dieffenbachia, Dogwood, Ficus Hibiscus, Japanese Holly, Juniper, Leather-leaf Fern, Monterey Pine, Peperomia, Photinia, Pittosporum, Philodendron, Pieris, Pothos, Rhododendron, Roses, Schefflera, Sedum, Sempervivum, Spathiphyllum, Taxus media, White Cedar, White Pine, Zygocactus.

Apply product prior to disease development and/or upon disease appearance, in conjunction with good agricultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the indicated application rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed.

Disease	Application method	Rate	Application program
<i>Pythium</i> Damping – Off and Root Rot	Soil drench	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank. Apply 2.5 to 5 gal. (10 -20 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 with normal irrigation systems (such as drip irrigation system). Apply shortly before, or after sowing/planting, and at 2-week intervals within 2 months from time of planting/sowing.
<i>Phytophthora</i> Root Rot of annuals	Soil drench	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank. Apply 2.5 to 5 gal. (10 -20 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 with normal irrigation systems (such as drip irrigation system) upon appearance of disease.
<i>Phytophthora</i> Root Rot of perennials	Application into planting holes	Dilute CALIRUS 150 1:15 (e.g., add 1 gal. to 14 gal; add 1 L to 14 L) in water in fertilizer tank. Apply about 7 fl. oz (200 ml) CALIRUS 150 per planting hole.	Apply diluted CALIRUS 150 to planting hole of existing mature tree upon appearance of disease. Repeat as needed.
<i>Pythium</i> Damping -Off	Seed and bulb dip	Dip the desired amount of seeds and bulbs into product for 10 – 20 minutes.	Apply immediately before sowing/planting.
<i>Pythium</i> and <i>Phytophthora</i> Root Rot	Spray	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; Add 1 L to 149 L) water in sprayer tank. Apply 2.5 to 5 gal. (10 -20 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 with sprayer on day of sowing/ planting, and at 2-week intervals within 2 months from time of planting/ sowing.
<i>Pythium</i> Damping –Off	Seed coating and pelleting	Use 10 fl. oz. (300 ml) of CALIRUS 150 for 2.2 lb. (1 kg) seeds.	Apply undiluted CALIRUS 150 to conventional equipment for seed coating and pelleting.
Downy Mildew and <i>Phytophthora</i> Blight	Foliar	Dilute CALIRUS 150 1:27 (add (e.g., add 3 qt. to 80 qt 380 L) in water in sprayer tank. Apply about 6 qt. (6 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 to foliage with conventional sprayers at 2-3 week intervals, and repeat as needed.
<i>Pythium</i> and <i>Phytophthora</i> Root Rot	Bare Root Dip	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) water in fertilizer tank	Dip transplants into diluted CALIRUS 150 to thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.

140 of 18
11/15

Bedding Plants

CALIRUS 150 fungicide is used for the control of *Pythium* Damping-Off, *Pythium* Root Rot, *Phytophthora* Root Rot, Downy Mildew and *Phytophthora* Blight diseases of bedding plants grown in commercial nurseries, greenhouses, parks and commercial landscapes.

Apply CALIRUS 150 to plants including: Ageratum, Algerian Ivy, Anthurium, Artemisia, Aster, Baby's Breath, Begonia, Caladium, Carnation, Chrysanthemum, Coleus, Columbine, Daisy Delphinium, Easter Lily, English Ivy, Foxglove, Gaillardia, Geranium, Gloxinia, Impatiens, Marigold, Pansy, Petunia, Phlox, Pinks, Poinsettia, Primrose, Prostrate Rosemary, Salvia, Snapdragon, Verbena, Vinca, Zinnia, etc.

Apply product prior to disease development and/or upon disease appearance, in conjunction with good agricultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the indicated application rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed.

Disease	Application method	Rate	Application program
<i>Pythium</i> Damping -Off and Root Rot	Soil drench	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank. Apply 2.5 to 5 gal. (10 -20 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 with normal irrigation systems (such as drip irrigation system). Apply shortly before, or after sowing/planting, and at 2-week intervals within 2 months from time of planting/sowing.
<i>Phytophthora</i> Root Rot	Soil drench	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank. Apply 2.5 to 5 gal. (10 -20 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 with normal irrigation systems (such as drip irrigation system) upon appearance of disease.
<i>Pythium</i> Damping -Off	Seed and bulb dip	Dip the desired amount of seeds and bulbs into product for 10 - 20 minutes.	Apply immediately before sowing/planting.
<i>Pythium</i> and <i>Phytophthora</i> Root Rot	Spray	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in sprayer tank. Apply 2.5 to 5 gal. (10 -20 liter CALIRUS 150 per acre.	Apply diluted CALIRUS 150 with sprayer on day of sowing/ planting, and at 2-week intervals within 2 months from time of planting/ sowing.
<i>Pythium</i> Damping -Off	Seed coating and pelleting	Use 10 fl. oz. (300 ml) of CALIRUS 150 for 2.2 lb. (1 kg) seeds.	Apply undiluted CALIRUS 150 to conventional equipment for seed coating and pelleting.
Downy Mildew and <i>Phytophthora</i> Blight	Foliar	Dilute CALIRUS 150 1:27 (e.g., add 3 qt. to 80 qt. .; add 3 L to 80 L) in water in sprayer tank. Apply about 6 quarts (6 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 to foliage with conventional sprayers at 2-3 week intervals, and repeat as needed.
<i>Pythium</i> and <i>Phytophthora</i> Root Rot	Bare Root Dip	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank	Dip transplants into diluted CALIRUS 150 to thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.

150418
12/15

CALIRUS 150 revised two sections to EPA 05 June 1

Potting Food and Non-Food Plants

CALIRUS 150 fungicide is used for the control of *Pythium*, *Plasmodiophora Brassica*, Downy Mildew and *Phytophthora* diseases of plants grown in pots or containers in commercial nurseries, greenhouses, parks and indoors.

Apply CALIRUS 150 to plants including: Aglaonema, Aphelandra, Arborvitae, Azaleas, Bougainvilleas, Boxwood, Cattelya skinneri, Ceanothus, Cotoneaster, Cissus, Dieffenbachia, Dogwood, Ficus Hibiscus, Japanese Holly, Juniper, Leather-leaf Fern, Monterey Pine, Peperomia, Photinia, Pittosporum, Philodendron, Pieris, Pothos, Rhododendron, Roses, Schefflera, Sedum, Sempervivum, Spathiphyllum, Taxus media, White Cedar, White Pine, Zygocactus, Cucurbits, Brassica (Cole) Leafy Vegetables, Onion, Lettuce, Celery, Coriander, Parsley, Begonia and Petunia.

Apply product prior to disease development and/or upon disease appearance, in conjunction with good agricultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the indicated application rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed.

Disease	Crop	Application method	Rate	Application program
<i>Pythium</i> Damping-Off and Root Rot	All crops except Cucurbits, Brassica, Onion, Lettuce, Coriander, Celery, Parsley, Begonia and Petunia	Soil drench	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank. Apply 2.5 to 5 gal. (10 -20 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 with normal irrigation systems (such as drip irrigation system). Apply shortly before, or after sowing/planting, and at 2-week intervals within 2 months from time of planting/sowing.
<i>Pythium</i> Damping-Off and Root Rot	Cucurbits, Brassica, Onion, Lettuce, Coriander, Celery, Parsley, Begonia and Petunia	Soil drench	Apply 1.0% CALIRUS 150. Dilute 1.0 gal. CALIRUS 150 in 100 gal. of water. Use a volume of 30-100 gal. of water per acre. Apply up to a full drainage (field capacity) of the container media.	Apply diluted CALIRUS 150 with normal irrigation systems used in nurseries. Apply shortly before, or after sowing. Apply at 1-2-week intervals within 2 months from time of sowing.
<i>Phytophthora</i> Root Rot	All crops	Soil drench	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank. Apply 2.5 to 5 gal. (10 -20 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 with normal irrigation systems (such as drip irrigation system) upon appearance of disease.
<i>Plasmodiophora</i> Brassica (Clubroot)	Brassica	Soil drench	Apply 0.75%-1.5% CALIRUS 150. Dilute 0.75-1.5 gal. CALIRUS 150 in 100 gal. of water. Use a volume of 30-100 gal. of water per acre. Apply up to a full drainage (field capacity) of the container media.	Apply diluted CALIRUS 150 with normal irrigation systems in the nurseries. Apply shortly before, or after sowing, and at 2-week intervals within 2 months from time of planting. Use the higher rate when disease severity is high.
<i>Pythium</i> Damping-Off	All crops	Seed and Bulb dip	Dip the desired amount of seeds or bulbs into undiluted CALIRUS 150 solution for 10 - 20 minutes.	Apply immediately before sowing/planting. Allow complete air-drying of the seeds or bulbs before sowing. After sowing, use the application program for the after sowing period, as indicated above.

16 of 18
13/15

CALIRUS 150 revised two sections to EPA 05 June 1

<i>Pythium</i> Damping -Off	All crops	Seed coating and pelleting	Use 10 fl. oz. (300 ml) of CALIRUS 150 for 2.2 lb. (1 kg) seeds.	Apply undiluted CALIRUS 150 to conventional equipment for seed coating and pelleting.
<i>Pythium</i> and <i>Phytophthora</i> Root Rot	All crops	Bare Root Dip	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank.	Dip transplants into diluted CALIRUS 150 to thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.
<i>Phytophthora</i> Late Blight	All crops	Foliar spray	Dilute CALIRUS 150 1:27 (e.g., add 3 qt. to 80 qt.; 3 L to 80 L) in water in sprayer tank. Apply about 6 qt. (6 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 to foliage with conventional sprayers at 2-3 week intervals, and repeat as needed
Downy Mildew	All crops except Cucurbits, <i>Brassica</i> , Lettuce, Coriander, Celery, Parsley	Foliar spray	Dilute CALIRUS 150 1:27 (e.g., add 3 qt. to 80 qt.; 3 L to 80 L) in water in sprayer tank. Apply about 6 qt. (6 L) CALIRUS 150 per acre.	Apply diluted CALIRUS 150 to foliage with conventional sprayers at 2-3 week intervals, and repeat as needed
Downy Mildew	Cucurbits, <i>Brassica</i> , Lettuce, Coriander, Celery, Parsley	Foliar spray	Apply 1.0% CALIRUS 150. Dilute 1.0 gal. CALIRUS 150 in 100 gal. of water. Use a volume of 30-100 gal. of water per acre.	Apply diluted CALIRUS 150 to foliage with conventional sprayers at 2-3 week intervals, and repeat as needed.
<i>Pythium</i> and <i>Phytophthora</i> Root Rot	All crops	Foliar spray	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in sprayer tank. Apply 2.5 to 5 gal. (10 -20 L) CALIRUS 150 per acre	Apply diluted CALIRUS 150 with sprayer on day of sowing/ planting, and at 2-week intervals within 2 months from time of planting/ sowing

17 of 18
14/15

Conifers and forest trees

CALIRUS 150 fungicide is used for the control of *Pythium* and *Phytophthora* Root Rot of conifers and forest trees including (but not limited to) Douglas Fir, spruce and pines.

Apply product prior to disease development and/or upon disease appearance, in conjunction with good agricultural management practices. Use the higher rate when disease pressure is severe. Do not exceed the indicated application rates or apply more frequently than the specified interval or plant injury can occur. Do not apply to plants that are dormant or heat or moisture stressed.

Disease	Application method	Rate	Application program
<i>Phytophthora</i> Root Rot	Application to roots	Dilute CALIRUS 150 1:15 (e.g., add 1 gal. to 14 gal.; add 1 L to 14 L) in water in fertilizer tank. Apply about 7 fl. oz. (200 ml) CALIRUS 150/tree	Apply diluted CALIRUS 150 to root system of existing mature tree upon appearance of disease. Repeat as needed.
<i>Phytophthora</i> Root Rot	Spray	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in sprayer tank. Apply 10 -20 liter CALIRUS 150 / acre	Apply diluted CALIRUS 150 with sprayer at 2-4 week intervals. Repeat as needed.
<i>Phytophthora</i> Root Rot	Bare Root Dip	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in fertilizer tank.	Dip transplants into diluted CALIRUS 150 to thoroughly wet plant root mass immediately before transplanting. Keep roots submerged for 1 to 2 minutes.

Turf

CALIRUS 150 fungicide is used for the control of *Pythium* Blight and Damping-Off diseases of turf grasses on sod farms, commercial turf production, golf courses, parks and commercial landscapes. Begin preventive applications when conditions first favor disease and repeat at indicated intervals. Use higher rates under severe disease conditions.

Disease	Application method	Rate	Application program
<i>Pythium</i> Blight (Yellow Turf)	Spray	Dilute CALIRUS 150 1:150 (e.g., add 1 qt. to 149 qt.; add 1 L to 149 L) in water in sprayer tank. Apply 2.5 to 5 gal. (10 -20 L) CALIRUS 150per acre	Apply diluted CALIRUS 150 on thoroughly wet foliage at 2-week intervals as long as needed. Do not mow or irrigate treated areas until spray has completely dried.

180418
15/15

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Keep this product in containers stored upright and secured with the original closure. Do not store this product near heat sources. Do not store near strong oxidants. If transfer to another container becomes necessary, ensure that the container is clearly labeled, is a type suitable for the product, and is clean and free of other materials.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal or excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Do not reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, keep out of smoke.

Warranty: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with label directions under normal conditions of use, but neither this warranty nor any other warranty of **MERCHANTABILITY** or **FITNESS FOR A PARTICULAR PURPOSE**, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.