



DANGER POISON PRECAUTIONS



CAN KILL YOU IF SWALLOWED

This product can kill you if swallowed even in small amounts; spray mist or dust may be fatal if swallowed.



CAN KILL YOU BY SKIN CONTACT

This product can kill you if touched by hands or spilled or splashed on skin, in eyes or on clothing (liquid goes through clothes).



CAN KILL YOU IF INHALED

This product can kill you if vapors, spray mist or dust are breathed.

Physicians' Note: Warning symptoms include weakness, headache, tightness in the chest, blurred vision, nonreactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps.

Treatment: Give atropine, preferably by injection, grains 1-100 three or more tablets at once and every hour as required until pupils dilate. Thirty or more tablets may be required during the first 24 hours. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests. 2-PAM is also antidotal and may be administered in conjunction with atropine.

NOTE: Protect from temperatures below 32° F.

Do Not Get On Skin, On Clothing

Wear clean heavy rubber gloves and clean waterproof or freshly-laundered protective clothing (coveralls, cap, etc.) Clothing must not get wet through. Destroy and replace gloves frequently.

Wash thoroughly with soap and warm water before eating or smoking. Bathe immediately after work and change all clothing. Wash clothing thoroughly with soap and hot water before re-use.

In case of contact, immediately remove contaminated clothing and wash skin thoroughly with soap and water.

Do Not Breathe Vapor Or Spray Mist

Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for methyl parathion protection. Airplane pilots should wear full-face canister-type mask.

If handled indoors, provide mechanical exhaust ventilation.

Remove unprotected persons and domestic animals from operating and adjacent areas where insecticide may drift. Prohibit re-entry until drift and vapor dissipate.

Do Not Store Near Food Or Feed Products

Cover spillage with an absorbent such as a soda ash, lime, clay or sawdust. Sweep up and bury. Wash area thoroughly with a full-strength liquid household chlorine bleach.

This product is toxic to fish, birds, and other wildlife.

Birds feeding on treated areas may be killed. Keep out of any body of water.

Do not apply where runoff is likely to occur.

Do not apply when weather conditions favor drift from areas treated.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.

DECONTAMINATE EMPTY CONTAINERS AS FOLLOWS: Add 1 pint of liquid chlorine bleach, close container and shake for 5 minutes. Allow to stand for 15 minutes. Flush with water and mutilate to prevent reuse.

When decontaminating spills or empty containers, observe personal safety precautions listed above.



Atropine is an antidote. Consult physician for emergency supply.

FIRST AID TREATMENT

Call A Physician At Once In All Cases Of Suspected Methyl Parathion Poisoning.

If symptoms or signs of poisoning (See Physicians' Note) include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets (each 1-100 gr.) at once.

Remove patient immediately from the area where methyl parathion is present. Remove contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of methyl parathion. If swallowed, induce vomiting by giving warm, salty or soapy water. Have patient lie down and keep quiet.

Never Give Anything By Mouth To An Unconscious Person

Insecticide

Active Ingredients:

O,O-dimethyl O-p-nitrophenyl phosphorothioate*

Aromatic Petroleum Solvent

Inert Ingredients

*methyl parathion

(1 gallon contains 4 lbs. methyl parathion)

44.4%

39.5%

16.1%

DIRECTIONS FOR USE

DISCLAIMER

Staple Cotton Services Association A.A.I. warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warranty shall be limited to direct damages, and shall not include consequential, commercial damages such as loss of profit or values, etc.

Staple Cotton Services Association A.A.I. makes no other express or implied warranty including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

BUYER assumes the risk of any use contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable by Staple Cotton Services Association A.A.I.

This product is to be applied by experienced personnel using ground or aircraft equipment. The hazards and precautions on this container are equally applicable to it after dilution with water for spray application. Pour specified amount of this product into nearly filled spray tank. Add balance of water to fill tank. Mix thoroughly either by means of a tank agitator or pump by-pass.

For best results, thoroughly cover all surfaces to be treated with spray. Consult your State Experiment Station or State Agricultural Extension Service for specific recommendations regarding application, dosage and timing of sprays. When applying by ground equipment, use specified dosage in not less than 10 gallons of water per acre, by aircraft equipment in 2.5 gallons of water per acre.

PEACH AND APRICOT: European red mite and green peach aphyd. Apply 1/2 pt. per 100 gallons of water. Apply twice at 5 day intervals for control of European red mites. Do not apply within 14 days of harvest.

VEGETABLE AND FIELD CROPS: Rates are given in terms of pints of METHYL PARATHION 4 E.C. per acre in sufficient water for good coverage. The number shown in parentheses () indicates the number of days which should be allowed between last application of harvest or grazing. Unless otherwise specified, this product will control aphids, armyworms (up to 3rd instar), flea beetles, leafhoppers and mites on the following crops:

DANGER
Keep out of reach
of children

POISON

See side panels
for antidote
and precautions

ALFALFA, CLOVER, VETCH: Spotted alfalfa aphids - 1/2-1 pt. Apply in early spring. Follow state recommendations. For the control of aphids, armyworms (up to 3rd instar), leafhoppers and mites - 2 1/2 pts. (20)

ALFALFA HAY: Alfalfa weevil larvae - Apply 1/4 pt. per acre by ground or air. Apply when buds and shoots first appear ragged or when larvae count first reaches 25 per sweep. (15)

ARTICHOKES, BEETS, CARROTS, CELERY, PEAS, PEPPERS: 1 1/4 pts. (17) artichokes (15) other crops listed except (21) days on beets if tops are to be used for food.]

BEANS green: 2 pts. (21)

BEANS (except green), TOMATOES: 2 pts. (15)

COLE CROPS (Cabbage, Cauliflower, Broccoli, Brussels Sprouts, Collards, Kale and Kohlrabi): Use 2 pts. (21)

COTTON: Aphids, boll weevil, cotton leafworm, fleahopper, lygus, red spider mites, thrips - 1/2-1 pt. Application should be made at 4-5 day intervals until control is obtained. For cotton leaf perforator control apply 1 qt. per acre. Do not apply within 7 days of handpicking or harvest.

For the control of armyworms (up to 3rd instar) on cotton: 2 1/2 pts. Workers entering treated fields within 24 hours of application should wear protective clothing (5).

LETTUCE, SPINACH: 1 1/4 pts. (21)

POTATOES: 2 pts. (5)

SMALL GRAINS (Barley, Oats, Rye and Wheat): Aphids (green bug), black grass bug (Irisia), brown wheat mite (Petrobia), and stink bugs - Apply 1/2-1 pt. For armyworms (up to 3rd instar) and leafhoppers, apply 2 1/2 pts. Use limitations: 1/2 pt. - none, over 1/2 and up to 1 1/2 pts. - 15 days; over 1 1/2 pts. - Do not apply after heads begin to form.

SOYBEANS: Stink bugs, three-cornered alfalfa hopper and velvet bean caterpillar.

Apply 1/4 pt. (20). Do not apply more than two applications per growing season. Do not apply within 20 days of harvest.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

COMBUSTIBLE

Keep Container Closed And Away From Heat Sparks and Open Flame.

Before using, study safety directions. Consult your physician about obtaining an adequate supply of 1-100 gr. atropine tablets for emergency use.

Insecticide

Active Ingredients:

O,O-dimethyl O-p-nitrophenyl
phosphorothioate*

Aromatic Petroleum Solvent

Inert Ingredients

*methyl parathion

(1 gallon contains 4 lbs. methyl parathion)

44.4%

39.5%

16.1%


Keep out of reach
of children

POISON

See side panels
for antidote
and precautions



CAN KILL YOU



DIRECTIONS FOR USE

DISCLAIMER

Staple Cotton Services Association A.A.I. warrant, only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use.

Any damages arising from a breach of this warrant shall be limited to direct damages, and shall not include consequential commercial damages such as loss of profits or values, etc.

Staple Cotton Services Association A.A.I. makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

BUYER assumes the risk of any use contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable by Staple Cotton Services Association A.A.I.

This product is to be applied by experienced personnel using ground or aircraft equipment. The hazards and precautions on this container are equally applicable to it after dilution with water for spray application. Read specified amount of this product into nearly filled spray tank. Add balance of water to fill tank. Mix thoroughly either by means of a tank agitator or pump by-pass.

For best results, thoroughly cover all surfaces to be treated with spray. Consult your State Experiment Station or State Agricultural Extension Service for specific recommendations regarding application, dosage and timing of sprays. When applying by ground equipment, use specified dosage in not less than 10 gallons of water per acre, or more if equipment in 20 gallons of water per acre.

PEACH AND APRICOT: European red mite, and green peach aoid. Appl. 1/2 pt. per 100 gallons of water. Apply twice at 5 day intervals for control of European red mites. Do not apply within 14 days of harvest.

VEGETABLE AND FIELD CROPS: Rates are given in terms of pints of METHYL PARATHION 4 EC per acre in sufficient water for good coverage. The number shown in parentheses () indicates the number of days which should be allowed between last application of harvest or grazing. Unless otherwise specified, this product will control aphids, armyworms (up to 3rd instar), leaf beetles, leafhoppers and mites on the following crops:

ALFALFA, CLOVER, VETCH: Spotted alfalfa aphids -- 1/2-1 pt. Apply in early spring. Follow state recommendations. For the control of aphids, armyworms (up to 3rd instar), leafhoppers and mites -- 2 1/2 pts. (20)

ALFALFA HAY: Alfalfa weevil larvae -- Apply 1/4 pt. per acre by ground or air. Apply when buds and shoots first appear ragged or when larvae count first reaches 25 per sweep. (15)

ARTICHOKES, BEETS, CARROTS, CELERY, PEAS, PEPPERS: 1 1/2 pts. (17) artichokes (15) other crops listed except (21) days on beets if tops are to be used for food.]

BEANS green: 2 pts. (21)

BEANS (except green), **TOMATOES:** 2 pts. (15)

COLE CROPS (Cabbage, Cauliflower, Broccoli, Brussels Sprouts, Collards, Kale and Kohlrabi): Use 2 pts. (21)

COTTON: Aphids, boll weevil, cotton leafworm, fleahopper, lygus, red spider mites, thrips -- 1/2-1 pt. Application should be made at 4-5 day intervals until control is obtained. For cotton leaf perforator control apply 1 qt. per acre. Do not apply within 7 days of handpicking or harvest.

For the control of armyworms (up to 3rd instar) on cotton 2 1/2 pts. Workers entering treated fields within 24 hours of application should wear protective clothing (5).

LETTUCE, SPINACH: 1 1/2 pts. (21)

POTATOES: 2 pts. (5)

SMALL GRAINS (Barley, Oats, Rye and Wheat): Aphids (green bug), black grass bug (tribia), brown wheat mite (Petrobia), and stink bugs. Apply 1/2-1 pt. For armyworms (up to 3rd instar) and leafhoppers, apply 2 1/2 pts. Use limitations: 1/2 pt. - none; over 1/2 and up to 1 1/2 pts. 15 days; over 1 1/2 pts. Do not apply after heads begin to form.

SOYBEANS: Stink bugs, three-cornered alfalfa hopper and velvet bean caterpillar -- Apply 1/2-1 pt. (20). Do not apply more than two applications per growing season. Do not apply within 20 days of harvest.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

COMBUSTIBLE

Keep Containers Closed and Away From Heat, Sparks and Open Flame

Before using, study safety directions. Consult your physician about obtaining an adequate supply of 1:100 gr. atropine tablets for emergency use.

CAN KILL YOU



WARNING
KEEP AWAY from FEED or FOOD PRODUCTS

POISON

CAUTION--DO NOT DROP

DON'T

BREATHE FUMES
TOUCH CONTENTS
SWALLOW

KEEP THIS PRODUCT OUT OF THE REACH OF CHILDREN. Containers of this product are highly toxic and are packed in such a manner as to prevent leakage. If leakage occurs, proper disposal should be made according to the Federal Hazardous Waste Management Act, as amended by the Interstate Commerce Commission.

For more information, contact your local health department or the nearest Federal Bureau of Investigation office.

SOLD BY:
STAPLE COTTON SERVICES ASSOCIATION A.A.I.
FARM CHEMICALS DIVISION
GREENWOOD, MISSISSIPPI

E.P.A. Reg. No. 8648-12

Made and Printed in U.S.A.