

TENOX[®] P

GRAIN PRESERVATIVE

ACCEPTED
 JUN 29 1978
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 8655-5

Active Ingredient Propionic acid 99.5%
 Inert Ingredients 0.5%



Marketed by
EASTMAN CHEMICAL PRODUCTS, INC., Kingsport, Tennessee 37629

KEEP OUT OF REACH OF CHILDREN

Made in U.S.A.

DANGER! CORROSIVE, CAUSES EYE AND SKIN DAMAGE
 COMBUSTIBLE DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

Do not get in eyes, on skin, or clothing. Rubber gloves, rubber aprons, and rubber footwear and goggles or face shield should be worn. Avoid breathing vapor; use approved respirator if working near mist or high concentration of vapors. Keep container closed. Use with adequate ventilation. Wash thoroughly after handling.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes while removing clothing and shoes. Call a physician. Wash clothing before reuse.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Bin Preparation As with other blends of acids, Tenox P is corrosive. Bins should be protected by lining with non-toxic plastic sheeting before filling.

Application Tenox P is suggested for application as directed below immediately after harvest to prevent fungal growth in stored corn, milo, barley, wheat, and oats:

Pounds per ton of grain with the following respective percent moisture content

Moisture Content (%)	16	18	20	22	24	26	28	30	32	34	36
Pounds of Tenox P	11	13	15	17	19	21	23	25	27	29	31

[Convert to "fluid oz. per 100 lbs. of grain" by use of the formula: (lbs. product per ton grain) × 0.77 = (fl. oz. product per cwt. grain)]

Apply undiluted, by use of low pressure spray nozzles achieving uniform coverage, as grain passes through applicator. Stop grain flow immediately in event of spray failure or nozzle plugging. Concentrations of untreated grain may develop fungal growth which may spread and spoil all grain in the bin. ONLY GRAIN WHICH IS TO BE USED AS ANIMAL FEED SHOULD BE TREATED since application will reduce grains to Sample Grade and reduce germination. Increasing the suggested rate of application in the Table by 25% for the top 10% of the grain placed in each storage bin will provide additional protection against the effect of normal moisture migration in the grain during storage.

— STORAGE AND DISPOSAL —

STORAGE: Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide, spray mixture, or rinsate that cannot be used or chemically processed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies. Open dumping is prohibited.

CONTAINER DISPOSAL: Reseal container and offer for reconditioning, or triple rinse (or equivalent) and offer for recycling, reconditioning, or disposal in approved landfill, or bury in a safe place.

GENERAL: Consult Federal, State or Local disposal authorities for approved alternative procedures.

WARRANTY AND LIMITATION OF LIABILITY

Tenox P Grain Preservative is warranted to conform to the chemical description hereon, and to be reasonably fit for the described use when applied strictly in accordance with directions. THERE ARE NO OTHER WARRANTIES, EXPRESS OR IMPLIED, INCLUDING THOSE OF MERCHANTABILITY OR FITNESS FOR PURPOSE. Factors beyond the control of Eastman Kodak Company and its marketing affiliates prevent them from assuming responsibility for damage to other property, including plants and crops, even when this product is applied according to directions. Buyer and User assume all risks beyond the above warranty. In no event shall the liability of Eastman Kodak Company and its marketing affiliates exceed the purchase price hereof.

Propionic acid 53 200 kg • 440 lb
 LOT NUMBER O.T. SHIPPING NAME GALLONS 96 NET WEIGHT

8655-5

6-29-1978