

preventer crabgrass WITH FERTILIZER

GUARANTEED ANALYSIS 25-3-3

Total Nitrogen (N)	25%
1.2% Ammoniacal Nitrogen	
6.5% Coated Slow Release Urea Nitrogen	
17.3% Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	3%
Soluble Potash (K ₂ O)	3%
Derived from Urea, Sulfur-Coated Urea, Diammonium Phosphate, and Muriate of Potash	

The Coated Slow Release Nitrogen is from Sulfur-Coated Urea. Feeds up to 130 days under normal growing conditions.

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS & DOMESTIC ANIMALS CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, ponds and streams. Do not apply when weather conditions favor runoff or drift from target area.

GENERAL INFORMATION

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, ponds and streams. Do not apply when weather conditions favor runoff or drift from target area.

GENERAL INFORMATION

When used as directed this product will provide plant food for your lawn and will control smooth and hairy crabgrass, green and yellow foxtail, barnyardgrass and goosegrass (silver crabgrass or crowsfoot) by killing plants as seeds germinate. It does not control established weeds.

Use only on established lawns listed below which are tolerant to the herbicide in this product when applied as directed. SEE PRECAUTIONS.

Bluegrass (perennial)
Ryegrass (perennial)
Fescue (except fine leaved)
Bahiagrass

Bermudagrass
Centipedegrass
St. Augustinegrass
Zoysiagrass

DIRECTIONS FOR USE

General Classification:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apply STA-GREEN Crabgrass Preventer to establish lawns in the spring before germination of annual weeds. Be sure the entire package (20 lbs.) is applied evenly over 5000 sq. ft. of lawn area.

Use a good lawn spreader and apply evenly over the turf without skips or misses. A more uniform application can be made by spreading half the required amount of product over the area to be treated and then applying the remaining half at right angles or crosswise.

SPREADER SETTINGS

STA-GREEN, CYCLONE	SCOTTS PRECISION FLOW	SEARS BROADCAST, IMPERIAL ROTO-SPRED MARK IV	SEARS 20", IMPERIAL MARK III
4 1/4	7	4 1/4	9

RESEEDING: Chemicals which effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applications. When reseeding, it is necessary that proper moisture be maintained until grass seedlings are established. To insure satisfactory reseeding results after using this product, the amount of seed used should be increased.

PRECAUTIONS: Do not apply in the spring to turfgrass planted the previous fall. Do not apply to bentgrass or dichondra turf. Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, or Centipedegrass until these turfs are well established. If verticutting is done, do not apply for 2 to 3 weeks. Do not let livestock graze treated areas. Do not feed forage from treated areas to livestock.

GENERAL INFORMATION

When used as directed this product will provide plant food for your lawn and will control smooth and hairy crabgrass, green and yellow foxtail, barnyardgrass and goosegrass (silver crabgrass or crow'sfoot) by killing plants and preventing them from becoming established weeds.

Use only on established lawns which are tolerant to the herbicide in this product when applied as directed. SEE PRECAUTIONS.

Bluegrass (perennial)
Ryegrass (perennial)
Fescue (except fine leaved)
Bahia grass

Bermudagrass
Centipedegrass
St. Augustinegrass
Zoysiagrass

DIRECTIONS FOR USE

General Classification

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Apply STA-GREEN Crabgrass Preventer to establish lawns in the spring before germination of annual weeds. Be sure the entire package (20 lbs.) is applied evenly over 5000 sq. ft. of lawn area.

Use a good lawn spreader and apply evenly over the turf without skips or misses. A more uniform application can be made by spreading half the required amount of product over the area to be treated and then applying the remaining half at right angles or crosswise.

SPREADER SETTINGS

STA-GREEN,
CYCLONE
4 1/4

SCOTTS
PRECISION FLOW
7

SEARS BROADCAST,
IMPERIAL ROTO-SPRED
MARK IV
4 1/4

SEARS 20",
IMPERIAL
MARK III
9

RESEEDING: Chemicals which effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after applications. When reseeding, it is necessary that proper moisture be maintained until grass seedlings are established. To insure satisfactory reseeding results after using this product, the amount of seed used should be increased.

PRECAUTIONS: Do not apply in the spring to turfgrass planted the previous fall. Do not apply to bentgrass or dichondra turf. Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, or Centipedegrass until these turfs are well established. If verticutting is done, do not apply for 2 to 3 weeks. Do not let livestock graze treated areas. Do not feed forage from treated areas to livestock.

STORAGE AND DISPOSAL

STORAGE — Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Keep in a dry place.

DISPOSAL — Product that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.

~~Do not use for any other purpose. Dispose of in an incinerator or landfill approved for pesticides, or bury in a safe place.~~

NOTICE — We make no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and/or handling of this material when such use and/or handling is contrary to label instructions.

NFT WT 20 LBS. (9.07 kg)

RESIDING

PRECAUTION

STORAGE AND DISPOSAL

STORAGE — Do not contaminate water, food or feed by storage, disposal or cleaning of equipment. Open dumping is prohibited. Keep in a dry place.

DISPOSAL — Product that cannot be used should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies

Do not reuse container for any purpose. Dispose of in an incinerator or landfill approved for pesticides or in a safe place.

NOTICE: The seller makes no warranty, expressed or implied, in this product. The product is sold as indicated on the label. Buyer assumes all responsibility and liability for the use of the product. The following is contrary to all of the terms.

NET WT. 20 LBS. (9.07 kg)

STA-GREEN PLANT FOOD COMPANY

SYLACAUGA, ALABAMA 35150

STA-GREEN

**crabgrass
preventer**

WITH FERTILIZER

25-3-3

COVERS 5,000 SQ. FT.

PREVENTICER WITH FERTILIZER

25-3-3

COVERS 5,000 SQ. FT.

ACTIVE INGREDIENT

N-butyl-N-ethyl-a, a, a-Trifluro-2, 6-dinitro-p-toluidine

INERT INGREDIENTS

TOTAL

By Weight

1.15%

98.85%


100.00%

Contains 2.5 lbs. benefin per ton.

Benefin is a product of Elanco Products Company, a division of Eli Lilly and Company.



31



ACTIVE INGREDIENT	By Weight
N-butyl-N-ethyl-a, a, a-Trifluro-2, 6-dinitro-p-toluidine	1.15%
INERT INGREDIENTS	<u>98.85%</u>
TOTAL	100.00%

Contains 23 lbs. benefin per ton

Benefin is a product of Elanco Products Company, a division of Eli Lilly and Company.



30

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED, induce vomiting by giving one tablespoon of salt in a glass of warm water.

IF INHALED, remove to fresh air.

IF ON SKIN, Wash with plenty of soap and water

IF IN EYES, flush with water for 15 minutes.

Get medical attention if irritation persists

(SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS)

31

IF ON SKIN, wash with plenty of soap and water.

IF IN EYES, flush with water for 15 minutes.

Get medical attention if irritation persists

(SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS)

EPA Reg. No. 8600

NET WT. 20 LBS. (9.07 kg)

STA-GREEN PLANT FOOD COMPANY

SYLACAUGA, ALABAMA 35150