

STA-GREEN

COVERS 2,500 SQ. FT.

weed & feed

FOR BAHIA, BERMUDA, CENTIPEDE, FESCUE,
AND ST. AUGUSTINE GRASSES.
CONTROLS GROWING BROADLEAF WEEDS.

20-3-6

weeds as it feeds

CAUTION: Keep out of reach of children.
See back panel for additional cautions.

ACTIVE INGREDIENTS:

*Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid	0.072%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) propionic acid	0.235%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.093%

INERT INGREDIENTS: 99.600%

TOTAL: 100.000%

Used at 22 lbs. on 2500 sq. ft. equivalent to:

*2,4-D Acid	-0.080%	-0.23 lbs. acre or 1.20 lbs. per ton
**MCPP Acid	-0.195%	-0.75 lbs. acre or 3.91 lbs. per ton
***Dicamba Acid	-0.024%	-0.09 lbs. acre or 0.48 lbs. per ton

CONTAINS TRIMEC® HERBICIDE +

*Reg. T.M. PBI — Gordon Corporation + U.S. Patent No. 3,264,189
EPA REG. NO. 8880-7
EPA EST. NO. 8880 ALD1

NET WT. 22 LBS.

STA-GREEN PLANT FOOD COMPANY
SYLACAUGA, ALABAMA

20

weed

GUARANTEED ANAL

Total Nitrogen (N)
 0.0% Nitrate Nitrogen
 1.3% Ammoniacal Nitrogen
 15.2% Other Water Soluble Nitrogen
 3.5% Water Insoluble Organic Nitrogen

Available Phosphoric Acid (P₂O₅)

Soluble Potash* (K₂O)
 Derived from Urea, Ammoniated Superphosphate,
 Ureaform, Sulfate of Potash

*In states requiring, reads "Water Soluble Potash"

Potential Acidity equivalent to 600 pounds of sulfuric acid

DIRECTIONS FOR USE

CONTROLS SUCH WEEDS AS: Florida Smartweed, Clover, Cocksfoot, Dandelion, Creechgrass, Lambsquarter, Mustard, Oxalis, Peppercorn, Purslane, Ragweed, Sida, Spurge, and other broadleaf weeds.

TIME TO APPLY: Apply any time broadleaf weeds are present on St. Augustine Grass in summer when the grass is growing rapidly, because temporary results are obtained when weeds are young and growing slowly, control will be slow and incomplete. Moisture from dew or watering on weeds is necessary for control. Do not water for 24 to 48 hours after application.

HOW MUCH TO APPLY: Use 22 lbs. or more per acre on most other grasses. Over-application will injure the grass.

SUGGESTED SPREADER SETTINGS: May vary with age, model and condition of your spreader by trial on a known area.

Make of Spreader
 Sta-Green, Cyclone or Sears Broadcast Spreader
 Scotts, Central or Sears Calibrated (18' or 24')

in states requiring, reads: Water Soluble Grass
Potential Acidity equivalent to 600 pounds of Calcium Carbonate per Ton.

DIRECTIONS FOR USE

CONTROLS SUCH WEEDS AS: Florida Parsley, Black Medic, Chickweed, Clover, Cocklebur, Dandelion, Creeping Charlie, Dock, Henbit, Knotweed, Lambsquarter, Mustard, Oxalis, Peppergrass, Pigweed, Plantains, Poison Ivy, Purslane, Ragweed, Sida, Spurge, Wild Onion, Wild Lettuce, and most other broadleaf weeds.

TIME TO APPLY: Apply any time broadleaf weeds are present, except to avoid treating St. Augustine Grass in summer months (June through September) when it is growing rapidly, because temporary foliage discoloration may occur. Best results are obtained when weeds are young and growing vigorously. If weeds are growing slowly, control will be slow and more than one application may be needed. Moisture from dew or watering on weed foliage at time of application aids control. Do not water for 24 to 48 hours **after** application.

HOW MUCH TO APPLY: Use 22 lbs. on 2,500 sq. ft. for Bahia, St. Augustine Grass and most other grasses. Over-application may cause temporary discoloration.


SUGGESTED SPREADER SETTINGS

May vary with age, model and condition of spreader and walking speed. Check your spreader by trial on a known area.

Make of Spreader

Spreader Settings

Sta-Green, Cyclone or Sears Broadcast (5 ft. swath)	5
Scotts, Central or Sears Calibrated (18")	4 3/4



NOTICE: The seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and/or handling of this material when such use and/or handling is contrary to label instructions.

KEEP IN A DRY PLACE

NET WT. 22 LBS.

STA GREEN PLANT FOOD COMPANY

1000 S. GULF BLVD. TAMPA, FLA. 33604

Available Phosphoric Acid (P₂O₅) 3.00%
Soluble Potash (K₂O) 1.00%
Derived from Urea Ammoniated Superphosphate
Sulfate of Potash
States requiring reads: Water Soluble Potash
Plant Acidity equivalent: 1000 pounds of material per acre

DIRECTIONS FOR USE

CONTROLS SUCH WEEDS AS: Flaxweed, Green Pigweed, Yellow Pigweed, Red Top, Dandelion, Ragwort, Black Top, Chickweed, Groundnut, Mustard, Xanthoxylon, Peppergrass, Yellow Top, Broadleaf, and other weeds.

TIME TO APPLY: Apply to the soil before or after the first rain. Apply in August or early in September. It is most effective when applied to the soil before or after the first rain. It is most effective when weeds are young and growing. It is most effective when weeds are young and growing. It is most effective when weeds are young and growing.

HOW MUCH TO APPLY: Use 1.5 to 2.0 lbs. per acre. Apply to the soil in the spring or fall. Over-application may cause injury to the soil.

SUGGESTED SPREADER SETTINGS
May vary with age, model and size of spreader. Refer to the instructions for the spreader by which it is to be used.

Make of Spreader	Spreader Settings
Sta-Green Cyclone or Sears Broadcast 18" or larger	
Sears Centratrol or Sears Calibrated 18"	

CAUTION: Keep out of reach of children. Keep away from food and feed. Do not take internally. Avoid inhalation or contact with eyes. Wash after use. Do not use on lawns or ornamentals.

NOTICE: The seller makes no warranty, expressed or implied, concerning the use of this product other than that stated on the label. Buyer assumes all risks of use and is liable for any damage or injury caused by the use of this product contrary to the instructions.

KEEP IN A DRY PLACE

NET WT. 22 LBS.

**STA-GREEN PLANT FOOD COMPANY
SYLACAUGA, ALABAMA**

STA-GREEN

COVERS 2,500 SQ. FT.

weed & feed

FOR BAHIA, BERMUDA, CENTIPEDE, FESCUE,
AND ST. AUGUSTINE GRASSES.
CONTROLS GROWING BROADLEAF WEEDS.

20-3-6

weeds as it feeds

CAUTION: Keep out of reach of children
See back panel for additional cautions.

ACTIVE INGREDIENTS:	
*Dimethylamine Salt of 2,4-dichlorophenoxyacetic acid	0.072%
**Dimethylamine Salt of 2-(2-methyl-4-chlorophenoxy) pro- pionic acid	0.235%
***Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.028%
INERT INGREDIENTS:	99.665%
TOTAL:	100.000%
Used at 22 lbs. on 2500 sq. ft. equivalent to:	
*2,4-D Acid	-0.060% -0.23 lbs. acre or 1.20 lbs. per ton
**MCPP Acid	-0.195% -0.75 lbs. acre or 3.91 lbs. per ton
***Dicamba Acid	-0.024% -0.09 lbs. acre or 0.48 lbs. per ton

CONTAINS TRIMEC® HERBICIDE +
*Reg. T.M. PBI - Gordon Corporation + U.S. Patent No. 3,284,186
EPA REG. NO. 8880-7
EPA EST. NO. 8880 AL01

NET WT. 22 LBS.

STA-GREEN PLANT FOOD COMPANY
SYLACAUGA, ALABAMA

8660-7

20-3-6

Weed & feed

GUARANTEED ANALYSIS

Total Nitrogen
 Nitrate Nitrogen
 Ammonia Nitrogen
 Water Soluble Phosphorus
 Water Soluble Potassium
 Available Phosphorus
 Available Potassium
 Sulfur
 Zinc
 Manganese
 Boron
 Copper
 Iron
 Magnesium
 Calcium

DIRECTIONS FOR USE

CONTROLS SUCH WEEDS AS ...

TIME TO APPLY ...

HOW MUCH TO APPLY ...

SUGGESTED SPREADER SETTINGS ...

Make of Spreader	Spreader Settings

CAUTION: Keep out of reach of children. Keep away from feed and food products. Do not take it orally. Avoid inhalation or contact with eyes, skin or clothing. Wash after use. Do not use on dichonra or lippe ground cover or similar ground covers. Do not burn or reuse empty bag. Destroy when empty. Do not apply in areas underlaid by roots of desirable trees and shrubs. Avoid drift of dust to desirable plants.

Avoid application on garden plants, flowers and shrubs. Do not use on newly seeded lawns. Do not contaminate irrigation ditches or water used for irrigation purposes.

Store in a dry place, away from plants, seeds, fertilizers, insecticides, etc.

NOTICE: The seller makes no warranty, expressed or implied, concerning the use of this product other than that stated on the label. Buyer assumes all risks of use and is to hold this seller harmless for any use not intended by the seller.

KEEP IN A DRY PLACE

NET WT. 22 LBS.

STA-GREEN PLANT FOOD COMPANY
 SYLACAUGA, ALABAMA

3

10