

7/10/97

Replanting: If replanting is necessary in a field previously treated with Tri-Scept, replant only soybeans. Rework soil no deeper than the treated zone. Do not reapply Tri-Scept.

All additional use directions, precautions and limitations applicable to the use of Tri-Scept, apply to the use of this tank-mix.

Rotational Crop Restrictions for Tri-Scept:

Use of Tri-Scept herbicide in accordance with label directions is expected to result in normal growth of rotational crops in most situations; however, various environmental and agronomic factors make it impossible to eliminate all risks associated with the use of Tri-Scept and therefore, rotational crop injury is always possible.

Certain severe restrictions apply to corn, wheat and other rotational crops following an application of Tri-Scept. Refer to the label for Tri-Scept for complete details of these and other restrictions.

1. Do not plant wheat within 4 months following an application of Tri-Scept in soybeans; and, do not plant rice until the spring of the year following an application of Tri-Scept.
2. Do not plant barley, field corn, edible beans, grain sorghum, oats, peanuts or tobacco within 11 months after an application of Tri-Scept.
3. Do not plant sugar beets for 26 months after an application of Tri-Scept.
4. Field corn planting restrictions: For the "Southern Use Area" as defined by the label for Tri-Scept; field corn may be planted as a rotational crop in the spring of the year following an application of Tri-Scept unless extreme drought conditions develop or less than 15 inches of rainfall or irrigation is received from the date of application through October of the year of application. For eastern Oklahoma, Arkansas, the Bootheel of Missouri, Tennessee, North Carolina and states south: field corn may be planted in the spring of the year following an application of Tri-Scept unless extreme drought conditions develop or less than 15 inches of rainfall or irrigation is received within 6 months following the date of application.
5. Do not plant cotton or rotational crops other than those listed above within 18 months following an application of Tri-Scept.
6. Do not apply products containing chlorimuron ethyl (e.g., Classic, Canopy, Gemini, Lorox Plus, Preview, etc.) or imazethapyr (e.g., Pursuit, or Pursuit Plus) the same year as Tri-Scept or injury to following crops may occur.

Refer to the labels for Trellan and Tri-Scept for additional use directions, precautions and limitations before use.

Revisions:

1. Added text box containing required FIFRA and DowElanco standard statements and limitations.
2. Minor editing to correct (one or more may apply): reference to "Trellan" as a DowElanco trademark; grammar, punctuation; or use of w/ case.

*Trademark of DowElanco

T2EAG017 Approved 12/06/90