

PM19

8660-10

ACCEPTED

STA-GREEN

APR 18 1995

20 Pounds Treats 5,000 Sq.Ft.

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under Reg. No. 8660-10

# lawn pest control & fertilizer 25-3-3

**For Control of SURFACE INSECTS such as:**  
Sod Webworms(Lawn Moth), Armyworms, Ants, Chinch Bugs, Brown Dog Ticks, Fleas, Cutworms, Chiggers, Clover Mites, and Crickets and  
**SOIL INSECTS such as:** "White Grub" Larvae of Japanese Beetle, European Chafer, Southern Chafer, Hyperodes Weevil and Billbugs.  
**CONTROLS PESTS AS IT FEEDS GRASS**

ACTIVE INGREDIENT: Chlorpyrifos: [O, O, diethyl O-(3,5,6 trichloro- 2-pyridyl) phosphorothioate] - 0.75%  
INERT INGREDIENTS: - 99.25%  
Total 100.00%

KEEP OUT OF REACH OF CHILDREN

## CAUTION

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Call a physician or Poison Control Center immediately. Have victim drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. **IF INHALED:** Remove victim to fresh air. Apply artificial respiration if indicated. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists.

**NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine by injection only is an antidote.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Harmful if swallowed. May be absorbed through the skin. Avoid inhalation or contact with skin, eyes and clothing. Do not apply by hand. Wash thoroughly with soap and water after using. Wash contaminated clothing before reuse. Avoid contamination of feed or foodstuffs. Keep children and pets off treated areas until this material is washed into the soil and turf is dry.

#### ENVIRONMENTAL HAZARDS

Product is toxic to birds, fish, and aquatic invertebrate. Birds feeding in treated areas may be killed. Clean up spilled product to reduce exposure to wildlife. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. Apply this product only as specified on this label.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not reuse bag. Wrap in several layers of newspaper and discard in trash.

## SUGGESTED SPREADER SETTINGS

Spreader may vary with age, model and condition, and walking speed. Check your spreader by trial on a known area.

20 lbs. to cover	STA-GREEN CYCLONE B-1	SCOTTS PRECISION FLOW	SEARS BROADCAST IMPERIAL ROTO-SPRED MARK IV	SEARS 20" IMPERIAL MARK III
5,000	4 1/4	7	4 1/4	9
3,300	5	8	5	10

## CONTROLS SURFACE INSECTS

**SUCH AS:** Sod Webworms(Lawn Moth), Armyworms, Ants, Chinch Bugs, Brown Dog Ticks, Fleas, Cutworms, Chiggers, Clover Mites and Crickets. **TIME TO APPLY:** Apply anytime pests appear and before severe damage can be done. **HOW MUCH TO APPLY:** Uniformly apply 20 lbs. on 5,000 sq.ft. for Bahia, Bermuda, Centipede, Fescue, St. Augustine grasses and Dichondra. Treat when insects first appear; retreat as needed. To help obtain uniform distribution, apply one-half the required amount of product in one direction; then apply the balance at a right angle. It is not necessary to water in the insecticide for control of surface feeding insects.

## FOR ROOT FEEDING INSECTS

**For ROOT FEEDING INSECTS such as Billbug Larvae, and the "White Grub" larvae of Japanese Beetle, European Chafer, Southern Chafer and Hyperodes Weevil** apply 6 lbs./1,000 sq.ft. of turf, (20 lbs. to treat 3,300 sq.ft.). Make sure your spreader is properly calibrated to apply this product. Irrigate the turf after treatment as soon as possible to wash the insecticide into the root zone where the insects are feeding. Apply when the White Grub larvae are young and actively feeding near the soil surface in mid-July and early August. For the Hyperodes (Turftgrass) Weevil apply in Mid-April and again in Mid-May at the same rate as above. After application do not allow children or pets on the treated areas until insecticide has been washed into the soil and the grass is dry.

## GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	Variable %
Variable % Ammoniacal	
Variable % Coated Slow Release Urea Nitrogen	
Variable % Urea Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	Variable %
Soluble Potash (K <sub>2</sub> O)	Variable %
Primary Plant Food derived from: Urea, Sulfur Coated Urea, Diammonium Phosphate, Muriate of Potash. Potential Acidity xxx lbs. CaCO <sub>3</sub> /ton.	

### IMPORTANT: PLEASE READ BEFORE USE

By using this product, the user accepts the following: **LIMITED WARRANTY:** Pursell warrants that (a) this product conforms to the chemical description on its label; (b) this product is reasonably fit for the purposes stated on its label, subject to the inherent risks referred to herein, when used in accordance with its directions; and (c) that the directions, warnings, cautions, and other statements on this label are based upon responsible experts' evaluations of reasonable tests of effectiveness, of toxicity to laboratory animals and plants, of residues on food crops, and upon reports of field experience. Treating has not been performed on all varieties of turfgrasses, in all states, or under all application, weather and crop conditions. There are no express warranties other than those set forth herein. Pursell neither makes nor intends, nor does it authorize any agent or representative to make, any other warranty, express or implied. Pursell expressly excludes and disclaims all implied warranties of merchantability, fitness for particular purpose, or any other warranty of quality or performance. This warranty does not extend to, and the user shall be solely responsible for, any loss or damage which results from the use of this product in any manner which is inconsistent with this label's directions, warnings or cautions. User's exclusive remedy and Pursell's or seller's exclusive liability for any claim, loss, damage, or injury resulting from the use or handling of this product, whether or not based in contract, negligence, strict liability in tort, or otherwise, shall be limited, at Pursell's option, to replacement of, or repayment of the purchase price for, the quantity of product with respect to which damages are claimed. In no event shall Pursell or Seller be liable for special, indirect, or consequential damages resulting from the use or handling of this product. (d) 8660-10 pm5/01-25-95)

EPA REG. NO. 8660-10 EPA EST. NO. 8660-AL-1  
NET WEIGHT 20 POUNDS (9.07 kg.)

# PURSELL

INCORPORATED 1904

BEST COPY AVAILABLE

1981