

Lawn Pest Control & Fertilizer 29-3-10

Treats 14,500 square feet By Weight **ACTIVE INGREDIENTS** Chlorpyrifos [O,O-Diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 00 50% TOTAL 100 00% **GUARANTEED ANALYSIS** 29-3-10 29% Total Nitrogen (N) ... 1.2% Ammoniacal Nitrogen 27.8% Urea Nitrogen*

 Available Phosphoric Acid (P2O5)
 3%

 Soluble Potash (K2O)
 10%

V.S. PATENT NO. 3,244,599

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by sticking finger down throat.

IF ON SKIN: Wash with soap and water.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation develops or persists

(SEE BELOW FOR ADDITIONAL PRECAUTION STATEMENTS)

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS & DOMESTIC ANIMALS

CAUTION:

Harmful if swallowed. May be absorbed through the skin. Avoid inhalation or contact with skin, eyes and clothing. Do not apply by hand. Wash thouroughly with soap and water after using. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Wash contaminated clothing before reuse. Avoid contamination of feed and foodstuffs. Keep children and pets off treated areas until this material is washed into the soil and turf is dry.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling CONTROLS SUCH PESTS AS: Sod Webworms (Lawn Moth). Armyworms, Ants. Chinch Bugs, Brown Dog Ticks, Fleas, Cutworms, Chiggers, Clover Mites and Crickets

TIME TO APPLY Apply anytime after pests appear and before severe damage can be done. Water in well for best results. Use 50 lbs. on 14,500 square feet for Bahia, Bermuda, Centipede, Fescue, St. Augustine grasses and Dichondra. To help obtain uniform distribution, Apply one-half the required amount of product in one direction; then apply the balance at a right angle. Consult your Cooperative Agricultural Extension Service for proper time to freat in your area.

SUGGESTED SPREADER SETTINGS

Spreader may vary with age, model and ground or PTCL speed. Check your spreader by trial on a known area.

SPREADER	GROUND OR PTC SPEED	COVERGE WIDTH	SETTINGS
Scotts Broadcast	3 M P H	6 ft	r
Cyclone	3 M P H ←	6 ft	4 5
Lely	5 M P H	39 ft	

STORAGE AND DISPOSAL

STORAGE: Do not contaminate water, food or feed by storage, dispusal or cleaning of equipment. Open dumping is prohibited. Keep in a dry place DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection

NOTICE: The seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions

EPA REG. NO.

Net Wt. 50 lbs. (22.68 Kg.) STA-GREEN PLANT FOOD COMPANY SYLACAUGA, ALABAMA 35150

EPA Est. No. 8F60-AL-1