

St. Augustine & Centipede Weed & Feed


Weed Control Plus Lawn Fertilizer

Use on St. Augustine, Centipede, Zoysia and Carpetgrass Lawns Only

28-3-7 Fertilizer

Controls Weeds While Fertilizing Your Lawn

16 lb. Bag Treats 5,000 Sq.Ft.



ACTIVE INGREDIENT:

Atrazine: 2 chloro-4-ethylamino-6-isopropylamino-s-triazine - - - - 1.406%

Related Compounds - - - - - 0.074%

INERT INGREDIENTS: - - - - - 98.520%
Total 100.000%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 8660-18

EPA EST. NO. 8660-AL-1

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Harmful if swallowed. Do not contaminate water used for irrigation or domestic purposes. Do not graze treated areas. Do not feed clippings to livestock.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Not for use on or under ornamental trees, shrubs, bedding plants, or garden plants. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner consistent with its labeling. Not for use on turf being grown for sale or for commercial use as sod, or for commercial seed production, or for research purposes. Apply only when grass is dry.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children and domestic animals. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not reuse empty bag. Securely wrap original container in several layers of newspaper and discard in trash.

WHAT THIS PRODUCT DOES

This product, "St. Augustine & Centipede Weed & Feed" controls and prevents weeds while feeding your lawn.

CONTROLS these common weed:

Annual Bluegrass (poa annua)	Chickweed (Common & Mouseear)	Fragrant Cudweed (Rabbit Tobacco)	Pepperweed
Annual Sowthistle	Common Purslane	Henbit	Purslane
Annual Clover	Crabgrass	Knotweed	Red Sorrel
Black Medic	Cudweed	Lespedeza	Rescue Grass
Blue Toadflax	Cutleaf Evening Primrose	Mat Lippia (Malchweed)	Sandspur
Bull Paspallum	Dandelion	Moneywort	Shepards Pursue
Buckhorn	Dichondra	Narrowleaf Vetch	Spurweed
Plantain	Dollarweed	Oxalis (Common, Yellow, Woodsorrel)	Slinging Nettle
Bur Clover	Florida Betony (Rattlesnake weed)	Parsley-Piert	Virginia Creeper
Carolina			Wild Mustard

IMPORTANT USE PRECAUTIONS

- Do not apply on or under the branch spread (root zone) area of trees, shrubs, bedding plants, flowers, or garden plants.
- Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.
- Do not apply to newly seeded Zoysia, Centipede, and Carpetgrass lawns until after the new grass has been mowed at least twice.
- Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds.
- Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80 F or above, water sufficiently to rinse particles off the grass blades.

TO CONTROL GROWING BROADLEAF WEEDS

Apply any time weeds are actively growing. If complete control is not obtained a second treatment may be required 2 months after the first. Apply in the fall to control Henbit, Chickweed and other winter weeds.

TO PREVENT ANNUAL WEEDS

Apply in January, February or early March to prevent germination of annual weeds such as crabgrass, Bull Paspallum, Sedges (from seed) and Spotted Spurge. Don't apply to Centipede or Carpetgrass lawns during the period of winter dormancy.

SPREADER SETTINGS

Always apply fertilizer with a spreader.

Sta-Green (R)Brdcst.	3.5
Cyclone B1-84	4.2
Central	3.5
Central Drop	6.0
Scotts Speedy Green	3.5
Ortho Rotary	2.5
Ortho Drop	4.0

For best results, water in product after application as fertilizers will not work without water. For a beautiful, healthy lawn, get on a program of lawn care. Apply fertilizer, weed controls, and pest controls on a regular basis.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N)	- - - - -	Variable %
X X% Ammoniacal Nitrogen		
X X% Urea Nitrogen		
Available Phosphoric Acid (P2O5)	- - - - -	Variable %
Soluble Potash (K2O)	- - - - -	Variable %
Sulfur (S) X %	Copper (Cu) X %	Iron (Fe) X %

... and Registration Act
... the pesticide
...
Reg. No. 8660-18

Related Compounds - - - - - 0.014%
INERT INGREDIENTS: - - - - - 98.520%
Total 100.000%

19 32

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
EPA REG. NO. 8660-18

EPA EST. NO. 8660-AL-1

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMAN AND DOMESTIC ANIMALS

CAUTION: Causes eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Harmful if swallowed. Do not contaminate water used for irrigation or domestic purposes. Do not graze treated areas. Do not feed clippings to livestock.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Not for use on or under ornamental trees, shrubs, bedding plants, or garden plants. Runoff and drift from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Atrazine can travel (seep or leach) through soil and can enter ground water which may be used as drinking water. Atrazine has been found in ground water. Users are advised not to apply Atrazine to sand and loamy soils where the water table (ground water) is close to the surface and where these soils are very permeable, i.e. well-drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of ground water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in manner consistent with its labeling. Not for use on turf being grown for sale or for commercial use as sod, or for commercial seed production, or for research purposes. Apply only when grass is dry.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in original labeled container in a cool, dry, locked place out of reach of children and domestic animals. Do not store near food or feedstuffs. To prevent cross contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides, or seeds. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not reuse empty bag. Securely wrap original container in several layers of newspaper and discard in trash.

WHAT THIS PRODUCT DOES

This product, "St. Augustine & Centipede Weed & Feed" controls and prevents weeds while feeding your lawn.

CONTROLS these common weed:

- | | | | |
|--------------------------------|-----------------------------------|-------------------------------------|---------------------|
| Annual Bluegrass (poa annua) | Chickweed (Common & Mouseear) | Fragrant Cudweed (Rabbit Tobacco) | Pepperweed |
| Annual Crabgrass | Common Purslane | Henbit | Purslane |
| Sowthistle | Cudweed | Knotweed | Red Sorrel |
| Annual Clover | Cutleaf Evening Primrose | Lespedeza | Rescue Grass |
| Black Medic | Dandelion | Mat Lippia (Maichweed) | Rescue Grass |
| Blue Toadflax | Dichondra | Moneywort | Sandspur |
| Bull Paspallum | Dollarweed | Narrowleaf Vetch | Shepards Purse |
| Buckhorn | Florida Betony (Rattlesnake weed) | Oxalis (Common, Yellow, Woodsorrel) | Spurweed |
| Plantain | Florida Pusley | Parsley-Piert | Stinging Nettle |
| Bur Clover | | | Swinecrest |
| Carolina Geranium (Cranesbill) | | | Venus Looking Glass |
| | | | Virginia Creeper |
| | | | Wild Mustard |

PREVENTS these annual weeds: Crabgrass, Sedges, Spotted Spurge, Bull Paspallum.

QUALITY FERTILIZER

Provides a quick-greening, long-feeding turf fertilizer. Contains quick release nitrogen to green your lawn in a few days and slow release nitrogen to feed for up to three months. Contains phosphorous and Potash to help build quality turf.

NET WEIGHT 16 POUNDS (7.3 Kg)

IMPORTANT USE PRECAUTIONS

- Do not apply on or under the branch spread (root zone) area of trees, shrubs, bedding plants, flowers, or garden plants.
- Do not apply to lawns containing grasses or mixtures of grasses other than those listed on the label or injury to these grasses may result.
- Do not apply to newly seeded Zoysia, Centipede, and Carpetgrass lawns until after the new grass has been mowed at least twice.
- Do not plant sprigs for one month, or sow new grass seed for six months following application of this product. May be applied to newly sprigged St. Augustine and Zoysia lawns one week after sprigging to prevent competition from germinating weeds.
- Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Apply this product only when grass blades are dry. When daytime temperatures are 80° F or above, water sufficiently to rinse particles off the grass blades.

TO CONTROL GROWING BROADLEAF WEEDS

Apply any time weeds are actively growing. If complete control is not obtained a second treatment may be required 2 months after the first. Apply in the fall to control Henbit, Chickweed and other winter weeds.

TO PREVENT ANNUAL WEEDS

Apply in January, February or early March to prevent germination of annual weeds such as crabgrass, Bull Paspallum, Sedges (from seed) and Spotted Spurge. Do not apply to Centipede or Carpetgrass lawns during the period of winter dormancy.

SPREADER SETTINGS

Always apply fertilizer with a spreader.	
Sta-Green (R) Brdcast	35
Cyclone B1-84	425
Central	35
Central Drop	60
Scotts Speedy Green	35
Ortho Rotary	25
Ortho Drop	40

For best results, water in product after application as fertilizers will not work without water. For a beautiful, healthy lawn, get on a program of lawn care. Apply fertilizer, weed controls, and pest controls on a regular basis.

GUARANTEED FERTILIZER ANALYSIS

Total Nitrogen (N) - - - - - Variable %
 X X% Ammoniacal Nitrogen
 X X% Urea Nitrogen
 Available Phosphoric Acid (P2O5) - - - - - Variable %
 Soluble Potash (K2O) - - - - - Variable %
 Sulfur (S) X %; Copper (Cu) X %; Iron (FE) X %; Zinc (Zn) X %
 Plant nutrients derived from Ammoniated Phosphate, Urea, Muriate of Potash, and Sulfur Coated Urea, Iron oxide, Copper Oxides, Zinc (Zn) Zinc Oxides Potential Acidity Equivalent to XXX lbs. of Calcium Carbonate/ton.
 Chlorine not more than . . . X %;
 Slowly Available Urea Nitrogen from Sulfur Coated Urea; . . . X %

NOTICE: BUYER AND USER ASSUMES ALL RISK AND LIABILITY OF USE, STORAGE AND/OR HANDLING OF THIS MATERIAL NOT IN ACCORDANCE WITH THE TERMS OF THIS LABEL (D-8660-18 PM512-21-94)

PURSELL
INCORPORATED 1904

P.O. Box 540 - Sylacauga, AL 35150

BEST AVAILABLE COPY