

(Fertilizer/Team Label)

10F4
PM 23
8660-19

FRONT PANEL

STA-GREEN (R)

CRABGRASS PREVENTER PLUS LAWN FERTILIZER

28-3-6

Treats 5,000 Square Feet

Prevents crabgrass, foxtail, barnyardgrass, Poa annua, goosegrass, and green and yellow foxtail.

Fertilizes your lawn and golf course turf with a quality, long-lasting fertilizer.

ACTIVE INGREDIENTS:

N-butyl - N-ethyl-alpha, alpha,	
alpha-trifluoro-2,6-dinitro-p-toluidine	0.85%
trifluralin: alpha, alpha,	
alpha-trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine	0.43%
INERT INGREDIENTS:	98.72%
TOTAL:	100.00%

EPA Reg. No. 8660-XX

EPA Est. No. 8660-AL-1

KEEP OUT OF THE REACH OF CHILDREN
CAUTION

See Back Panel for Additional Precautions

NET WT. 18 LBS. (8.2 kg)

REAR PANEL

STA-GREEN (R)

CRABGRASS PREVENTER PLUS LAWN FERTILIZER

GUARANTEED ANALYSIS 28-3-6


Total Nitrogen (N)	28%
1.2% Ammoniacal Nitrogen	
26.8% Urea Nitrogen*	
Available Phosphoric Acid (P205)	3%
Soluble Potash (K20)	6%

Primary Plant Food Sources: Sulfur Coated Urea, Urea, Diammonium Phosphate, and Muriate of Potash.

*Part of the nitrogen in this product has been sulfur-coated to provide 6.0% coated slow release nitrogen.

Chlorine (Cl), not more than 5.2%

FLA. No. _____



KEEP OUT OF REACH OF CHILDREN
CAUTION
PRECAUTIONARY STATEMENTS ✓

Hazards to Humans and Domestic Animals

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust or contact with skin or eyes. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt, and impermeable gloves when handling this product.

✓ **Statement of Practical Treatment**

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

If on skin: Wash with plenty of soap and water. Get medical attention if irritation persists.

If in eyes: Flush with plenty of water. Get medical attention.

If inhaled: If discomfort occurs, remove individual to fresh air. Get medical attention if breathing difficulty occurs.

Environmental Hazards ✓

Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes.

STORAGE AND DISPOSAL ✓

Storage: Store in original container only. In case of spill, contain material and dispose as waste.

Pesticide Disposal: Do not contaminate water, food, or feed by storage and disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. ✓

Container Disposal: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. ✓

✓ STA-GREEN Crabgrass Preventer Plus Lawn Fertilizer is a preemergence herbicide which is applied to established turfgrasses for the control of annual weeds such as crabgrass, Poa annua, goosegrass, foxtail, and banyardgrass. Apply this product before these annual weeds begin to germinate since it does not control established weeds. Contains a quality turf fertilizer to green your lawn in a few days and keep it green for two to three months.

For use on: ✓

Kentucky bluegrass Fescue St. Augustinegrass
Perennial ryegrass Zoysiagrass Bahiagrass
Centipedegrass Bermudagrass Bentgrass Fairways

✓ This product will not harm most nearby established ornamental trees and shrubs.

RATES, REQUENCY AND TIMING OF APPLICATION

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply to prevent crabgrass and other annual weeds in the Spring at the NORMAL rate one to two weeks before germination (before dandelions reach the puffball stage).

Apply in August or September to prevent germination of Poa annua. Apply at the NORMAL rate on cool season grasses (fescue, bluegrass, etc.) and at the HEAVY rate on warm season grasses (bermuda, zoysia, centipede, St. Augustine, Bahia, etc.).

Do not aerify, rake, verticut or otherwise disturb the soil after application. This can disturb the chemical barrier on the soil surface.

✓ Do not reseed for 12 to 16 weeks after application as the preemergence herbicide will also prevent grass seed from germinating.

Do not apply to newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass or zoysiagrass until these turfs are well established. Do not apply to dichondra.

Always apply product evenly over the turfgrass area with a spreader. Never apply by hand.

Suggested Spreader Settings

	<u>NORMAL</u>	<u>HEAVY</u>
Sta-Green	5½	6
Scotts Drop	6½	6 3/4
Scotts Rotary	6	H
Ortho Drop	6½	7
Ortho Rotary	1	1½
Sears Drop	8	9
Sears Rotary	4½	5
Central Drop	8	8½
Central Rotary	4½	5
Cyclone Drop	3	3½
Cyclone Rotary	4½	4½

Note: The above spreader settings are approximate. Differences may occur due to condition of spreader, speed of application, or roughness of area treated. Calculate the size of area to be treated and do not over or under apply. Close spreader when filling, turning, or stopping.

For Best Results With Your Lawn: Get on a program of lawn care. Apply crabgrass preventer in the Spring, Weed & Feed in the late Spring/Summer, insect controls when needed, and a good fall/winter fertilizer (this is the most important feeding of the year). Water your lawn during periods of dry weather. If we can be of further assistance, please write for our free lawn care information guide.

Money Back Guarantee

Net Wt. 18 lbs (8.2 Kg)

STA-GREEN
A Division of Parker Fertilizer Company
P.O. Box 540
Sylacauga, Alabama 35150