

8660-21

05/08/2000

1/2



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

WASHINGTON, D.C. 20460

Randy R. Simon
Purcell Industries, Inc.
P.O. Box 540
Sylacauga, AL 35150

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

MAY 8 2000

Subject: Sta-Green Sevin brand Carbaryl Insecticide Lawn Insect Control
EPA Registration No. 8660-21
Resubmission Amendment to update label as per the Carbaryl GDE.
Amendment dated February 4, 2000.

Dear Mr. Simon:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling change indicated below before you release the product for shipment bearing the amended labeling:

1. The chart containing the spreader settings must be revised so that the setting for the three different spreaders at all four application rates is not the same. It presently has a "XX" for each spreader at each of the four application rates.

Submit two copies of your final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records.

Sincerely,

A handwritten signature in cursive script that reads "George Tompkins".

George Tompkins, Ph.D., Entomologist
Insecticide-Rodenticide Branch
Registration Division (7505C)

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente*

FIRST AID - STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician of Poison Control Center immediately. Induce vomiting by giving victim 1 or 2 glasses of water and touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person. **IF IN EYES:** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN:** Remove contaminated clothing and wash affected area with soap and water. Get medical attention if irritation persists. **IF INHALED:** Remove victim to fresh air. Apply Artificial Respiration if indicated. Get medical attention.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is Antidotal.

SEE SIDE/BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact eyes, skin and clothing. Wash thoroughly after handling and before eating or smoking. Change to clean clothing daily. Do not contaminate feed or foodstuffs.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use such as sod, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in interior plantscapes, ornamental gardens or parks, or on golf courses or lawns and grounds.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open container. In case of spill; Avoid contact, isolate area and keep out animals and unprotected persons. Confine spills.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved wastes disposal facility. **CONTAINER DISPOSAL:** Do not reuse empty bag. Completely empty bag into equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

GENERAL INFORMATION

FOR LAWNS, ORNAMENTAL & RECREATIONAL TURF

Use when insects first appear and/or plant damage is imminent. For best control of Root Feeding insects such as grubs mow lawn first; apply material and then irrigate 1/4" to move insecticide down to soil level. It is not necessary to "water in" the granules for surface insect control. *See watering limitations for Sod Webworms. Keep children and pets off treated areas until grass is dry. Repeat applications may be made at a 30 day intervals up to 3 treatments per season for control of new infestations of Sod Webworms.

SURFACE FEEDING INSECTS

FOR ANTS, ARMYWORMS, CENTIPEDES, CHIGGERS, CUTWORMS, EARWIGS, FALL ARMYWORMS, FIERY SKIPPERS, GRASSHOPPERS, JUNE BEETLES, LEAFHOPPERS, MILLIPEDES, SOW BUGS, SPITTLE BUGS, SPRINGTAILS, TICKS, AND YELLOWSTRIPED ARMYWORMS.

Apply 0.75 lb./1,000 sq.ft. when insects become a problem or their damage is imminent. This is equivalent to 2 lbs. carbaryl per acre. Refer to General Directions.

FOR CHINCH BUGS AND SOD WEBWORMS

Apply 0.75 lb./1,000 sq.ft. of turf which is equal to 2 lbs. Carbaryl/Acre. For Chinch Bugs treat entire area to prevent their migration. *For Sod Webworms do not irrigate treated areas for 24 to 48 hours after treatment. Repeat applications may be made at 30 day intervals or as needed for control of new infestations. For heavy infestations increase rate to 1.5 lb. per 1,000 sq.ft. and treat entire area.

EPA REG. NO. 8660-21

EPA Est. No. 8660-AL-1

Net Weight 4; 10; 15 Pounds

In EPA Letter Dated:

MAY 8 2000

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 8660-21

MURSELL INDUSTRIES, INC.

P.O. Box 540 · Sylacauga, AL 35150

Nuisance pests around the outside of homes; such as Fire Ants and Fleas. Also for control of Ticks that Vector Lyme Disease

FOR ROOT FEEDING INSECTS

FOR CONTROL OF WHITE GRUBS (SUCH AS JAPANESE BEETLE, CHAFER BEETLES, AND PHYLLOPHAGA SPP. LARVAE), BLUE-GRASS BILLBUGS, FLEAS, EUROPEAN CRANE FLY LARVAE, AND BLACK TURFGRASS ATAENIUS

Apply 1.5 lb. per 1,000 sq.ft. of turf area (equal to 4 lbs. A./Acre) during late March through May or July to early September when Grubs are feeding near the soil surface or as recommended by local agricultural extension agents. Water or irrigate turf thoroughly soon after treatment to move the insecticide into the root feeding zone. See General Directions.

For CRANE FLY LARVAE apply at the same rate in early spring; April 1 to April 15th or as recommended by the local Extension Service. Irrigate area after treatment to move granules into root feeding zone.

FOR FLOWERING PLANTS AND SHRUBS IN LANDSCAPE and FOUNDATION PLANTINGS

For CUTWORMS, ARMYWORMS, SOWBUGS, MILLIPEDES and CENTIPEDES which attack ornamental and flowering plants in shrubbery and foundation plantings apply 1.5 lb. granules evenly over 1,000 sq.ft. (20 x 50 ft.). Repeat as needed for control up to 3 times per year.

SPREADER SETTINGS

Spreader	Pounds of Product to use per 1,000 Sq.Ft.			
	0.75 lb.	1.5 lbs.	2.0 lbs.	3.0 lbs.
Cyclone B1	X X	XX	XX	XX
Scott Rotary R-8	X X	XX	XX	XX
Scott Drop Spreader	X X	XX	XX	XX

TREAT LAWNS AND PERIMETER WOODED AREAS FOR CONTROL OF TICKS THAT VECTOR "LYME DISEASE"

For control of juvenile and adult Ixodes spp. (Deer Tick and Black Legged Tick) and Amblyomma spp. ticks (Lone Star Ticks) which vector Lyme Disease. Apply granules at the rate of 1.5 to 3.0 pounds of product per 1000 sq.ft. with appropriate equipment to ensure uniform coverage. Target applications to control juvenile ticks (larvae and nymphs) in late spring or early summer (May/June). Control of adult ticks can be achieved with late summer and fall applications (Aug/Sept). Make applications to lawns, shrub beds, ornamental plantings, wooded areas and around outside perimeters of home and other buildings. Do not make spot applications. Treat entire lawn and perimeter, wooded areas and property boundaries where exposure to ticks may occur. For optimum results, irrigate treated areas after application to apply chemical from the granules. Ticks may be reintroduced from surrounding areas on host animals. Retreat as necessary to maintain adequate control levels.

NUISANCE PEST CONTROL AROUND THE OUTSIDE OF HOMES, PORCHES, PATIOS AND BUILDINGS.

For control of Ants, Armyworms, Ticks, Cutworms, Centipedes, Crickets, Earwigs, Firebrats and Silverfish, Fleas, Millipedes, Sowbugs, Scorpions and Spiders; Apply granules uniformly in an 8 foot band around the building. Apply granules at the rate of 2 pounds per 1,000 sq.ft. to shrub and flower beds, foundations, ornamental plantings and lawn and soil areas immediately adjacent to the building. For most effective control treated area should be lightly watered after application to remove the chemical from the granules. Repeat treatments as necessary to control new infestations.

FIRE ANT CONTROL /MOUND TREATMENT.

For control of Fire Ants in mounds occurring in lawns, turf and premise areas, apply 2 to 3 ounces (8 to 14 tablespoonfuls) of granules over and around each mound area. Then gently sprinkle the treated areas with a small amount of water (1/4 to 1/2 gal) to release the chemical from the granules. Do not wash granules away from mound. For best results apply in cool weather (65° to 80°) or early morning or late afternoon. Retreat new mounds as they may appear.

NOTICE: This product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use. Manufacturer is not responsible for the use of this product contrary to the label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to the manufacturer and/or seller, and buyer assumes the risk of any such use. (8660-21-NU-LAB-P88-102-04-00)