

8660-21

STA-GREEN CARBARYL INSECTICIDE LAWN INSECT CONTROL

Controls Fleas, Ticks, Chinch Bugs, White Grubs, and Webworms, Armyworms, Ants, Fire Ants, Billbugs and Other Listed Pests • For Fire Ants • For Ticks That [Vector] [Transmit] Lyme Disease • Kills Nuisance Pests Around Homes • Kills Nuisance Pests Around The Outside Of Home [Such As Fire Ants And Fleas] • Also For Control Of Ticks That [Vector] [Transmit] [Carry] Lyme Disease • Kills Grubs and Lawn Invading Insects • Kills Insects Above and Below the Soil Surface [Ground] • Effective Grub Control • Kills Grubs on Contact • Penetrates Deeply [Into The Soil] • Kills Insects That Damage Lawns • Penetrates The Ground to Kill Root Eating Insects • Releases Into Soil To Kill Hard To Reach Grubs & Insects • Attacks [Kills] Grubs Where They Live • Deep Penetrating Formula • 15 lbs. treats up to 10,000 Sq.Ft.

ACCEPTED
DEC 6 2002
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the purpose of registration under EPA Reg. No. 8660-21

Active Ingredient: Carbaryl 6.30%
Other Ingredients: 93.70%
[TOTAL] 100.00%

KEEP OUT OF REACH OF CHILDREN
CAUTION

[*PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente*]
See [back] [side] panel for additional precautionary statements.

NET WT 4 LB. (1.8 kg); 10 LB. (4.5 kg); 15 LB (6.8 kg)

STOP. READ ENTIRE LABEL BEFORE USE

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. For outdoor use around the home only.

GENERAL INFORMATION

[FOR] LAWNS

Use when insects first appear and/or plant damage is [noticed] [imminent]. For best control of **root feeding insects** such as grubs mow lawn first; apply material and then apply 1/4" of water to move insecticide down to soil level. It is not necessary to "water in" the granules for surface insect control. *See watering limitations for Sod Webworms. Keep children and pets off treated areas until grass is dry. Repeat applications may be made at 30-day intervals up to 3 treatments per season for control of new infestations of Sod Webworms.

SURFACE FEEDING INSECTS

For Ants, Armyworms, Centipedes, Chiggers, Cutworms, Earwigs, Fall Armyworms, Flary Skippers, Grasshoppers, June Beetles, Leafhoppers, Millipedes, Sowbugs, Spittlebugs, Springtails, Ticks And Yellowstriped Armyworms: Apply 1.5 lbs. per 1,000 sq. ft. when insects become a problem or their damage is imminent. Refer to General Information.
For Chinch Bugs And Sod Webworms: Apply 3 lbs. per 1,000 sq. ft. of turf. For Chinch bugs treat entire area to prevent their migration.

*For Sod Webworms do not irrigate treated areas for 24 to 48 hours after treatment. Repeat applications may be made at 30-day intervals or as needed for control of new infestations.

FOR ROOT FEEDING INSECTS

For Control of White Grubs (such as Japanese Beetle, Chafer Beetles, and *Phyllophaga spp.* Larvae), Bluegrass Billbugs, Fleas, European Crane Fly Larvae, and Black Turfgrass *Ataenius*: Apply 3 lbs. per 1,000 sq. ft. of turf area during late March through May, or July to early September when grubs are feeding near the soil surface or as recommended by local agricultural extension agents. Water or irrigate turf thoroughly soon after treatment to move the insecticide into the root feeding zone. See General Directions.

FOR CRANE FLY LARVAE: Apply 3 lbs. per 1,000 sq. ft. of turf area in early spring: April 1 to April 15 or as recommended by the local Extension Service. Irrigate area after treatment to move granules into root feeding zone.

12/6/2002

1/3

FOR FLOWERING PLANTS AND SHRUBS

For Cutworms, Armyworms, Sowbugs, Millipedes and Centipedes which attack ornamental and flowering plants in shrubbery and foundation plantings: Apply 1.5 lbs. of granules evenly over 1,000 sq. ft. (20 ft x 50 ft). Repeat as needed for control up to 3 times per year.

SPREADER SETTINGS		
Spreader Make and Model	1.5 lbs. per 1000 sq. ft.	3 lbs. per 1000 sq. ft.
Scotts SpeedyGreen 3000	2	4
Scotts AccuGreen 3000	5	8

NOTE: These settings are approximate due to variability and wear of spreader. Check settings and calibrations on a small area before making total application.

TREAT LAWNS AND PERIMETER WOODED AREAS FOR CONTROL OF TICKS THAT VECTOR LYME DISEASE

For Control of juvenile and adult *Ixodes spp.* (Deer Tick and Black Legged Tick) and *Amblyomma spp.* Ticks (Lone Star Ticks) which vector Lyme Disease: Apply granules at the rate of 3 lbs. of product per 1,000 sq. ft. with appropriate equipment to ensure uniform coverage. Target applications to control juvenile ticks (larvae and nymphs) in late spring or early summer (May/June). Control of adult ticks can be achieved with late summer and fall applications (Aug/Sept). Make applications to lawns, shrub beds, ornamental plantings, wooded areas and around outside perimeters of home and other buildings. Do not make spot applications. Treat entire lawn and perimeter wooded areas and property boundaries where exposure to ticks may occur. For optimum results, irrigate treated areas after application to remove chemical from the granules. Ticks may be reintroduced from surrounding area on host animals. Retreat as necessary to maintain adequate control levels.

NUISANCE PEST CONTROL AROUND THE OUTSIDE OF HOMES, PORCHES, PATIOS AND BUILDINGS

For control of Ants, Armyworms, Ticks, Cutworms, Centipedes, Crickets, Earwigs, Firebrats and Silverfish, Fleas, Millipedes, Sowbugs, Scorpions, and Spiders: Apply granules at the rate of 1.5 lbs. per 1,000 sq. ft. to shrub and flowerbeds, foundations, ornamental plantings and lawn and soil areas immediately adjacent to the building. For most effective control, treated area should be lightly watered after application to remove the chemical from the granules. Repeat treatment as necessary to control new infestations.

FIRE ANT CONTROL / MOUND TREATMENT

For control of Fire Ants in mounds occurring in lawns, turf and premise areas apply 2 to 3 ounces (8 to 14 Tbsp.) of granules over and around each mound area. Then gently sprinkle the treated areas with a small amount of water (¼ to ½ gal) to release the chemical from the granules. Do not wash granules away from mound. For best results apply in cool weather (65° to 80°) or early morning or late afternoon. Retreat new mounds as they may appear.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Keep out of reach of children and animals. Store in original container in a cool, dry place and avoid excess heat. **DISPOSAL: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain

PRECAUTIONARY STATEMENTS**Hazards To Humans And Domestic Animals**

CAUTION: Harmful if swallowed or inhaled. Avoid breathing dust. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling and before eating or smoking.

Continued on page 3.....

First Aid

The table below describes the first aid procedures for incidents involving Sta-Green Carbaryl Insecticide Lawn Insect Control.

<i>In the case of accidental ...</i>	<i>Immediately ...</i>
Swallowing	<ul style="list-style-type: none"> • Call a Poison Control Center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by a Poison Control Center or doctor. • Do not give anything by mouth to an unconscious person.
Contact with Eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 10-15 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a Poison Control Center or doctor for treatment advice.
Contact with Skin or Clothing	<ul style="list-style-type: none"> • Take off contaminated clothing • Rinse skin immediately with plenty of water for 15-20 minutes • Call a Poison Control Center or doctor for treatment advice
Inhalation of dust	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, and then give artificial respiration, preferably mouth-to-mouth if possible. • Call a Poison Control Center or doctor for further treatment advice.
Have the product container or label with you when calling a Poison Control Center doctor or going for treatment.	
Note to Physician:	
Carbaryl is an N-methyl carbamate that inhibits cholinesterase. Atropine is antidotal.	
Hot Line Number: 1-800-332-5553	

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

[Questions or Comments? Call 1-800-332-5553]

Made in the USA by
Earth Care
Division of United Industries Corporation
P.O. Box 142642
St. Louis, Missouri 63114-0642

EPA REG. NO. 8660-21

EPA EST. NO. 9688-MO-1

© 2002