8660 - 34 Pm 23

page 18 5

18 JUN 1992

Mr. Daniel Paradiso, ur. alble farker fertilizer 201 west Forth Street Sylacoute At. ..5156

cear Nr. Paradisu:

Subject: Vertegreen Pricem 71

EFA Registration No. 8660-25

vertogand MCPP diover - Chickneed Killer

SPA Registration No. 8660-24

Vertacreen Reed And Feed

TPA Pogistration No. 8668-34

Protect o Way

IPA Resistration No. 6660-104

Ye.tu-Creen St. Augustine Reen & feed

LPA Registration No. 8660-147

Re: Latel Amendment

Your Soumission Dates May Lc. 1992

The labeling referred to above submitted in connection with registration under the Federal Insectivide rungicide and Rogenticide act, as akended is acceptable, provided that under the Environmental Hazers Section of the label replace the statement" Do not apply directly to water or wetlands" to read "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark." Please submit five (5) copies of your final printed labels incorporating this amendment. A stapped copy of labeling is enclosed for your records.

Sincerely

BEST AVAILABLE COPY

Joanne I. Miller Product Hanager (73) Fungicide-Herbicide Branch Registration Division (H-1505C)

CONCURRENCES										
SYMBOL	H-750556	osure								
	Metra	Į į		{	Į.		{	\		
DATE	6/17/12			}						
EPA Form	1320-1 (12-70)	 _			·		OFFICI	AL FILE COPY		

eu.s. Gpo 1985 467 853

8660-34 1 of 5

(FRONT PANEL)

VERTAGREEN®

FERTILIZER

FOR PROFESSIONAL USE

WITH 2,4-D & MCPP

ACTIVE INGREDIENTS:			
*Isooctyl Ester of 2	2,4-Dichlorophenoxyacetic Acid		0.87%
**Isooctyl Ester of 2	2-(2-methyl-4-chlorophenoxy) propionic acid		0.92%
	TOTAL	1	L00.00%
Equivalent to: *0 577% of 2 4-Dichl	orophenoxyacetic Acid		

**0.60% of 2-(2-methyl-4-chlorophenoxy) propionic acid

KEEP OUT OF THE REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 50 LBS.

Treats Up To 12,500 Sq. Ft.

MONORPH ME DEALTH COURT MEPA LAND COURT

EPA Reg. No. 8660-34

Minder to Francisco de la Managarda de la Caracter de la Caracter

EPA Est. 9198-OH-1

BEST AVAILABLE COPY

(BACK PANEL)

NET WEIGHT 50 LBS.

TREATS UP TO 12,500 SQ. FT.

VERTAGREEN®

FERTILIZER

FOR PROFESSIONAL USE

WITH 2,4-D & MCPP

GUARANTEED ANALYSIS

Total Nitrogen (N)	15.00%
15.00% Ammoniacal Nitrogen Available Phosphoric Acid (P_2O_5)	5.00% 5.00%
Sources of Plant Nutrients: Ammonium Sulfate, Ammonium Phosphate, Potash	, Muriate of
Chlorine (Cl), Maximum	5.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash skin with soap and water.

IF IN EYES: Flush eyes with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS

Do not apply to lakes, streams, or public waters. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes.

BEST AVAILABLE COPY

8660-34 3 of 5

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is designed for control of broadleaf weeds in lawns such as: Bindweed, Black Medic, Blue Lettuce, Buckhorn, Burdock, Chickory, Chickweed, Cinquefoil, Ground Ivy, Hawkweed, Heal-All, Henbit, Jewelweed, Jimson Weed, Knotweed, Kochia, Lambsquarter, Mallow, Morning Glory, Mustard, Nettles, Oxalis, Pennywort, Peppergrass, Pigweed, Plantain, Poverty Weed, Purslane, Ragweed, Shepherd's Purse, Smartweed, Speedwell, Spotted Spurge, Thistle, Vervains, Vetch, White Aster, Wild Radish, Yarrow.

READ THE FOLLOWING CAREFULLY BEFORE APPLICATION.

This product may be used on Bluegrass, Fescue, Rye, Bermuda, Bahia, and Zoysia lawns. DO NOT USE on Dichondra, Carpetgrass, St. Augustinegrass, Bentgrass, Centipede or ground cover. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables, or ornamental shrubbery. If accidentally deposited on flowers or shrubs, hose off immediately. Vapors from this product may injure susceptible plants in the immediate vicinity.

WHEN TO APPLY

Best results are obtained when weeds are growing rapidly in spring or early fall. To avoid injury to new growth, new lawns should not be treated until after the fourth mowing. Established lawns may be injured if treated when in need of water. Grass may be injured if treated in periods of high heat and humidity. For this reason, apply only when the daytime temperature is not expected to exceed 80°F within 48 hours. Delay planting grass seed until 4 weeks after treatment.

HOW TO APPLY

This 50 pound bag should be used on 12,500 sq. ft. Apply evenly with a lawn spreader, either as a one pass application or a two pass application, crisscrossing the area at half the rate each time. Apply to lawn when the grass is wet after rain, sprinkling or dew so granules will stick to weeds. Do not water or mow lawn for at least two days after treatment. Lawns that are heavily overrun by weeds and that are in poor condition may require more than one application of this product. A suggested second application should be made in approximately 30 days.

STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

8660-34 4 of 5

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the rearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Completely empty bog into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

MANUFACTURER warrar that this product (1) conforms to the ingredient statement on the labland (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CALEFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by:

PARKER FERTILIZER CO. 201 4TH STREET SYLACAUGA, AL 35150

Date Printed: 05/07/92

Date Approved: Supersedes: Label Change: