

8660-37
1 of 5

(FRONT PANEL)

VERTAGREEN®

FERTILIZER

FOR PROFESSIONAL USE

WITH BALAN®

26-4-4

ACTIVE INGREDIENT:

N-butyl-N-ethyl-a,a,a,-trifluoro-2,6-dinitro-p-toluidine 2.41%

INERT INGREDIENTS: 97.59%

Total 100.00%

This product contains 2.41 lbs. Benefin per 100 lbs.
Balan® is the Trademark for Benefin, DowElanco


KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 40 LBS.

COVERS UP TO 20,000 SQ. FT.



2 8 5

8660-37
2 of 5

(BACK PANEL)

NET WEIGHT 40 LBS.

COVERS UP TO 20,000 SQ. FT.

VERTAGREEN®

FERTILIZER

FOR PROFESSIONAL USE

WITH BALANCE

26-4-4

GUARANTEED ANALYSIS

Total Nitrogen (N)	26.00%
1.6% Ammoniacal Nitrogen	
24.4% Urea Nitrogen*	
Available Phosphoric Acid (P ₂ O ₅)	4.00%
Soluble Potash (K ₂ O)	4.00%
Sulfur (S) Free	1.30%
Manganese (Mn)	0.05%
Zinc (Zn)	0.05%

Sources of Plant Nutrients: Urea, Sulfur Coated Urea, Ammonium Phosphates, and Muriate of Potash

Manganese Oxide, Zinc Oxide

BEST AVAILABLE COPY

Chlorine (Cl), Maximum 4.00%

*Includes 4.0% Coated Slow Release Urea Nitrogen derived from Sulfur Coated Urea

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes or clothing. In case of contact, flush with water. Do not contaminate foodstuffs or feed. Do not let livestock graze treated areas. Do not feed forage from treated areas to livestock.

STATEMENT OF PRACTICAL TREATMENT:

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, ponds or wetlands. Do not contaminate water by direct application, cleaning of equipment, or disposal of wastes. Apply this product only as specified on the label.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product provides plant nutrients and controls Smooth and Hairy Crabgrass, Green and Yellow Foxtail, Barnyardgrass and Goosegrass (Silver Crabgrass or Crowsfoot) by killing plants as seeds germinate. It does not control established weeds.

Use only on established lawns listed below which are tolerant to the herbicide in this product when applied as directed. SEE PRECAUTIONS.

Bluegrass (perennial)	Bermudagrass
Ryegrass (perennial)	Centipedegrass
Fescue (except fine-leaved)	St. Augustinegrass
Bahiagrass	Zoysiagrass

WHEN TO APPLY

Apply to established turf in the spring before germination of annual weeds listed above.

AMOUNT TO APPLY

Use 83 lbs. per acre or 1.9 lbs. per 1000 sq. ft. This rate is equivalent to 2.0 lbs. of active ingredient per acre.

HOW TO APPLY

Apply evenly over the turf without skips or misses. A more uniform application can be made by spreading half the required amount of product over the area to be treated and then applying the remaining half at right angles or crosswise.

RESEEDING

Chemicals which effectively control annual weed seeds may also affect new desirable turfgrass seedlings. Reseeding should be delayed for at least 6 weeks after application. When reseeding, it is necessary that proper moisture be maintained until grass seedlings are established. To assure satisfactory results after using this product, the amount of seed used should be increased.

4 8 5

8660-37

4 of 5

PRECAUTIONS

Do not apply in the spring to turfgrass planted the previous fall. Do not apply to Bentgrass putting greens or Dichondra turf. Do not apply to newly sprigged areas of Bermudagrass, St. Augustinegrass, or Centipedegrass until these turfs are well established. If verticutting is done, do not apply for 2 to 3 weeks.

STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by:

PARKER FERTILIZER CO.
201 4TH STREET
SYLACAUGA, AL 35150

Date Printed: 05/07/92
Date Approved:
Supersedes:
Label Change: