

yellow copy

PM 19 8660-43

10/13/6

JUL 2 1992

Mr. Daniel Paradiso, Jr.
The Andersons
d/b/a Parker Fertilizer Co.
201 4th Street
Sylacauga, AL 35150

Dear Mr. Paradiso:

Subject: Revised labeling
Vertagreen 25% Methoxychlor
EPA Registration No. 8660-43
Amendment application of May 22, 1992

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended is acceptable. A stamp copy is enclosed for your records.

Add the appropriate statement of practical treatment for "IF SWALLOWED....."

The statement "Do not apply directly to wetlands" as it appears in the environmental hazard statement must be changed to: "Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark."

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (H-7505C)

Enclosure

(FRONT PANEL)

VERTAGREEN®

25% METHOXYCHLOR

EMULSIFIABLE CONCENTRATE

ACTIVE INGREDIENTS:

| | |
|--|---------|
| Methoxychlor: 1,1,1-trichloro-2,2-bis(p-methoxyphenyl) | 25.00% |
| *INERT INGREDIENTS: | 75.00% |
| Total | 100.00% |

*Contains Xylene

KEEP OUT OF THE REACH OF CHILDREN


CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash skin with plenty of soap and water.
IF IN EYES: Flush with water for 15 minutes and get medical attention.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT ___ GALLONS



(BACK PANEL)

NET WEIGHT ___ GALLONS

VERTAGREEN®

25% METHOXYCHLOR

EMULSIFIABLE CONCENTRATE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes, or clothing. Wash thoroughly after handling.

WORKER PROTECTION

Do not enter or allow entry into treated areas until sprays have dried/dusts have settled/vapors have dispersed, as applicable, to perform hand labor tasks. A person may enter the area to perform other tasks only if the person is wearing the personal protection equipment listed on the label for a pesticide handler.

Do not apply this product in a way that will contact unprotected workers, either directly or through drift. Only protected handlers wearing long pants, long-sleeved shirts, socks, shoes and chemical resistant gloves may be in the area during application.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Aerial application over such sites is permissible only when they are not visible from above the tree canopy. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Do not graze or feed treated pea vines or pea hay to livestock.
Do not graze or feed treated cowpea vines or cowpea hay to livestock.
Do not graze or feed treated soybean hay or straw to livestock.
Do not feed treated corn to livestock.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

BEEF CATTLE, HOGS, SHEEP (except dairy animals): For controlling flies, lice, ticks and fleas, spray or dip using 1 to 2 gals. per 50 gals. of water. Make sure animals are wetted to the skin. Usually, one treatment is effective for about 3 weeks. For use in rubbing devices, mix 1 gal. with 4 gals. of fuel oil or kerosene.

FARM BUILDINGS: To control stable flies, house flies and fleas, mix 1 gal. in 5 gals. of water and apply 1 gal. of mixture to approximately 500 sq. ft. of surface; treating walls, floors, cracks, crevices, baseboards, bedding, coops, stalls, and pens. Such heavy deposits usually are effective for 3-6 weeks, unless removed by weathering or other means. Proper sanitary measures should also be followed, including spraying or removal of breeding sources such as manure in and around buildings. **CAUTION:** When treating dairy barns, dairy animals should be removed, and feed and water protected during the treatment.

GRAIN STORAGE BINS: Spray empty bins thoroughly (after thorough cleaning) before storage of barley, corn, oats, rice, rye, sorghum or wheat, with spray composed of 12 gals. mixed with 10 gals. of water at the rate of 1 gal. of spray per 500 sq. ft. of surface. This treatment will aid in control of Cadelle, Flat Grain Beetle, Saw-Toothed Grain Beetle, Lesser Grain Borer, Red Flour Beetle, Long-Headed Flour Beetle, Rice Weevil, Granary Weevil, Confused Flour Beetle, Foreign Grain Beetle and Hairy Fungus Beetle.

NOTE: In the following paragraphs, the figure in parentheses following each crop name represents the maximum number of gallons of Methoxychlor Emulsifiable Concentrate that should be applied per acre.

TREE FRUITS: Apply to Apples (7-1/2), Apricots* (3), Cherries (4), Nectarines* (1), Peaches* (3), Pears (4), Plums (4), Prunes (4), and Quinces (3) for control of Apple Maggot, Cankerworm, Cherry Fruit Fly, Cherry Fruitworm, Codling Moth, Japanese Beetle, Leafhoppers, Pear Slug, Plum Curculio and Tent Caterpillar. Start applications at petal-fall stage, using 1/2 to 3/4 gals. per 100 gals. of water (1-1/2 to 2 tablespoonsful per gallon) and repeat at 7 to 14 day intervals as needed. Do not apply later than 7 days (*21 days) before harvest.

SMALL FRUITS: Apply to blackberries (1-1/2), Blueberries (1), Cranberries (1-1/2), Currants (1-1/2), Gooseberries (1-1/2), Grapes (3), Loganberries (1-1/2), Raspberries (1-1/2), and Strawberries (1), to control Cherry Fruitworm, Cranberry Fruitworm, Grape Berry Moth, Grape Leaf Skeletonizer, Japanese Beetle, Leafhoppers, Plum Curculio, Rose Chafer, San Jose Scale (Crawlers), Spittlebug and Strawberry Weevil. Start applications at first sign of infestation and repeat at 7 to 14 day intervals as needed, using 1/2 to 3/4 gal. per 100 gals. of water (1-1/2 to 2 tablespoonsful per gallon). Do not apply later than 14 days before harvest.

BEST AVAILABLE COPY

GARDEN CROPS: Apply to Asparagus** (1), Beans (1), Beets* (1), Blackeyed Peas (1-1/2), Broccoli* (1), Brussels Sprouts* (1), Cabbage (1), Carrots* (1), Cauliflower (1), Collards* (1), Corn (1), Cucumbers (1), Eggplant (1), Kale* (1), Kohlrabi (1), Lettuce* (1), Melons (1), Peas (1), Peppers (1), Potatoes (1), Pumpkin (1), Squash (1), Tomatoes (1), and Turnips* (1) for control of Asparagus Beetle, Bean Leaf Beetle, Blister Beetle, Cabbageworm, Colorado Potato Beetle, Cucumber Beetle, Fall Armyworm, Flea Beetles, Japanese Beetle, Leafhopper, Mexican Bean Beetle, Pea Weevil, Squash Vine Borer and Webworm. Mix 1/2 to 3/4 gal. per 100 gals. of water (1-1/2 to 2 tablespoonsful per gallon) and apply at first sign of infestation, repeating at 7 to 14 day intervals as needed. Do not apply later than 7 days (*14 days, **3 days) before harvest.

FIELD AND FORAGE CROPS: Apply on Alfalfa (3/4), Cowpeas (3/4), Clover (3/4), Soy Beans (3/4), and foliage grasses (3/4) to control Alfalfa Caterpillar, Alfalfa Weevil Larvae, Blister Beetle, Armyworm, Cowpea Curculio, Clover Leaf Weevil, Fall Armyworm, Flea Beetles, Japanese Beetle, Leafhopper, Pea Weevil, Soybean Caterpillar, Spittlebug and Velvet Bean Caterpillar. Use 1/2 to 3/4 gal. per 100 gals. of water (1-1/2 to 2 tablespoonsful per gallon), starting at the first sign of infestation and repeating at 7 to 14 days intervals as needed. Do not apply later than 7 days before harvest or grazing.

STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY

WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. **FOLLOW DIRECTIONS CAREFULLY.** Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured By:

**PARKER FERTILIZER CO.
201 4TH STREET
SYLACAUGA, AL 35150**

Date Printed: 05/27/92

Date Approved:

Supersedes:

Label Change: