

8660-60


1 of 4

(FRONT PANEL)

GREENUP®

50% SEVIN®

WETTABLE POWDER



**ACTIVE INGREDIENT:**

Carbaryl (1-Naphthyl N-Methylcarbamate) . . . . .	50.00%
INERT INGREDIENTS: . . . . .	<u>50.00%</u>

TOTAL	100.00%
-------	---------

SEVIN® is the registered trademark of Union Carbide Corporation for 1-Naphthyl N-Methylcarbamate.  
U.S. Patent 2,903,476

KEEP OUT OF REACH OF CHILDREN

CAUTION


SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 6 OZ.

MAKES UP TO 20 GALLONS OF SPRAY

EPA Reg. No. 8660-60

EPA Est. 9198-QH-1



(BACK PANEL)

NET WEIGHT 6 OZ.

MAKES UP TO 20 GALLONS OF SPRAY

GREENUP●

50% SEVIN●

WETTABLE POWDER

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

##### CAUTION

HARMFUL IF INHALED OR SWALLOWED. Do not take internally. Avoid breathing of dust or spray mist. Skin contact may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after work. Wear regular long-sleeved work clothing. Change to clean clothing daily.

#### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting and seek medical attention immediately.  
IF IN EYES: Flush with water for 15 minutes and get medical attention.  
IF ON SKIN: Wash skin with plenty of soap and water.  
IF INHALED: Move from contaminated atmosphere and call a physician.  
NOTE TO PHYSICIAN: Carbaryl is a moderate reversible, cholinesterase inhibitor. Atropine is antidotal.

#### ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. This product is highly toxic to bees exposed to direct treatment of residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds when bees are visiting the treatment area. Consult your Cooperative Agricultural Extension Service for further information.

#### PHYSICAL OR CHEMICAL HAZARDS

Do not store or use near heat or open flame.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Controls Japanese Beetles, Mexican Bean Beetles, Squash Bugs and certain other insect pests, as listed, of fruit, vegetables and ornamentals.

50% Sevin Wettable Powder is a wettable spray powder. Measure in level tablespoonfuls. Keep spray mixture well agitated. Do not mix with lime, Bordeaux Mixture or other alkaline materials. It is compatible with most fungicides and miticides. Do not mix with oil base sprays.

**ORNAMENTALS:** Use 2 tablespoonfuls per gallon for Japanese Beetles, Flea Beetles, Bagworms, Leafhoppers, Elm Leaf Beetles, Apple Aphids, Blister Beetles, Leaf Rollers, Plant Bugs, Scale Insects, Box Elder Bugs and Tent Caterpillars. Apply when pests appear. Repeat as needed.

**APPLES AND PEARS:** Use 2 tablespoonfuls per gallon for Codling Moth, Pear Psylla, Apple Maggot, Plum Curculio, Fruit Tree Leaf Rollers, Japanese Beetles, Apple Aphids and San Jose Scale. Do not apply before second cover spray to avoid fruit thinning. Apply at 10 day intervals in July for Apple Maggot. May be used up to 1 day before picking.

**PEACHES AND APRICOTS:** Use 2 tablespoonfuls per gallon for Oriental Fruit Moth, Plum Curculio, Peach Twig Borer, Cat-Facing Bugs, San Jose Scale and Japanese Beetles. May be used 1 day before harvest on Peaches; 3 days on Apricots.

**GRAPES:** Use 1 tablespoonful per gallon for Leafhoppers, Grape Berry Moth and Japanese Beetles. May be applied up to harvest.

**STRAWBERRIES:** Use 2 tablespoonfuls per gallon up to 1 day before picking for Leaf Roller and Meadow Spittlebug control. Make 3 to 5 post bloom summer sprays as needed. May be used 1 day before harvest.

**BEANS:** Use 2 tablespoonfuls per gallon. Spray until all foliage is wet. Repeat in 7 to 10 days. For Mexican Bean Beetles, Leafhoppers, Flea Beetles, Stink Bugs and Lima Bean Pod Borers. May be used up to harvest. Do not feed vines to dairy or meat animals.

**CUCUMBERS, MELONS, AND SQUASH:** Use 2 tablespoonfuls per gallon. Apply as full wetting spray for Squash Bugs, Melon Worms, Cucumber Beetles and Pickle Worms.

**POTATOES:** Use 2 tablespoonfuls per gallon for Colorado Potato Beetle, Leafhoppers and Flea Beetles.

**TOMATOES, PEPPERS, EGGPLANTS:** Use 2 tablespoonfuls per gallon for Tomato Fruit Worm, Flea Beetles, Fall Armyworms, Stink Bugs and Leafhoppers. Apply up to day of picking.

8660-60  
4 of 4**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

**WARRANTY STATEMENT**

**MANUFACTURER** warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. **FOLLOW DIRECTIONS CAREFULLY.** Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by:

**PARKER FERTILIZER CO.  
201 4TH STREET  
SYLACAUGA, AL 35150**

Date Printed: 01/28/93  
Date Approved:  
Supersedes:  
Label Change:

