

8660-68
1 of 5

(FRONT PANEL)

PATTERSON'S

GREENUP®

FERBAM

FUNGICIDE

ACTIVE INGREDIENT:

Ferbam (Ferric dimethyl-dithiocarbamate) 76.00%

INERT INGREDIENTS: 24.00%

TOTAL 100.00%

11 SEP 1992

KEEP OUT OF REACH OF CHILDREN

CAUTION

8660-68

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 8 OZ.

MAKES UP TO 28 GALLONS OF SPRAY

295
8660-68

2 of 5

(BACK PANEL)

NET WEIGHT 8 OZ.

MAKES UP TO 28 GALLONS OF SPRAY

PATTERSON'S

GREENUP®

FERRAM

FUNGICIDE

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

May cause irritation of eyes, nose, throat or skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Store away from children, domestic animals and foodstuffs. Keep away from fire or sparks.

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Flush with plenty of water.

IF IN EYES: Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

SPRAYS: For best results, add enough water to the proper amount of Ferbam to make a thin, smooth paste; add paste to water in sprayer and mix thoroughly. Use 2 level tablespoonfuls per gallon of water, spraying thoroughly to cover bottom and top surfaces of foliage and fruit.

Ferbam can be dusted, if desired, for which the usual mixture contains 1 part Ferbam with 9 parts of diluent, preferably talc or fine clay.

Begin applications before diseases are apparent and continue at 7 to 14 day intervals as long as necessary. Do not apply later than 7 days before harvest or picking, unless otherwise indicated below.

Use Ferbam to control these diseases:

<u>CROP</u>	<u>CONTROL</u>
AZALEAS	Leaf Scorch
CAMELLIAS	Petal Blight
CARNATIONS	Leaf Spot, Alternaria and Septoria Blights, Rust
CHRYSANTHEMUMS	Septoria Leaf Spot, Rust
CYCLAMEN	Botrytis Rot, Leaf Spot
GERANIUMS	Bacterial Leaf Spot
JUNIPER	Rust
ROSES	Black Spot, Rust
SNAPDRAGONS	Rust, Leaf Spot
TULIPS	Botrytis Blight (Tulip Fire)
ASTER, BEGONIA, GARDENIA, HYDRANGEA, IVY, LAUREL	Leaf Spots
CUTTING-ROT, DAMPING-OFF Diseases (Except Hydrangea)	Dip cutting in spray mixture before planting

BEST AVAILABLE COPY

8660-68

4 of 5

<u>CROP</u>	<u>CONTROL</u>
APPLES	Scab, Cedar-Apple Rusts, Black Rot (Fruit and Leaf Spots), Brooks Spot, Apple Blotch, Bitter Rot, Sooty Blotch, Fly Speck
*APRICOTS	Green Rot, Shot Hole (Fungus)
**BLUEBERRIES	Botrytis Blight
CHERRIES	Brown Rot, Leaf Spot
**GRAPES	Black Rot
*PEACHES	Brown Rot (Early Season Only), Scab, Leaf Curl, Leaf Spot
PEARS	Scab, Leaf Blight, Fruit Spot, Sooty Blotch
**RASPBERRIES	Anthracnose, Rust, Leaf Spot
TOBACCO	Blue Mold in plant beds. Begin applications when leaves are the size of a dime or when Blue Mold is first reported in the area and repeat twice weekly until plants are set into field.
CABBAGE	Downy Mildew in plant beds.
LETTUCE	Damping-Off in plant beds.
SNAP or STRING BEANS	Anthracnose

Do not feed treated plant parts to meat or dairy animals.

*Do not apply later than 21 days before harvest.

**Do not apply later than 40 days before harvest. On Grapes, 7 days before harvest.

BEST AVAILABLE COPY

STORAGE AND DISPOSAL**FOR HOMEOWNER USE ONLY:**

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. **EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.**

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. **FOLLOW DIRECTIONS CAREFULLY.** Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by:

**PARKER FERTILIZER CO.
201 4TH STREET
SYLACAUGA, AL 35150**

Date Printed: 06/09/92
Date Approved:
Supersedes:
Label Change:

BEST AVAILABLE COPY