8660-83 1 of 4

(FRONT PANEL)

#### **VERTAGRENO**

# WHITE FLY SPRAY

ACCEPTED

ACCEPT

**READY-TO-USE** 

For Greenhouse Pests

(Contains 2 lbs. SBP-1382 per gallon)

ACTIVE INGREDIENT:

*Resmethrin														٠					24.30%
Related Compounds.						_													3.30%
**INERT INGREDIENTS: .																			72.40%
	TOLAL													100.00%					

\*Cis/trans isomers ratio maximum 30% ( $\pm$ ) cis and minimum 70% ( $\pm$ ) trans. \*\*Contains petroleum hydrocarbons.

# KEEP OUT OF REACH OF CHILDREN

# CAUTION

# SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET CONTENTS 1 QT. (32 FL. OZ.)

# (BACK PANEL)

# NET COMMENTS 1 QT. (32 FL. OZ.)

### VERTAGREEN

#### WHITE FLY SPRAY

#### PRECAUTIONARY STATEMENTS

# HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Harmful if swallowed. Avoid contact with eyes, skin or clothing. Avoid contamination of feed and foodstuffs. Avoid breathing of spray mist. Do not use, pour, spill or store near heat or open flame. Keep container closed.

### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Get medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation develops.

IF IN EYES: Flush with water for 15 minutes. Get medical attention if irritation develops or persists.

NOTE TO PHYSICIAN: Induced vomiting as first aid for this toxic substance may result in increased risk of chemical pneumonia or edema caused by aspiration of the hydrocarbon solvent. Vomiting should be induced only under professional supervision.

# ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Drift and runoff from treated sites may be hazardous to fish in adjacent waters. Do not contaminate water when disposing of equipment washwaters.

# DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

GREENHOUSE PESTS: For infestations of Whiteflies on Ageratum, Aster, Azalea, Begonia, Calendula, Chrysanthemums, Coleus, Fuchsia, Geraniums, Iv/, Marigold, Salvia, Petunia, Poinsettia, and Roses.

8660-83 3 of 4

CONTACT SPRAY: For Whiteflies, apply Vertagreen White Fly Spray at 1 pint per 100 gallons of water or 1 teaspoonful per gallon. Spray upper and lower surfaces of plant foliage to point of runoff. Spray in late afternoon and at night only, when the temperature ranges from 50°F to 72°F. Repeat application weekly for four weeks or longer for control of adults, pupae, and nymphs. Complete spray schedule on Poinsettia before blooms form. Where fewer applications are preferred, spray at the rate of 1 quart per 100 gals. of water or 2 tsp. per gallon. Repeat as necessary. Apply sprays as mentioned above.

NON-THERMAL APPLICATORS: Close off greenhouse area to be treated for White Fly infestations. Direct mist first toward upper corners of the greenhouse, then apply evenly throughout the area. Keep nozzle at least three feet from plant foliage. Apply Vertagreen White Fly spray at the rate of 1-1/2 fl. oz. thoroughly mixed in 3 fl. oz. of methylene chloride per 10,000 cu. ft. This is equivalent to 1 gram of active ingredient per 1,000 cu. ft. Keep treated area closed off for at least two hours. Do not remain in treated areas. Ventilate the area after treatment is completed. Apply at weekly intervals for at least four weeks. Applications should be made during the late afternoon or at night when the temperature range is 50°F to 72°F.

#### STORAGE AND DISPOSAL

#### FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by scorage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

### FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration. or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

8660-83 4 of 4

### WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured by:

PARKER FERTILIZER CO. 201 4TH STREET SYLACAUGA, AL 35150

Date Printed: 07/16/92

Date Approved: Supersedes: Label Change: