PM 25

}

8660-99

3660-99 1 of 4

# (FRONT PANEL)

# **GREEN UP**

# PROTECK-PLUS

### CONTAINS BALAN"

# PREEMERGENT HERBICIDE + 18-3-5 FERTILIZER

For Use On Ornamental Lawns And Turf. Will Control Crabgrass, Fo<sup>--</sup>ail, Watergrass, Annual Bluegrass (Poa annua), and other grasses as listed

FC? USE ON THE FOLLOWING TURF

Bluegrass (perennial)	Bermudagrass						
Ryegrass (perennial)	Centipedegmass						
Fescue (except fine-leaved)	St. Augustinegrass						
Bahiagrass	Zoysiagrass						

# ACTIVE INGREDIENTS:

N-butyl	-N-ethyl-a	a,a	,a,	-t:	rif	E11	101	co-	-2	,6	-di	ini	Ĺtz	co-	- P-	to	<b>)l</b> u	uid	lin	e	•	٠	•	-	•	. 78%
INERT INGR	EDIENTS:						-	-		-			-	-			•			•	•	٠	•			<u>99.22%</u>

Total 100.00%

Balan" is the trademark for Benefin, DowElanco.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT	OF	PRACTICAL	TREATMENT
-----------	----	-----------	-----------

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists. IF IN EYES: Flush with water for 15 minutes.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 29 LBS.


COVERS UP TO 5,000 SQ. FT. OF TURF

EPA Reg. No. 8660-99

EPA Est. 9193-OH-1

. .

المحاد المحاد المحاد المحاد



3.1

8660-99 2 of 4

## (BACK PANEL)

# NET WEIGHT 29 LBS.

# COVERS UP TO 5,000 SQ. FT. OF TURF

# GREEN UP

# PROTECK-PLUS

#### CONTAINS BALAN

# PREEMERGENT HERBICIDE + 18-3-5 FERTILIZER

#### PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, ponds, and streams. Do not apply when weather conditions favor run off or drift from target area.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

ProTeck-Plus is a preemergence herbicide and lawn food which should be applied directly to established turf for the control of Smooth Crabgrass, Hairy Crabgrass, Goosegrass (Silver Crabgrass), Watergrass (Barnyardgrass), Yeilow and Green Foxtail, and Annual Bluegrass (Poa annua).

It controls annual grass weeds by killing their seeds as they germinate. It does not control established weeds. When it is applied as directed the following established grasses are tolerant to Balan: Perennial Bluegrasses, Perennial Ryegrass, Centipedegrass, Fescue, Zoysiagrass, Bermudagrass, St. Augustinegrass and Bahiagrass.

Apply to turf before the germination of annual grass weeds. Do not reseed within 42 days after application.



8660-99 3 of 4

#### GUARANTEED ANALYSIS

Total Nitrogen (N)	18%
Available Phosphoric Acid $(P_2O_5)$	3%
Soluble Potash (K <sub>2</sub> O)	5 <b>%</b>

Sources of Plant Nutrients: Urea, Diammonium Phosphate, and Muriate of Potash

When applied to 5,000 sq. ft. it supplies 1.04 lbs. Nitrogen per 1,000 sq. ft. Supplies 1.58 lbs. "N" per 1,000 sq. ft. on 3,300 sq. ft. application.

#### CRABGRASS, GOOSEGRASS, FOXTAIL:

Northern Cool Season Grass - Apply bag contents to 5,000 sq. ft. area.

Southern Warm Season Grass - Apply bag contents to 3,300 sq. ft. area. Note: Apply in the spring before Crabgrass germinates (mid-April, south to early May, north). Apply evenly with a drop or rotary spreader.

#### ANNUAL BLUEGRASS (Poa annua):

Northern areas - Apply contents to 5,000 sq. ft. of turf. This provides 2 lbs. of active Balan per acre. Treat in late summer or early fall before seed germinates.

Southern Areas - On grasses such as Bermuda, St. Augustine, and Bahia, apply contents to 3,0-3 sq. ft. This provides 3 lbs. of active Balan per acre. Treat before Poa annua germinates in early fall.

#### SUGGESTED SYREADER SETTINGS

Make sure of your spreader's calibration. Use a drop or cyclone type.

	NORTHERN	SOUTHERN
	<u>5.000 Sq. Ft.</u>	<u>3.300 Sq. Ft.</u>
Scotts	5	7
Cyclone	3 1/2	5

### PRECAUTIONS

This product may thin fine-leaved Fescues at rates above 60 lbs. per acre (1.5 lbs. active ingredient). It should not be applied in the spring to Bluegrass planted the previous fall. Do not apply to Bentgrass putting greens or Dicnondra turf. Do not apply to newly sprigged areas of Bermudagrass, St. Augustinggrass, or Centipedegrass until these turfs are well established.



8660-99 4 of 4 4 % 4

# STORAGE AND DISPOSAL

FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE** DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs, and domestic water supplies. **PESTICIDE DISPOSAL**: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Concrol Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

### WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS S'LD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXFRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

Manufactured By:

PARKER PERTILIZER CO. 201 4TH STREET SYLACAUGA, AL 35150

Date Printed: 05/07/92 Date Approved: Supersedes: Label Change:

TUEST ATE-TABLE COPY