

P# 13 8111-1-15-94

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

NOV 16

Mr. Sanford R. Simon
 The Andersons
 201 West Fourth Street
 Sylacauga, AL 35150

Dear Mr. Simon:

Subject: Amendment- Label Revisions and Confidential Statement of Formulas
 Vertagreen 5% Diazinon Insecticide
 EPA Registration No. 8660-101
 Your Letter Dated July 8, 1994

This amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable, provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. On the front panel change "Ants Army worms" to "Ants, Armyworms".
 - b. Under DIRECTIONS FOR USE change "federal law" to "Federal law".
 - c. Under Lawn Spot Treatment change "55-80 degrees" to "65-80 degrees".

d. Under Band Treatment ~~change~~ ^{change} ~~to~~ ^{to} ~~5' by 20'~~ ^{5' by 20'} to ~~(2' by 50')~~ ^(2' by 50').

SYMBOL	412505						
SURNAME	KAG						
DATE	11/25/94						

e. Under **PRECAUTIONARY STATEMENTS** delete "May cause contact sensitization...contact a physician.

Your Confidential Statement of Formula dated July 8, 1994 has been reviewed and is acceptable.

3. Submit three (3) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager (13)
Insecticide-Rodenticide Branch
Registration Division (H7505C)

Enclosure

front panel

ACCEPTED
with COMMENTS
in EPA Letter Dated

NOV 16 1994

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

8660-101

VERTAGREEN®

5% DIAZINON INSECTICIDE

LAWN AND GARDEN INSECT CONTROL

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and in the home garden only.

Ready-to-use granules. Contains 5 % diazinon insecticide.

Treats up to 5,000 square feet.

Vertagreen 5% Diazinon Insecticide Kills: Ants Army worms, Bermudagrass mites, Billbugs, Brown dog ticks, Chiggers, Chinch bugs Crickets, Cutworms, Earwigs, Fire ants* Fleas, Onion maggots, Sod webworms, White grubs of:

- European chafer ● Japanese beetle ● Southern chafer, Wire worms, and others.

* Aids in control.

See Directions for Use

Active Ingredient:

Diazinon®: O,O-Diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)	
phosphorothioate	5.00 %
Inert Ingredients:	95.00 %
Total:	100.00 %

EPA Registration Number: 8660-101


EPA Establishment Number: 9198-OH-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements on back of bag.

NET WEIGHT 10 lbs. (4.5 kg)



back panel

In Lawns Vertagreen 5% Diazinon Insecticide kills:

Anis	Cutworms	Sowbugs (Pillbugs)
Armyworms	Digger wasps	Springtails
Bermudagrass mites	Earwigs	White grubs of:
Billbugs	Fire ants*	• European chafer
Brown dog ticks	Fleas	• Japanese beetle
Chiggers	Lawn moths	• Southern chafer
Chinch bugs	Leafhoppers	
Clover mites	Millipedes	
Crickets	Sod webworms	

*Aids in control

In Gardens Vertagreen 5% Diazinon Insecticide kills:

Cutworms	Onion maggots	Wireworms
Mole crickets	Root maggots	

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use and the Conditions of Sale and Warranty before using this product. If terms are not acceptable, return the unopened product container at once.

CONDITIONS OF SALE AND WARRANTY

The Directions for Use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness, or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Pursell Industries, Inc. or the Seller. All such risks shall be assumed by the Buyer.

Pursell Industries, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use subject to the inherent risks referred to above. **Pursell Industries, Inc. makes no other express or implied warranty of Fitness or Merchantability or consequential, special, or indirect damages resulting from the use or handling of this product.** Pursell Industries, Inc. and the Seller offer this product, and the Buyer and user accept it, subject to the foregoing Conditions of Sale and Warranty, which may be varied only by agreement in writing signed by a duly authorized representative of Pursell Industries, Inc.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

OVERALL LAWN APPLICATION

Special Precautions To Reduce Risk To Birds

Birds, especially waterfowl (ducks or geese), feeding or drinking on treated areas may be killed. To reduce risks to birds:

1. Do not treat areas larger than 15,000 sq. ft.
2. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20.
3. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl.
4. Avoid overlapping treated areas.
5. Follow all watering instructions in the directions for use, however, stop watering before puddling occurs.
6. Do not apply this product to sites occupied by waterfowl or to sites within 75 ft. of any body of water that may attract waterfowl, such as lakes or ponds.
7. If it is necessary to repeat application, wait a minimum of 3 weeks between applications. Make no more than 4 applications per year.

BEST AVAILABLE COPY

Vertagreen 5% Diazinon Insecticide controls certain lawn insects on both grass and dichondra lawns. Apply with fertilizer spreader. Refer to the tables that follow for use rates and sprayer settings.

Pest	Rate*	Timing/Special Directions
Ants, Armyworms, Bermudagrass mites, Brown dog ticks, Chiggers, Chinch bugs, Clover mites, Crickets, Cutworms, Earwigs, Fleas, Lawn moths (sod webworms), Leafhoppers, Millipedes, Sowbugs (Pillbugs), Springtails (Collembola)	2 lbs. per 1,000 sq. ft.	Apply when grass is dry. Water lawn lightly after application. If necessary, repeat application 3- week intervals. Make no more than 4 applications per year.
Billbugs, White grubs of Japanese beetle, European chafer, Southern chafer, Ateneus spretulus (dung beetle)	2 lbs. per 1,000 sq. ft.	Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. For billbug control, apply when activity is first observed or when chewed grass indicates damage. If necessary, repeat application at 3-week intervals. Make no more than 4 applications per year. For grub control, except dung beetle, treat any time between late July and early October. For dung beetle grub control, apply in Spring or Summer when damage is first noticed. Water grass with at least 1/4 inch of water application, however, stop watering before puddling occurs. Treat again in late Summer or early Fall, if a second generation occurs.
Hyperodes weevils	2 lbs. per 1,000 sq. ft.	Water grass with at least 1/4 inch of water after application, however, stop watering before puddling occurs. Treat problem areas in mid-April and again in mid-May.

BEST AVAILABLE COPY

* 1,000 sq. ft. = an area 20 ft. x 50 ft.

100 sq. ft. = an area 10 ft. x 10 ft.

2 lbs. per 1,000 sq. ft. = 3.2 oz. per 100 sq. ft.

NOTE: The spreader settings listed below are approximate. Accurately weigh a few pounds into your spreader, and apply to a measured area. Then adjust the spreader setting, if necessary, in order to ensure correct rate of application.

LAWN SPREADER SETTINGS NO.**

(To deliver 2 lbs. per 1,000 sq. ft.)

Republic EZ Drop	True Temper Rotary	Republic EZ Rotary	Scotts Easy Green
Ortho Drop (5000)	Scotts Speedy Green	Ortho Rotary (6000)	Precision Rotary (SB 50)
Scotts Drop (PF3)	Republic EZ Grow	Ev-N-Spred Rotary (2600)	Quaker Rotary (10500)

**For application with other types of equipment, apply at the rate of 2 lbs. per 1,000 sq. ft.

Lawn Spot Treatment

Ants: Apply 1½ teaspoons of Vertagreen 5% Diazinon Insecticide over and around each ant hill.

Fire ants: To aid in the control of fire ants in lawns and other outdoor home recreation areas, apply ½ cup over, and 2 ft. around, each fire ant mound. Immediately after application, water-in thoroughly. Apply gently to avoid disturbing ants. Use of high pressure watering equipment may disturb the ants and cause migration, reducing product effectiveness. A minimum of 1 gal. of water should be applied to each mound area. Reapply as necessary.

Note: For best results, apply in cool weather, 55-80°F, or in early morning or late evening hours. Works within 24 hours. No waiting for days or weeks. Treat new mounds as they appear.

Digger Wasps: Apply 1½ teaspoons of Vertagreen 5% Diazinon Insecticide in and around each nest opening. Apply at dusk when wasps are not active.

Band Treatment Around House Foundation

The following pests are commonly found around or near house foundations, and may enter the house:

Ants	Crickets	Millipedes
Brown dog ticks	Earwigs	Sowbugs (Pillbugs)
Clover mites	Fleas	Springtails (Collembola)

Apply a 2-foot band of Vertagreen 5% Diazinon Insecticide next to the house foundation. Distribute granules uniformly, using a rate of 3.2 oz. for each 100 sq. ft. of area (5' by 20') treated. If necessary, repeat application at 3-week intervals.

VEGETABLES - INSECT CONTROL

Special Precautions To Reduce Risk To Birds

Birds feeding on treated areas may be killed. To reduce risk to birds:

1. Thoroughly mix Vertagreen 5% Diazinon Insecticide into the soil during or immediately after application. Refer to the Soil incorporation table that follows.
2. Granules that are spilled or are visible on soil surfaces should be covered with soil or thoroughly mixed into the soil.

Vertagreen 5% Diazinon Insecticide controls certain garden insects on crops listed in the following tables. Apply uniformly as directed for each crop. When a range of rates is given, use the higher rate only for heavier infestations.

Do not apply this product to food crops grown in greenhouses.

Unless direction under Timing, Special Directions state otherwise, applications for control of pests listed below should be thoroughly mixed into the soil during or immediately after application.

BEST AVAILABLE COPY

SOIL INCORPORATION

INSECT	Depth of Mixing or Incorporation into Soil
Surface cutworms*	2-3 inches
Subterranean cutworms*	3-6 inches
Mole Crickets	1-2 inches
Root maggots	3-4 inches
Wireworms	4-8 inches

*If type of cutworm cannot be identified, mix into soil 4-6 inches deep.

Crop	Pest	Rate	Timing/Special Directions
Beans (Lima, Pole and Snap) (succulent only)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Note: Do not use on dried bean varieties, such as pintos, dried limas, or navy beans.			
Beets, Red (Table)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Broccoli	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Brussels Sprouts	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.

BEST AVAILABLE COPY

CROP	PEST	RATE	Timing/Special Directions
Cabbage	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Carrots	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Cauliflower	Root Maggots	7-10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting or transplanting.
	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Celery	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Collards	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Cucumbers	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Endive (Escarole)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Kale	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Lettuce	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Melons (Cantaloupes, Casabas, Crenshaws, Honeydew, Muskmelons, Persians, and Hybrids of these), and Watermelons	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Mustard	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Onions (Bulb and Green)	Onion Maggots	7-14 oz. per 500 sq. ft.	Broadcast just before planting and mix into the top 3-4 inches of soil. Note: Resistant onion maggots will not be controlled by diazinon or similar type insecticides.
	Wireworms	10½-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Parsley	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Peas (succulent only)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Note: Do not use on dried pea varieties such as dried blackeyed peas.			
Peppers	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Potatoes	Cutworms (Surface and Subterranean)	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Radishes	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Spinach	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	
Squash (Summer and Winter)	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½-14 oz. per 500 sq. ft.	

BEST AVAILABLE COPY

Crop	Pest	Rate	Timing/Special Directions
Sweet Corn	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½ -14 oz. per 500 sq. ft.	
	Seed Corn Maggots	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil 2 inches just before planting.
Sweet Potatoes	Wireworms, Flea Beetles	10½ oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
Swiss Chard	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½ -14 oz. per 500 sq. ft.	
Tomatoes	Cutworms	7-14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.
	Wireworms	10½ -14 oz. per 500 sq. ft.	
	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
Turnips	Mole Crickets	3½ oz. per 500 sq. ft.	Broadcast and mix into soil 1-2 days before planting.
	Wireworms	10½ -14 oz. per 500 sq. ft.	Broadcast and mix into soil just before planting.

STORAGE AND DISPOSAL

STORAGE: Store in original container in a cool, dry, well-ventilated area. Store product in an area inaccessible to children and domestic animals. Do not store near fertilizers, herbicides, other insecticides, seeds, food, or feed.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes. In the event of a major spill, fire or other emergency, call 1-800-874-8892.

CAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating or smoking. May cause contact sensitization following repeated contact with skin of susceptible individuals. If sensitization reactions result, contact a physician. Do not store near food or feed products. Food utensils such as spoons or measuring cups must not be used for food purposes after use in measuring pesticides. Do not contaminate ornamental fish ponds. Do not allow children or pets on treated areas until granules have been watered into the soil and the grass or soil is dry.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed: Call a physician immediately. Drink one or two glasses of water and induce vomiting by touching the back of the throat with finger. Repeat until vomit is clear. Do not induce vomiting or give anything by mouth if a person is unconscious or convulsing.

If Inhaled: Remove victim to fresh air and apply artificial respiration, if indicated.

If On Skin: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

If In Eyes: Rinse eyes with plenty of water and call a physician immediately.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl (ducks or geese), feeding or drinking on treated areas may

be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increases potential hazards to birds. Avoid overlapping of granules. Where irrigation (watering) is recommended in the directions, or if waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after applying this product, however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring area. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

Manufactured by:

PURSELL INDUSTRIES, INC.
201 WEST FOURTH STREET
SYLACAUGA, AL 35150