UNITED STATES ENVIRONMENTAL PROTECTION AGENCY PM 23 8660-104

18 JUN 1992

Mr. Ganiel Paradiso, Jr. d\b\s Parker Fertilizer 201 West Forth Street Sylacauba, AL. 35150

Dear Mr. Paraciso:

Subject: Vertagreen Infram 75 EFA Registration Wolfbour - Chickween Siller EFA Registration No. 80c0-24 Vertagreen Keed And Feed EFA Registration No. 8060-14 Proteck - Way EFA Registration No. 8060-104 Verta-Green St. Augusting ween & Feed EFA Registration No. 8060-14 Perta-Green St. Augusting ween & Feed EFA Registration No. 8000-147 Ref Label Abenquent Your Submission Dated Day 10, 1092

The labeling referred to above submitted in connection with registration under the Federal insectionde fungicide and Rodenticide act, as amended is acceptable, provided that under the Environmental Hazard Section of the label replace the statement" Do not apply directly to water or wetlands" to read "Bo not apply directly to water, to areas where surface water is present or to intertidal areas below the mean migh water mark." Please submit five (5) copies of your final printed labels incorporating this dmendment. A stamped copy of labeling is enclosed for your records.

Sincerely

## BEST AVAILABLE COPY

Joanne I. Miller Product Hanager (23) Fungicide-Herbicide Branch Registration Division (H+7505C)

CONCURRENCES												
SYMBOL	Facil	beuro						<u> </u>				
DATE	••••••	*****	*******	• • • • • • • • • • • • • • • • • • • •	•••••••••••••••••••••••••••••••••••••••	••••••						
EPA Form	1320-1 (12-70)	L		I		<u></u>	L	L				

OFFICIAL FILE COPY

pape 195

.s. GPO 1985-467-853

100.00%

8660-104 1 of 4

(FRONT PANEL)

#### VERTAGREEN

#### PROTECK 3-WAY

#### CONTAINS PREEMERGENT HERBICIDE + 18-3-5 FERTILIZER + BROADLEAF WEED KILLER

For Lawns, Fairways and Ornamental Turf

-Pre-Emergence Control (Balan®): Crabgrass, Foxtail, Watergrass, Annual Bluegrass

-Post-Emergence Control: Dandelions, Chickweed, Henbit, Knotweed, Spurge and other weeds as listed.

-Feeds lawn with 18-3-5 analysis.

Use on Bluegrass, Perennial Rye, Fescue, Bermuda and Zoysiagrass.

#### ACTIVE INGREDIENTS.

N-butyl-N-ethyl-a,a,a,-trifluoro-2,6-dinitro-p-toluidine	1.15%
*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid	1.50%
**Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic acid)	0.28%
INERT INGREDIENTS:	<u>97.07%</u>

Equivalent to: \*1.24% of 2,4-Dichlorophenoxyacetic Acid \*\*0.23% of 3,6-dichloro-o-anisic acid

Balan<sup>™</sup> is the trademark for Benefin, DowElanco.

KREP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF IN EYES: Flush with water for 15 minutes.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 20 LBS.

#### COVERS UP TO 5,000 SQ. FT. OF TURF

EPA Reg. Nc. 8660-104

EPA Est. 9198-OH-1

# BEST AVAILABLE COPY

Total

Sec. 1. 1. 18

ere Storage

395

8660-104 2 of 4

(BACK PANEL)

#### NET WEIGHT 20 LBS.

COVERS UP TO 5,000 SQ. FT. OF TURF

#### VERTAGREEN®

PROTECK 3-WAY

#### CONTAINS

#### PREEMERGENT HERBICIDE + 18-3-5 FERTILIZER + BROADLEAF WEED KILLER

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.

This product can reach groundwater from imprope handling. To minimize contamination from spills during loading and cleaning of equipment, take the following steps:

<u>Handling</u>: When handling this product, wear chemical resistant gloves. Wash nondisposable gloves thoroughly with soap and water before removing. If spills occur, collect the material and dispose of by following disposal instructions on this label.

<u>Cleaning of Equipment</u>: When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

8660-104 3 of 4

4 03

5

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Super Proteck is a pre-emergence herbicide, lawn food and a broadleaf weed killer. It should be applied to established turf for the control of: Grasses (Smooth Crabgrass, Hairy Crabgrass, Goosegrass, Watergrass, Yellow and Green Foxtail and Annual Bluegrass) as well as broadleaf weeds: Buckhorn, Burdock, Bur Clover, Chickory, Common Chickweed, Mouse Ear Chickweed, Clover (White), Creeping Charley (Ground Ivy), Curly Dock, Dandelion, Dog Fennel, Frenchweed, Heal-All, Henbit, Knawel, Knotweed, Lambsquarter, Mallow, Morning Glory, Mustard, Oxalis, Peppergrass, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Red Sorrel, Sheep Sorrel, Sow Thistle, Spotted Spurge, and Yarrow.

#### WHEN TO APPLY

For preemergence action, apply this product prior to the germination of weed seeds in spring or early summer (April 1st thru May 1st; or later if the preemergence is being applied in two separate applications; one early and the other the last of May.) Broadleaf weeds present at the time of application will be controlled. Apply when there is adequate soil moisture and the turf (weeds) are growing actively. Do not apply to newly seeded lawns. For Annual Bluegrass, treat in late summer or early fall before Poa annua seed germinates.

HOW TO APPLY Apply 20 lbs. to 5,000 sq. ft. of turf. This rate feeds the lawn with .72 lbs. Nitrogen per 1,000 sq. ft.

#### PRECAUTIONS

Do not use on Centipedegrass, St. Augustinegrass, or Dichondra; nor on lawns or turf where desirable Bentgrass or clovers are present. Do not allow dust to come in contact with shrubs, flowers, trees, vegetables, or other valuable plants. Avoid application to areas under the branches or drip line of trees and shrubs. Do not reseed within 42 days after application.

This product may thin fine-leaved Fescues at rates above 60 lbs. per acre (1.5 lbs. active ingredient).

Do not apply to Bentgrass putting greens or Dichondra turf. Do not apply to newly sprigged areas of Bermudagrass.

#### SUGGESTED SPREADER SETTINGS

 Scotts
 5
 1/2

 Cyclone
 3
 1/2

NOTE: The above suggested spreader settings are only approximate. Age and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application. Close spreader when filling, stopping, or turning. DO NOT KAND SPREAD.

### BEST AVAILABLE COPY

8660-104 4 of 4

595

Avoid getting on concrete, stucco or asbestos siding, as the composition of certain fertilizer elements can cause permanent stain.

#### STORAGE AND DISPOSAL

#### FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

#### FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

#### WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

> The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

> > Manufactured by:

PARKER FERTILIZER CO. 201 4TH STREET SYLACAUGA, AL 35150

Date Printed: 05/06/92 Date Approved: Supersedes: Label Change:

BEST AVAILABLE COPY

;

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY 8660-104 PJ 125

18 JUN 1992

Mr. Daniel Paraciso, ur. GNDNa Parker Fertilizer 201 West Forth Street Sylacauba AL, US150

Lear Mr. Faradiso:

Ł

Subjects Switzgreen Thison (E Fra Registration Rollted-in Scrtagard RCPP Clover - Chickbeen Differ CPA Registration No. 8000-24 vertaurec: Reed And Feen CPA Registration No. 8000-14 Protect o Bay CPA Registration No. 8000-14 Frotect o Bay CPA Registration No. 8000-14 Certo-Creen St. Augusting Reed Steve CFP Registration No. 8000-14 Her Cobel Amenguent Your Submission Dated Nay 22, 1592

The iduality referred to adove submitted in connection with registration under the Federal insections of Fungicide and controlice act as anended is acceptible provided that under the Environmental Hazard Section of the label replace the statement" Do not apply directly to water or wetlands" to read "So not apply directly to water, to areas where surface water is present or to intertigal areas below the mean migh water mark." Please submit five is copies of your final printed labels incorporating this amendment. A stamped copy of labeling is enclosed for your records.

Sincercly

Joanne I. Hiller Product Hanager (23) Fungloide-Herbicige Branch Registration Division (H-7505C)

	CON	CURRENCES		
MOOL				
THANE D		*****	• • • • • • • • • • • • • • • • • • • •	
		*****	·····	• • • • • • • • • • • • • • • • • • • •
1320-1 (12-70)	<b>BESTA</b>	VAILABL	ECOPY	OFFICIAL FILE COPY

8660-104 1 of 4 285

#### (FRONT PANEL)

#### VERTAGREEN®

#### PROTECK 3-WAY

#### CONTAINS PREEMERGENT HERBICIDE + 18-3-5 FERTILIZER + BROADLEAF WEED KILLER

For Lawns, Fairways and Ornamental Turf

-Pre-Emergence Control (Balan®): Crabgrass, Foxtail, Watergrass, Annual Bluegrass

-Post-Emergence Control: Dandelions, Chickweed, Henbit, Knotweed, Spurge and other weeds as listed.

-Feeds lawn with 18-3.5 analysis.

Use on Bluegrass, Perennial Rye, Fescue, Bermuda and Zoysiagrass.

#### ACTIVE INGREDIENTS:

4 <sup>- 1</sup>

-

N-butyl-N-ethria, a, a, fluoro-2,6-dinitro-p-toluidine	1.15%
*Dimethylamine Salt of 2,4-Dicidorophenoxyacetic Acid	1.50%
**Dimethylamine Salt of Dicamba (3,dichloro-o-anisic acid)	0.28%
INERT INGREDIENTS:	<u>97.07%</u>

Equivalent to: \*1.24% of 2,4-Dichlorophenoxyacetic Acid \*\*0.23% of 3,6-dichloro-o-anisic acid

Balan<sup>w</sup> is the trademark for Benefin, DowElanco.

KEEP OUT OF REACH OF CHJ LDREN

CAUTION

## STATEMENT OF PRACTIC L TREATMENT

IF ON SKL': Wash with plenty of \* ap and water. Get medical attention if irritation presists. IF IN EYES: Flush with water. for 15 minutes.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

NET WEIGHT 20 LBS.

#### COVERS UP TO 5,000 SQ. FT. OF TURF

EPA Reg. No. 8660-104

EPA Est. 9198-0H-1

100.00%

Jotal

. . . **.** . .

يعدده وحد المرد

385

8660-104 2 of 4

#### (BACK PANEL)

#### NET WEIGHT 20 LBS.

COVERS UP TO 5,000 SQ. FT. OF TURF

#### VERTAGREEN®

#### PROTECK 3-WAY

#### CONTAINS

)

#### PREEMERGENT HERBICIDE + 18-3-5 FERTILIZER + BROADLEAF WEED KILLER

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters.

This product can reach groundwater from improper handling. To minimize contamination from spills during loading and cleaning of equipment, take the following steps:

Handling: When handling this product, wear chemical resistant gloves Wash nondisposable gloves thoroughly with soap and water before removing. If spills occur, collect the material and dispose of by following disposal instructions on this label.

<u>Cleaning of Equipment</u>: When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

8660-104 3 of 4

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Super Proteck is a pre-emergence herbicide, lawn food and a broadleaf weed killer. It should be applied to established turf for the control of: Grasses (Smooth Crabgrass, Hairy Crabgrass, Goosegrass, Watergrass, Yellow and Green Foxtail and Annual Bluegrass) as well as broadleaf weeds: Buckhorn, Burdock, Bur Clover, Chickory, Common Chickweed, Mouse Ear Chickweed, Clover (White), Creeping Charley (Ground Ivy), Curly Dock, Dandelion, Dog Fennel, Frenchweed, Heal-All, Henbit, Knawel, Knotweed, Lambsquarter, Mallow, Morning Glory, Mustard, Oxalis, Peppergrass, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Red Sorrel, Sheep Sorrel, Sov Thistle, Spotted Spurge, and Yarrow.

#### WHEN TO APPLY

 $\mathbf{b}$ 

For preemergence action, apply this product prior to the germination of weed seeds in spring or early summer (April 1st thru May 1st; or later if the preemergence is being applied in two separate applications; one early and the other the last of May.) Broadleaf weeds present at the time of application will be controlled. Apply when there is adequate soil moisture and the turf (weeds) are growing actively. Do not apply to newly seeded lawns. For Annual Bluegrass, treat in late summer or early fall before Poa annua seed germinates.

HOW TO APPLY Apply 20 lbs. to 5,000 sq. ft. of turf. This rate feeds the lawn with .72 lbs. Nitrogen per 1,000 sq. ft.

#### PRECAUTIONS

Do not use on Centiredegrass, St. Augustinegrass, or Dichondra; nor on lawns or turf where desirable Bentgrass or clovers are present. Do not allow dust to come in contact with shrubs, flowers, trees, vegetables, or other valuable plants. Avoid application to areas under the branches or drip line of trees and shrubs. Do not reseed within 42 days after application.

This product may thin fine-leaved Fescues at rates above 60 lbs. per acre (1.5 lbs. active ingredient).

Do not apply to Bentgrass putting greens or Dichondra turf. Do not apply to newly sprigged areas of Bermudagrass.

#### SUGGESTED SPREADER SETTINGS

Scotts .	•	•	•	•	•	•	•	•	•	•	•	•	٠	•	•	•	•	•	•	•	5 1/2
Cyclone		٠	•	٠	٠	٠	•	٠	•	•	•	•	•	•	•	٠	•	•	•	٠	3 1/2

NOTE: The above suggested spreader settings are only approximate. Age. and condition of spreader can cause wide variation. Be sure to calibrate your spreader with each application. Close spreader when filling, stopping, or turning. DO NOT HAND SPREAD.

8660-104 4 cf 4

Avoid getting on concrete, stucco or asbestos siding, as the composition of certain fertilizer elements can cause permanent stain.

#### STORAGE AND DISPOSAL

#### FOR HOMEOWNER USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

#### FOR PROFESSIONAL/COMMERCIAL USE ONLY:

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL**: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate, is a violation of Federal Law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instruction, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance. **CONTAINER DISPOSAL**: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State or local authorities, by burning. If burned, stay out of smoke.

#### WARRANTY STATEMENT

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions for Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

> The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer. Buyer assumes all risks of use, storage or handling of this product.

> > Hamufactured by:

PARKER FERTILIZER CO. 201 4TH STREET SYLACADCA, AL 35150

Date Printed: 05/06/92 Date Approved: Supersedes: Label Change: